

REAL GROUPS TRANSITIVE ON COMPLEX FLAG MANIFOLDS

JOSEPH A. WOLF

(Communicated by Rebecca A. Herb)

ABSTRACT. Let $Z = G/Q$ be a complex flag manifold. The compact real form G_u of G is transitive on Z . If G_0 is a noncompact real form, such transitivity is rare but occasionally happens. Here we work out a complete list of Lie subgroups of G transitive on Z and pick out the cases that are noncompact real forms of G .

0. THE PROBLEM

Let $Z = G/Q$ be a complex flag manifold where G is a complex connected semisimple Lie group and Q is a parabolic subgroup. Let G_0 be a real form of G . If G_0 is the compact real form, then it is transitive on Z . On a number of occasions the question has come up as to whether any noncompact real form of G can be transitive on Z . Here I'll record the answer. The rough answer is "yes, but just a few." The precise answer, Corollaries 1.7 and 2.3 below, follows from a more general classification, Theorems 1.6 and 2.2. This more general classification uses a technique of D. Montgomery [M], together with some results of [W1] that depend in an essential way on a classification [O1] of A. L. Onishchik.

After this paper was written I learned of Onishchik's book [O2]. There is some overlap for compact groups, but there are no inclusions.

1. THE SOLUTION FOR IRREDUCIBLE FLAGS

We formulate the problem in terms of transitive subgroups. Let G_u be the compact real form of G , so $Z = G_u/(G_u \cap Q)$ and $G_u \cap Q$ is the compact real form of the reductive part of Q . Let $A \subset G$ be a closed subgroup that is transitive on Z . The identity component A^0 of A is transitive on Z , because Z is connected, so a maximal compact subgroup $B^0 \subset A^0$ already is transitive on Z , according to Montgomery [M]. We may replace A by a conjugate and assume $B = A \cap G_u$. So

Received by the editors July 28, 1999 and, in revised form, December 9, 1999.

2000 *Mathematics Subject Classification*. Primary 22E15; Secondary 22E10, 32E30, 32M10.

Key words and phrases. Semisimple Lie group, semisimple Lie algebra, representation, flag manifold, flag domain.

The author's research was supported by the Alexander von Humboldt Foundation and by NSF Grant DMS 97-05709. The author thanks the Ruhr-Universität Bochum for hospitality.

now we have several expressions:

$$(1.1) \quad \begin{aligned} Z &= G/Q = G_u/(G_u \cap Q) = A/(A \cap Q) = B/(B \cap Q) \\ &= A^0/(A^0 \cap Q) = B^0/(B^0 \cap Q). \end{aligned}$$

According to [W1, Prop. 3.1] there are just a few possibilities for a homogeneous almost-hermitian manifold Z to have distinct expressions such as G_u/L_u and $B^0/(B^0 \cap L_u)$, where G_u is the identity component of the group of all almost-hermitian isometries, G_u is simple, L_u is the centralizer of a torus subgroup of G_u , and $B^0 \subsetneq G_u$ with B^0 connected. They are :

$$(1.2) \quad Z = P^{2n-1}(\mathbb{C}) = SU(2n)/U(2n-1) = Sp(n)/(Sp(n-1) \cdot U(1)), \text{ complex projective space,}$$

$$(1.3) \quad Z = SO(2r+2)/U(r+1) = SO(2r+1)/U(r), \text{ unitary structures on } \mathbb{R}^{2r+2},$$

$$(1.4) \quad Z = SO(7)/(SO(5) \cdot SO(2)) = G_2/U(2), \text{ 5-dimensional complex quadric, and}$$

$$(1.5) \quad Z = SO(8)/(SO(6) \cdot SO(2)) = \{Spin(7)/Z_2\}/U(3), \text{ 6-dimensional complex quadric.}$$

This applies in our situation because $L_u = G_u \cap Q$ is the centralizer of a torus subgroup of G_u , and Z has a G_u -invariant hermitian metric.

Now return to the expression $Z = G/Q$. G (and thus G_u) is simple. Let $A \subsetneq G$ be a closed subgroup that is transitive on Z and let B be its maximal compact subgroup. We may assume $B = A \cap G_u$. Then $B \subsetneq G_u$, B^0 is transitive on Z , and the expression $Z = G_u/L_u = B^0/(B^0 \cap L_u)$ is given above. In each case the group B^0 is simple, so A^0 has Levi decomposition $A^0 = A_{ss}^0 A_{rad}^0$ into semisimple part and solvable radical, where B^0 is a maximal compact subgroup of A_{ss}^0 . We run through the 4 possibilities listed above.

For (1.2), $G = SL(2n; \mathbb{C})$ and $B^0 = Sp(n)$. The semisimple Lie groups with maximal compact subgroup $Sp(n)$ are $Sp(n), Sp(n; \mathbb{C})$, the quaternionic linear group $SL(n; \mathbb{H})$, and, for $n = 4$, the real group F_{4,C_4} . But F_4 does not have a representation of degree 8, in other words $F_4 \not\subset G$, so now A_{ss}^0 is one of $Sp(n), Sp(n; \mathbb{C})$ and $SL(n; \mathbb{H})$. Each of them is irreducible on \mathbb{C}^{2n} , so the unipotent radical of the algebraic hull of A^0 acts trivially on \mathbb{C}^{2n} and the center of the reductive part of A^0 acts by scalars. As G acts effectively and by transformations of determinant 1 on \mathbb{C}^{2n} now $A_{ss}^0 = A^0$, so A^0 is one of $Sp(n), Sp(n; \mathbb{C})$ and $SL(n; \mathbb{H})$. If $g \in G$ normalizes A^0 , then some element $g' \in gA^0$ centralizes A^0 , because A^0 has no rational outer automorphism. As A^0 is irreducible on \mathbb{C}^{2n} now g' is scalar (and thus acts trivially on Z). Thus $A = A^0 F$ where F can be any subgroup of the center $\{e^{2\pi i k/2n} I \mid 0 \leq k < 2n\}$ of G .

For (1.3), $G = SO(2r+2; \mathbb{C})$ and $B^0 = SO(2r+1)$. The semisimple Lie groups with maximal compact subgroup $SO(2r+1)$ are $SO(2r+1), SO(2r+1; \mathbb{C}), SO(1, 2r+1)$, and $SL(2r+1; \mathbb{R})$. But $A_{ss}^0 = SL(2r+1; \mathbb{R})$ would give $SL(2r+1; \mathbb{C}) \subset SO(2r+2; \mathbb{C})$, so the respective dimensions would satisfy $4r^2 + 4r \leq 2r^2 + 3r + 1$, forcing $r = 0$ and $Z = (\text{point})$. Thus¹ $A_{ss}^0 \neq SL(2r+1; \mathbb{R})$. Now A_{ss}^0 is one of $SO(2r+1), SO(2r+1; \mathbb{C})$, and $SO(1, 2r+1)$. The last one acts irreducibly on \mathbb{C}^{2r+2} , and there $A_{ss}^0 = A^0$ as above. For the first two, recall that $SO(2r+1)$ is absolutely irreducible on the tangent space $\mathfrak{so}(2r+2)/\mathfrak{so}(2r+1)$ of the sphere S^{2r+1} , so A_{rad}^0 has Lie algebra reduced to 0, and again $A_{ss}^0 = A^0$. Now A^0 is one of $SO(2r+1), SO(2r+1; \mathbb{C})$, and $SO(1, 2r+1)$. If $g \in G$ normalizes A^0 , then some

¹ The author thanks the referee for a comment that improved and clarified his treatment of this $SL(2r+1; \mathbb{R})$ case.

element $g' \in gA^0$ centralizes A^0 , because A^0 has no rational outer automorphism. Thus either $A = A^0$ or A/A^0 has order 2 where A is one of $O(2r + 1)$, $O(2r + 1; \mathbb{C})$, and $SO(1, 2r + 1) \cdot \{\pm I\}$.

For (1.4), $G = SO(7; \mathbb{C})$ and $B^0 = G_2$. The semisimple Lie groups with maximal compact subgroup G_2 are G_2 and its complexification $G_{2,\mathbb{C}}$. They are irreducible on \mathbb{C}^7 and have no rational outer automorphisms, so, as before, A^0 is either G_2 or $G_{2,\mathbb{C}}$, and if $g \in G$ normalizes A^0 , then some element $g' \in gA^0$ centralizes A^0 . This forces g' to be central in $SO(7; \mathbb{C})$, so $g' = 1$ and $A = A^0$. Thus A is either G_2 or $G_{2,\mathbb{C}}$.

Finally, (1.5) is obtained from the case $r = 3$ of (1.3) by applying the triality automorphism, so it does not give us anything more.

In summary,

Theorem 1.6. *Consider a complex flag manifold $Z = G/Q$. Suppose that Z is irreducible, i.e., that G is simple. Then the closed subgroups $A \subset G$ transitive on Z , $G_u \neq A \neq G$, are precisely those given as follows:*

1. $Z = SU(2n)/U(2n - 1) = P^{2n-1}(\mathbb{C})$ complex projective $(2n - 1)$ -space; $G = SL(2n; \mathbb{C})$ and $A = A^0F$ where A^0 is one of $Sp(n)$, $Sp(n; \mathbb{C})$ and $SL(n; \mathbb{H})$, and F is any subgroup of the center $\{e^{2\pi ik/2n} I \mid 0 \leq k < 2n\}$ of G . Here F acts trivially on Z , so A and A^0 have the same action on Z .

2. $Z = SO(2r + 2)/U(r + 1)$, unitary structures on \mathbb{R}^{2r+2} ; $G = SO(2r + 2; \mathbb{C})$ and $A = A^0F$ where A^0 is one of $SO(2r + 1)$, $SO(2r + 1; \mathbb{C})$, and $SO(1, 2r + 1)$, and where F is any subgroup of the center $\{\pm I\}$ of G . Here F acts trivially on Z , so A and A^0 have the same action on Z .

3. $Z = SO(7)/(SO(5) \cdot SO(2))$, 5-dimensional complex quadric; $G = SO(7; \mathbb{C})$ and A is either the compact connected group G_2 or its complexification $G_{2,\mathbb{C}}$.

Picking out the cases where A is a real form of G we have

Corollary 1.7. *Consider a complex flag manifold $Z = G/Q$. Suppose that Z is irreducible, i.e., that G is simple. Then the (connected) noncompact real forms $G_0 \subset G$ transitive on Z are precisely those given as follows:*

1. $Z = SU(2n)/U(2n - 1) = P^{2n-1}(\mathbb{C})$ complex projective $(2n - 1)$ -space; $G = SL(2n; \mathbb{C})$ and G_0 is the quaternion linear group $SL(n; \mathbb{H})$, which has maximal compact subgroup $Sp(n)$.

2. $Z = SO(2r + 2)/U(r + 1)$, unitary structures on \mathbb{R}^{2r+2} ; $G = SO(2r + 2; \mathbb{C})$ and G_0 is the Lorentz group $SO(1, 2r + 1)$, which has maximal compact subgroup $SO(2r + 1)$.

2. THE SOLUTION FOR FLAG MANIFOLDS IN GENERAL

We complete the solution of the problem by reducing it to the case where Z is irreducible.

Proposition 2.1. *Decompose $G = \prod G_i$, the local direct product of complex connected simple Lie groups. Thus $Z = \prod Z_i$, the product of irreducible flag manifolds $Z_i = G_i/Q_i$ where $Q_i = Q \cap G_i$. Then $A^0 = \prod A_i^0$ with $A_i^0 = A^0 \cap G_i$ and $B^0 = \prod B_i^0$ with $B_i^0 = B^0 \cap G_i$. The groups A_i^0 and B_i^0 are connected, simple, and transitive on Z_i .*

Proof. The solvable radical of A^0 is contained in a Borel subgroup of G , and thus has a fixed point on Z . It is normal in the transitive group A^0 so it fixes every point. Thus A^0 is semisimple. Similarly B^0 is semisimple.

Let $\pi_i : G \rightarrow G_i$ denote the projection. The compact connected group $\pi_i(B^0)$ is transitive on Z_i . So it must be the compact real form $G_{u,i} = G_i \cap G_u$ of G_i or one of the compact connected transitive groups described in (1.2), (1.3) or (1.4). (Recall that (1.5) is in fact a special case of (1.3).) In all cases, $\pi_i(B^0)$ is nontrivial and simple. Now π_i annihilates all but one of the simple factors of B^0 . Obviously no simple factor of B^0 is annihilated by every π_i . So now $B^0 = \prod B_\alpha^0$ where the B_α^0 are simple and where the index set I for $G = \prod_I G_i$ is a disjoint union of subsets I_α with $B_\alpha^0 \subset \prod_{i \in I_\alpha} G_i$. The proof of Proposition 2.1 is reduced to the case where B^0 (and thus also A^0) is simple, and there it is reduced to the proof that G_u is simple.

We may now assume B^0 simple. Suppose that G_u is not simple. Projecting to $G_1 \times G_2$ we may assume $G = G_1 \times G_2$. View the isomorphisms $\pi_i : B^0 \cong \pi_i(B^0)$ as identifications. Denote $E_i = \pi_i(B_\mathbb{C}^0)$, the complexification of the image of B^0 in G_i . Denote $E_{u,1} = \pi_i(B^0)$, the compact real form of E_i . Denote $P_i = E_i \cap Q_i$, the parabolic subgroup of E_i that is its isotropy subgroup in Z_i , so $Z_i = E_i/P_i$. Now $B_\mathbb{C}^0 = \{(e, e) \mid e \in E_1\}$, $B_\mathbb{C}^0 \cap Q = \{(p, p) \mid p \in (P_1 \cap P_2)\}$, and $Z = B_\mathbb{C}^0/(B_\mathbb{C}^0 \cap Q) \cong E_1/(P_1 \cap P_2)$. In particular $P_1 \cap P_2$ is a parabolic subgroup of E_1 . Compute complex dimensions: $\dim E_1 - \dim(P_1 \cap P_2) = \dim B^0 - \dim(B^0 \cap Q) = \dim Z = \dim Z_1 + \dim Z_2 = (\dim E_1 - \dim P_1) + (\dim E_1 - \dim P_2)$. On the Lie algebra level this says $\dim \mathfrak{e}_1 = \dim \mathfrak{p}_1 + \dim \mathfrak{p}_2 - \dim(\mathfrak{p}_1 \cap \mathfrak{p}_2)$, in other words $\mathfrak{p}_1 + \mathfrak{p}_2 = \mathfrak{e}_1$. As $\mathfrak{p}_1 \cap \mathfrak{p}_2$ is a parabolic subalgebra of \mathfrak{e}_1 we have a Cartan subalgebra \mathfrak{h} and a Borel subalgebra \mathfrak{s} with $\mathfrak{h} \subset \mathfrak{s} \subset \mathfrak{p}_1 \cap \mathfrak{p}_2$. In the root order such that \mathfrak{s} is the sum of \mathfrak{h} and the negative root spaces, no parabolic containing \mathfrak{s} can contain the root space for the maximal root. This contradicts $\mathfrak{p}_1 + \mathfrak{p}_2 = \mathfrak{e}_1$. The contradiction proves G_u simple and completes the proof. \square

Combining Proposition 2.1 with Theorem 1.6 we have

Theorem 2.2. *Let $Z = G/Q$, the complex flag manifold, where G is a complex connected semisimple Lie group acting with finite kernel on Z . Then the closed subgroups $A \subset G$ transitive on Z are precisely those given as follows. Decompose $G = \prod G_i$ with G_i simple, so $Z = \prod Z_i$ with $Z_i = G_i/(Q \cap G_i)$. Then $A = A^0 F$ where $A^0 = \prod A_i$ with $A_i = (A \cap G_i)^0$, and A_i is equal to G_i , or to its compact real form $G_{u,i}$, or to one of the three types listed in Theorem 1.6, and F is any subgroup of the center of G . Here F acts trivially on Z , so A and A^0 have the same action on Z .*

Picking out the cases where A is a real form of G we have, as in Corollary 1.7,

Corollary 2.3. *Let $Z = G/Q$, the complex flag manifold, where G is a complex connected semisimple Lie group acting with finite kernel on Z . Then the (connected) real forms $G_0 \subset G$ transitive on Z are precisely those given as follows. Decompose $G = \prod G_i$ with G_i simple, so $Z = \prod Z_i$ with $Z_i = G_i/(Q \cap G_i)$. Then $A = \prod A_i$ where $A_i = A \cap G_i$ either is the compact real form $G_{u,i}$ of G_i or is one of the two types listed in Corollary 1.7.*

REFERENCES

- [M] D. Montgomery, Simply connected homogeneous spaces, Proc. Amer. Math. Soc. **1** (1950), 467–469. MR **12**:242c
- [O1] A. L. Onishchik, Inclusion relations among transitive compact transformation groups. Trudy Moskov. Mat. Obšč. **11** (1962), 199–142.

- [O2] A. L. Onishchik, *Topology of Transitive Transformation Groups*, Johann Ambrosius Barth, Leipzig/Berlin/Heidelberg, 1994. MR **95e**:57058
- [W1] J. A. Wolf, The automorphism group of a homogeneous almost complex manifold. *Trans. Amer. Math. Soc.* **144** (1969), 535–543. MR **41**:956

INSTITUT FÜR MATHEMATIK, RUHR-UNIVERSITÄT BOCHUM, D-44780 BOCHUM, GERMANY

(Permanent address) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY, CALIFORNIA 94720-3840

E-mail address: `jawolf@math.berkeley.edu`