

TOPOLOGICAL INSIGHTS FROM THE CHINESE RINGS


JÓZEF H. PRZYTYCKI AND ADAM S. SIKORA

(Communicated by Ronald A. Fintushel)

ABSTRACT. L. Kauffman conjectured that a particular solution of the Chinese Rings puzzle is the simplest possible. We prove his conjecture by using low-dimensional topology and group theory. We notice also a surprising connection between the Chinese Rings and Habiro moves (related to Vassiliev invariants).

1. INTRODUCTION

In some of the popular puzzles one is supposed to take a ring off a rope which is usually tangled with the rigid part of the puzzle. Typically, such puzzles possess ingenious solutions which, if considered carefully enough, lead to interesting problems in low-dimensional topology. This phenomenon can be observed in the Chinese Rings which are shown below. The purpose of the puzzle is to take the loose ring off the rope.



Received by the editors September 10, 1999 and, in revised form, August 11, 2000.

1991 *Mathematics Subject Classification*. Primary 57M25; Secondary 05C10.

Key words and phrases. Chinese Rings, puzzle, Habiro move.

Although this puzzle was for a long time a toy of the first author's children, we became seriously interested in it only after L. Kauffman pointed out to us some interesting mathematical aspects of it, [K2]. The history of the Chinese Rings can be found in [K2, BC]. The goal of this paper is to prove that a particular solution of the Chinese Rings puzzle is the simplest possible, as conjectured in [K2].

If you have not thought about this puzzle before, then we suggest that you try to solve it now in order to gain an appreciation for this beautiful problem. Do not be discouraged by the initial difficulty in finding a solution. Indeed it is almost impossible to see all the necessary moves to be performed on the rope just by staring at the picture above. However, it is easy to see that a solution exists: Imagine for a moment that the toy is made out of a flexible material and press the longer columns down so that all columns have the same length, as shown in Figure 1.




FIGURE 1.

Now the solution becomes obvious! Since now the rope is completely separated from the columns, it can also be untangled in the original puzzle. (Actually, we need to be sure that the rope is long enough for this theoretical solution to be correct; this can be confirmed by direct experimentation.)

The above solution shows how a simple topological idea can yield a beautiful solution to what seems to be a complicated problem. The same idea shows that the Chinese Rings can be presented in a somewhat more regular form (see Figure 2). Note that we replaced the rope by another column with an attached ring. The problem now is to separate the loose ring (which is assumed to be infinitely elastic) from the solid part of the puzzle. The problem addressed in this paper is to find the simplest solution to this puzzle. As we will soon see, the solution to this problem involves an inductive argument in the number of columns. For that reason we




FIGURE 2.




FIGURE 3.

consider a more general version of the puzzle of Figure 2 composed of an arbitrary number of columns (all but one with an attached ring at the top).

A precise answer to our problem requires an objective measure of the complexity of possible solutions to the puzzle. For that, imagine an arc A (drawn in Figure 3 with a dashed line), joining the highest column of the toy with the base. The complexity of a solution for the puzzle is the minimal number of times the elastic ring passes through the arc A in the process of that solution.


Theorem 1.1. *The minimal complexity of a solution for the Chinese Rings with n columns presented in Figure 3 is 2^{n-1} .*

This result was conjectured by L. Kauffman in [K2], who wrote:

“This problem in the topology of the Chinese Rings is a useful test case for questions that can arise in applications of knot theory to natural structures where there is always a mixture of topology and mechanical/geometrical modeling.

A solution to the Ring Conjecture will probably involve the discovery of new techniques for understanding topology of graph embeddings in three-dimensional space. It is fun to be able to take a classical puzzle as fascinating as the Chinese Rings and find within it a significant topological problem. Let us find the solution!”

Habiro moves. There is a surprising connection between the Chinese Rings (known for hundreds of years) and a very recent theory of Vassiliev invariants of knots, [BL]. K. Habiro [H] proved that two knots cannot be distinguished by Vassiliev invariants of order $\leq n$ if and only if they are related by a sequence of moves (called Habiro moves) presented below:



Note that the Habiro move corresponds exactly to the operation of untying the loose ring in the version of Chinese Rings presented in Figure 3!

2. PROOF

The solution of the puzzle presented at the beginning of the paper requires that we first untangle the rings attached to the columns. Note that as a result of this deformation of the puzzle, the arc A will assume the form presented in Figure 4.

Because we can now remove the ring by passing through the 2^{n-1} strands of A , the complexity of this algorithm is obviously at most 2^{n-1} . We are going to show that this is the simplest solution to the Chinese Rings; i.e. the complexity of any other solution is not less than 2^{n-1} . For the purpose of the proof we are allowed to relax the conditions of the problem and consider it in a topological setting by allowing arbitrary deformations of the dimensions and the shape of the toy. Moreover we add a point at infinity to the ambient three-space and hence consider the problem in S^3 . These assumptions make the problem simpler and surely do not increase the complexity of the minimal solution.

The body of the Chinese Rings with n columns (and with the loose ring excluded) is a handlebody H_{n-1} of genus $n - 1$ embedded in the standard way into S^3 . Its complement, H'_{n-1} , is also a handlebody of genus $n - 1$. The loose ring is contractible in H'_{n-1} , and this is the reason for which the puzzle has a solution. The arc A assumes a complicated position in H'_{n-1} which can be deduced from Figure 4; see Figure 5. If we remove the arc A from H'_{n-1} , then the loose ring, S^1 , will no longer be contractible.

Recall that the complexity of a solution for the puzzle is the minimal number of passes of the ring through A necessary for contracting the ring to a point in $H'_{n-1} \setminus A$. We need to show that this number is at least 2^{n-1} . Unfortunately, the position of the ring S^1 in $H'_{n-1} \setminus A$ is very complicated; try to figure it out by yourself to see that, indeed, it is not an easy problem! In order to avoid this problem we use three tricks: First, we take a dual approach: we fix the ring S^1 in H'_{n-1} and count the number of times the arc A has to pass through S^1 in order to make S^1 contractible in $H'_{n-1} \setminus A$. Second, we slightly relax the conditions of the




FIGURE 4.




FIGURE 5. Complement of the puzzle = H'_{n-1}




FIGURE 6.




FIGURE 7.



FIGURE 8. $a_1 = g_1$ and $a_2 = g_2g_1g_2^{-1}$

puzzle, by assuming that A can be deformed by an arbitrary homotopy fixing its endpoints. We assume that the endpoints of A are at some $x_0 \in \partial H'_{n-1}$. Therefore we consider A as an element of $\pi_1(H'_{n-1} \setminus S^1, x_0)$. Our intention is to prove that under these relaxed conditions the complexity of the puzzle is at least 2^{n-1} . This will surely imply that the complexity of the puzzle under original assumptions is also at least 2^{n-1} .

Third, we modify the puzzle by attaching the loose ring to the base of the puzzle by an additional column, and we try to untangle A in the modified puzzle. This seems to be a more difficult problem, but actually it is not. Observe that the modified puzzle is homeomorphic to a handlebody of genus n , H_n and that its complement is homeomorphic also to a handlebody, H'_n . Notice also that we have an embedding $i : H'_n \rightarrow H'_{n-1} \setminus S^1$ corresponding to the fact that the modified toy differs from the original only by an additional column. One can prove using Van-Kampen's Theorem that the map i induces an isomorphism of the fundamental groups $i_* : \pi_1(H'_n) \rightarrow \pi_1(H'_{n-1} \setminus S^1)$. Therefore the additional column does not create any new obstacle for A ! (Recall that we consider the arc A up to homotopy only.) The benefit of this trick is that the modified puzzle is much easier to solve because it does not have any loose rings. Observe that the arc A lies in H'_n in the pattern presented in Figure 6.

If we denote the loops going around the holes in H'_n in the manner presented in Figure 7 by g_1, g_2, \dots, g_n , then $F_n = \pi_1(H'_n, x_0)$ is the free group on generators g_1, \dots, g_n . We want to determine the presentation of the element $a_n \in F_n$ representing the arc A given as in Figure 6, with, say, an anti-clock orientation. The presentations of a_1 and a_2 are as in Figure 8.

Observe that a_{n+1} can be built inductively from a_n by adding the $n + 1$ -th hole and twisting it with the n -th hole by 180° ; see Figure 6. This operation corresponds to replacing all $g_n^{\pm 1}$ in the presentation of a_n by $g_{n+1}g_n^{\pm 1}g_{n+1}^{-1}$. Therefore

$$a_3 = g_3g_2g_3^{-1}g_1g_3g_2^{-1}g_3^{-1}, \quad a_4 = g_4g_3g_4^{-1}g_2g_4g_3^{-1}g_4^{-1}g_1g_4g_3g_4^{-1}g_2^{-1}g_4g_3^{-1}g_4^{-1}, \text{ etc.}$$

Note that each such presentation is reduced, i.e. none of the words a_n has a subword of the form $g_i^{\pm 1}g_i^{\mp 1}$. Indeed, this is true for $n = 1, 2, 3, 4$. In general, if this is true for some n , then it is also true for $n + 1$: Suppose that a_n is reduced. The only difference between a_n and a_{n+1} is that g_n in a_n is replaced by $g_{n+1}g_ng_{n+1}^{-1}$. Therefore, a_{n+1} cannot contain $g_i^{\pm 1}g_i^{\mp 1}$ for $i \leq n$. Hence, if a_{n+1} was not reduced, it would have to have a reduction in g_{n+1} 's. That is, it would have to contain a subword composed of $g_{n+1}g_n^{\pm 1}g_{n+1}^{-1}$ and its inverse. But this is impossible, since this would mean that a_n contains the subword $g_n^{\pm 1}g_n^{\mp 1}$.

Observe also that the number of appearances of $g_n^{\pm 1}$ in a_n is 2^{n-1} . This can also be proved by induction: g_1 appears once in a_1 , and g_{n+1} appears in a_{n+1} twice as many times as g_n in a_n .

Therefore we proved the following:

Proposition 2.1. *The above inductively defined presentation of $a_n \in F_n$ is reduced and $g_n^{\pm 1}$ appears 2^{n-1} times in it.*

Recall that we are supposed to count the minimal number of times that the ring S^1 has to pass through the arc A (in the original puzzle) in a process of contracting it to a point. Equivalently, we can calculate the number of times the arc A has to pass through the n -th hole in H'_n in order to be placed inside $H'_{n-1} \subset H'_n$ (in the situation in which the modified puzzle is considered). We claim that this number is at least 2^{n-1} .

Whenever A passes through the n -th hole, $xg_n^{\pm 1}x^{-1}$ is inserted in a word representing A or deleted from it. Intuitively, since $g_n^{\pm 1}$ appears 2^{n-1} times in a_n one has to repeat this process 2^{n-1} times. This may seem as an obvious fact at first, but after a closer look one can realize that it requires a proof. We will see in the next section that this problem is a special case of an interesting problem in the combinatorial group theory. In order to finish our proof we need the following fact.

Proposition 2.2. *If letters $b_1, b_2, \dots, b_k \in \{g_1^{\pm 1}, \dots, g_{n-1}^{\pm 1}\}$ form a reduced word $b_1b_2\dots b_k$, then the minimal number of insertions or deletions of conjugates of $g_n^{\pm 1}$ into a word $w = g_n^{\alpha_0}b_1g_n^{\alpha_1}b_2\dots b_kg_n^{\alpha_k} \in F_n$ necessary for transforming w into an element of F_{n-1} is $\sum_{i=0}^k |\alpha_i|$.*

We will prove the above proposition in the next section. Right now we complete the proof of the main theorem: Recall that a_n can be obtained from a_{n-1} by replacing all $g_{n-1}^{\pm 1}$ appearing in a_{n-1} by $g_n g_{n-1}^{\pm 1} g_n^{-1}$. Therefore, $a_n = g_n^{\alpha_0} b_1 g_n^{\alpha_1} b_2 \dots b_k g_n^{\alpha_k}$, where, by Proposition 2.1, $b_1 b_2 \dots b_k = a_{n-1}$ is reduced and $\sum_{i=0}^k |\alpha_i| = 2^{n-1}$. By the last proposition one needs at least 2^{n-1} insertions of conjugates of $g_n^{\pm 1}$ into a word representing a_n or deletions of such conjugates from a_n in order to eliminate all appearances of g_n in it. Therefore the arc A has to pass at least 2^{n-1} times through the n -th hole in H'_n in order to be placed inside $H'_{n-1} \subset H'_n$.

Final remark. We found the complexity of the Chinese Rings by the study of the homotopical properties of the arc A in a handlebody and by the use of the fundamental group. One can, however, consider more sophisticated puzzles, whose




FIGURE 9.

solutions require more advanced tools than homotopy theory. For example in order to show that the arc A of Figure 9 cannot be contracted to a point (without crossing changes) one can use the Kauffman bracket skein module of the solid torus [P] – an algebraic construction which generalizes the Kauffman bracket polynomial, [K1].

3. PROBLEMS IN GROUP THEORY INVOLVED
IN THE PROOF OF KAUFFMAN’S CONJECTURE

The group theoretical problem which we encountered at the end of the proof of the main theorem can be presented in a more general form as follows: Consider a free group $F = \langle g_1, g_2, \dots \mid \rangle$ and a quotient map $\pi : F \rightarrow G = \langle g_1, g_2, \dots \mid r_1, r_2, \dots \rangle$. By an elementary operation on $w \in F$ we mean an insertion of a conjugate of $r_i^{\pm 1}$ into a word representing w . Observe that a deletion of $xr_i^{\pm 1}x^{-1}$ from a word representing w can be also realized as an insertion of $x^{-1}r_i^{\mp 1}x$ into this word (followed by a reduction) and, therefore, it is also an elementary operation. Observe also that two elements $x, y \in F$ have the same image, $\pi(x) = \pi(y)$, if and only if they are related by a sequence of elementary operations.

Problem 1. Given $x, y \in F, \pi(x) = \pi(y)$, what is the minimal number of elementary operations necessary for transforming x into y ? We denote this number by (x, y) . If $\pi(x) \neq \pi(y)$, then we set $(x, y) = \infty$.

The above problem can be stated in a different form: Any $w \in Ker(\pi)$ can be presented as $w = \prod_{i=1}^k p_i r_{\alpha_i}^{\pm 1} p_i^{-1}$, and we define $\|w\|$ to be the minimal number k of conjugates of $r_{\alpha}^{\pm 1}$ appearing in such a presentation. If $w \notin Ker(\pi)$, then $\|w\| = \infty$.

Problem 2. Given $w \in Ker(\pi)$ calculate $\|w\|$.

It turns out that that Problems 1 and 2 are equivalent and that the symbols (\cdot, \cdot) and $\|\cdot\|$ have several interesting properties:

Proposition 3.1. (1) $\|\cdot\| : F \rightarrow \{0, 1, 2, \dots, \infty\}$ is a “norm” on F , i.e. (a) $\|w\| = 0$ iff $w = e$ in F , (b) $\|vw\| \leq \|v\| + \|w\|$, (c) $\|w^{-1}\| = \|w\|$.

(2) $(x, y) = \|xy^{-1}\|$ and (\cdot, \cdot) is a metric on F , which is finite on $\pi^{-1}(g)$, for any $g \in G$.

Proof. The proof of (1) is straightforward. Also, the claim that (\cdot, \cdot) is a metric on F follows immediately from (1) and the equality $(x, y) = \|xy^{-1}\|$. Therefore we include only the proof of that equality: If $\pi(x) \neq \pi(y)$, then $(x, y) = \infty = \|xy^{-1}\|$. Hence we can assume that $\pi(x) = \pi(y)$. Since $xy^{-1} = \prod_{i=1}^k p_i r_{\alpha_i}^{\pm 1} p_i^{-1}$, where $k = \|xy^{-1}\|$, $x = \prod_{i=1}^k p_i r_{\alpha_i}^{\pm 1} p_i^{-1} \cdot y$ can be transformed into y by k elementary operations. Therefore, $\|xy^{-1}\| \geq (x, y)$.

In order to prove the opposite inequality we need the following fact:

If z is obtained from 1 by k elementary operations, then z can be presented as $\prod_{i=1}^k p_i r_{\alpha_i}^{\pm 1} p_i^{-1}$. We prove this fact by induction. The statement is true for $k = 1$. Suppose it is also true for $k - 1$, and suppose that z is obtained from 1 by k elementary operations. By the inductive assumption, after $k - 1$ operations we get a product of $k - 1$ conjugates, z' . The word z is obtained by inserting a word $qr_{\beta}^{\pm 1}q^{-1}$ into z' . Therefore $z = z_1qr_{\beta}^{\pm 1}q^{-1}z_2$, for some z_1, z_2 such that $z' = z_1z_2$. Thus $z = z'z_2^{-1}qr_{\beta}^{\pm 1}q^{-1}z_2$ is a product of k conjugates.

Since x is obtained from y by (x, y) elementary operations, xy^{-1} may be obtained from $yy^{-1} = 1$ also by (x, y) elementary operations. Therefore, by the fact proved above, xy^{-1} can be presented as a product of (x, y) conjugates of $r_i^{\pm 1}$'s. Thus $\|xy^{-1}\| \leq (x, y)$. □

Notice that Problems 1-2 generalize the word problem in G and may be very difficult to solve in general. However, fortunately for us, the group theoretic problem encountered at the end of the previous section is a special, solvable, case of Problems 1-2:

From now on $\pi : F_n \rightarrow F_{n-1} \subset F_n$ will be a projection given by $\pi(g_i) = g_i$, for $i < n$ and $\pi(g_n) = 1$. Let $b_1b_2\dots b_k$ be a reduced word in F_{n-1} . We consider the problem of determining the minimal number of elementary operations on $w = g_n^{\alpha_0}b_1g_n^{\alpha_1}b_2\dots b_kg_n^{\alpha_k}$ necessary for transforming this word into an element of F_{n-1} . Since $w \in \pi^{-1}(b_1b_2\dots b_k)$, any word obtained by applying elementary operations to w will still be in $\pi^{-1}(b_1b_2\dots b_k)$. Observe that the only element of $\pi^{-1}(b_1b_2\dots b_k)$ which can be presented as a word in the letters $g_1^{\pm 1}, \dots, g_{n-1}^{\pm 1}$ (i.e. a word without the letter $g_n^{\pm 1}$ in it) is $b_1b_2\dots b_k \in F_{n-1} \subset F_n$. Therefore Proposition 2.2 can be restated as

Proposition 2.2'. *If letters $b_1, b_2, \dots, b_k \in \{g_1^{\pm 1}, \dots, g_{n-1}^{\pm 1}\}$ form a reduced word $b_1b_2\dots b_k$, then*

$$(g_n^{\alpha_0}b_1g_n^{\alpha_1}b_2\dots b_kg_n^{\alpha_k}, b_1b_2\dots b_k) = \sum_{i=0}^k |\alpha_i|.$$

Let P be a word of the form $s_1s_2\dots s_k$. Place points corresponding to s_i 's on the x -axis, s_{i+1} after s_i ; compare Figure 10. A connection, C , on P is a set of pairwise disjoint arcs in the upper half plane. Each arc connects some g_i with g_i^{-1} in such a way that only some $g_n^{\pm 1}$'s may be left unconnected. The norm of the connection, $|C|$, is the number of unconnected letters. Figure 10 shows two different connections for the word $P = g_1g_n g_1^{-1}g_n g_1 g_n^{-1}g_1^{-1}$, one of norm 3 and another of norm 1.

We denote by $|w|_c$ the minimum of $|C|$, where C varies over all connections on a word w . If w has no connection, then $|w|_c = \infty$.

Theorem 3.2. *For any word w , $|w|_c = \|w\|$.*

Observe that since each word has only a finite number of connections on it and it is easy to construct all of them, the above theorem gives an explicit method of calculating $\|w\|$.

Proof. 1. We prove $|w|_c \geq \|w\|$ first.

If w has no connection, then $|w|_c = \infty$ and the inequality is obvious. Therefore we may assume that w has a connection C . Consider two kinds of operations on w :




FIGURE 10.

(I) If there is an unconnected letter $g_n^{\pm 1}$ in w , then we delete this letter and we obtain a new word w' with a connection C' (composed of the same arcs as C).

(II) If there is a pair g_i, g_i^{-1} of letters in w connected by an arc which is not nested (i.e. g_i, g_i^{-1} are neighbors), then we remove these letters and the arc connecting them and we obtain a new word w' of a shorter length with a connection C' .

Observe that each of the above operations decreases the length of w , and we can always apply at least one of them to w , unless w is the trivial word 1. Therefore any word w can be reduced to 1 by a sequence of operations of the first and the second type. Observe that Operation (I) changes $\|w\|$ by at most one and decreases the norm of the connection on w by 1. Operation (II) does not change $\|w\|$ or the norm of the connection on w . Since at the end of the process (when $w = 1$) both $\|w\|$ and the norm of the connection on w are 0, $|C| \geq \|w\|$. Since this inequality holds for any connection C on w , we also have $|w|_c \geq \|w\|$.

2. We also claim that $|w|_c \leq \|w\|$.

We can assume that $w \in Ker(\pi)$, since otherwise $\|w\| = \infty$. There is a word $w' = \prod_{i=1}^k p_i g_n^{\pm 1} p_i^{-1}$ representing the same element of the group F_n as the word w , with $k = \|w\|$. Each factor $p_i g_n^{\pm 1} p_i^{-1}$ has a connection of a norm 1 (a nested family of arcs). Therefore the product w' has a connection of a norm $\|w\|$. The word w' can be transformed into the word w by a sequence of insertions and deletions of subwords of the form $g_i^{\pm 1} g_i^{\mp 1}$, $i = 1, 2, \dots, n$. Observe that after each insertion we obtain a new word with a connection of the same norm. Moreover, it is not difficult to see that after each deletion we obtain a new word with a connection of the same norm, if $i < n$, or lower or equal norm, if $i = n$. Therefore w has a connection C of norm $|C| \leq \|w\|$ and, hence, $|w|_c \leq \|w\|$. \square

Proof of Proposition 2.2'. Since

$$(g_n^{\alpha_0} b_1 g_n^{\alpha_1} b_2 \dots b_k g_n^{\alpha_k}, b_1 b_2 \dots b_k) \leq \sum_{i=0}^k |\alpha_i|,$$

we only need to prove the opposite inequality. By Proposition 3.1 and Theorem 3.2 it is enough to prove that each connection on

$$g_n^{\alpha_0} b_1 g_n^{\alpha_1} b_2 \dots b_k g_n^{\alpha_k} b_k^{-1} \dots b_2^{-1} b_1^{-1}$$

has its norm greater than or equal to $\sum_{i=0}^k |\alpha_i|$. Let C be any such connection. Observe that it is enough to prove that C does not connect any pair of letters $g_n^{\pm 1}, g_n^{\mp 1}$. Suppose that C does connect letters $g_n^{\pm 1}$ and $g_n^{\mp 1}$ enclosing a word $b_i g_n^{\alpha_i} b_{i+1} \dots g_n^{\alpha_{j-1}} b_j$. The connection C restricts to a connection on this word and therefore $b_i g_n^{\alpha_i} b_{i+1} \dots g_n^{\alpha_{j-1}} b_j \in \text{Ker}(\pi)$. Hence $b_i b_{i+1} \dots b_j = e$ in $F_{n-1} = \langle g_1, \dots, g_{n-1} \rangle$ and therefore $b_1 b_2 \dots b_k$ is reducible, which contradicts our assumption. \square

REFERENCES

- [BC] W.W. Rouse Ball and H.S.M. Coxeter, *Mathematical Recreations and Essays*, University of Toronto Press, Toronto, 1974 (First edition - 1892). MR **50**:4229
- [BL] J. S. Birman, X. L. Lin, Knot polynomials and Vassiliev's invariants, *Invent. Math.* **111** (1993), 225–270. MR **94d**:57010
- [H] K. Habiro, Claspers and finite type invariants of links, *Geom. Topol.* **4** (2000), 1–83, <http://www.maths.warwick.ac.uk/gt/GTVol4/paper1.abs.html> CMP 2000:07
- [K1] L. H. Kauffman, An invariant of regular isotopy, *Trans. Amer. Math. Soc.* **318** (1990), no. 2, 417–471. MR **90g**:57007
- [K2] L. H. Kauffman, Tangle complexity and the topology of the Chinese rings, in *Mathematical approaches to biomolecular structure and dynamics*, IMA Vol. 82, Springer, New York, 1996, 1–10. MR **98b**:57013
- [P] J. H. Przytycki, Fundamentals of Kauffman bracket skein modules, *Kobe J. Math.*, 16(1), 1999, 45–66, <http://xxx.lanl.gov/abs/math>. GT/9809113 MR **2000i**:57015

DEPARTMENT OF MATHEMATICS, THE GEORGE WASHINGTON UNIVERSITY, WASHINGTON, DC 20052

E-mail address: przytyck@gwu.edu

Current address: Department of Mathematics, University of Maryland, College Park, Maryland 20742

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MARYLAND, COLLEGE PARK, MARYLAND 20742

E-mail address: asikora@math.umd.edu