

FINITE DIMENSIONAL REPRESENTATIONS OF THE SOFT TORUS

SØREN EILERS AND RUY EXEL

(Communicated by David R. Larson)

ABSTRACT. The soft tori constitute a continuous deformation, in a very precise sense, from the commutative C^* -algebra $C(\mathbb{T}^2)$ to the highly non-commutative C^* -algebra $C^*(\mathbb{F}_2)$. Since both of these C^* -algebras are known to have a separating family of finite dimensional representations, it is natural to ask whether that is also the case for the soft tori. We show that this is in fact the case.

1. INTRODUCTION

Knowing that a given C^* -algebra has many representations on finite dimensional Hilbert spaces is of great importance to understanding structural properties of it. Among C^* -algebras, those who possess a separating family of finite dimensional representations are called *residually finite dimensional* or just *RFD*. This class was studied in [12], [11] and [1], and more recent insight about it has led to important advances in classification theory and the theory of quasidiagonal C^* -algebras (see, e.g., [2], [5] and [6]).

For any $\varepsilon \geq 0$ we define a C^* -algebra A_ε as the universal (unital) C^* -algebra defined by the generators u, v subject to the relations

$$uu^* = u^*u = 1, \quad vv^* = v^*v = 1, \quad \|uv - vu\| \leq \varepsilon.$$

As recorded in [8], A_0 is the commutative C^* -algebra of functions over the torus \mathbb{T}^2 , and A_ε is the full C^* -algebra of the free group of two generators \mathbb{F}_2 whenever $\varepsilon \geq 2$. For ε between 0 and 2 we get a class of C^* -algebras which are commonly referred to as *soft tori*. These C^* -algebras are of relevance to several problems in operator algebra theory (see [10]) and have been extensively studied in [3], [8], [9], [7].

The starting point of the investigation reported on in the present paper is a result from [9], stating that the soft tori form a continuous field interpolating between the (hard) torus and the group C^* -algebra of the free group. Since $C(\mathbb{T}^2)$ is obviously RFD, and since $C^*(\mathbb{F}_2)$ was proved to be RFD in [4] — a surprise at the time — we are naturally led to the question of whether the same is true for the interpolating family A_ε . We are going to prove that this is the case.

Received by the editors November 19, 1998 and, in revised form, August 28, 2000.
2000 *Mathematics Subject Classification*. Primary 46L05; Secondary 46L65, 46L85, 47B20.
This work was partially supported by the Carlsberg Foundation.

2. METHODS

We prove that A_ε is RFD using an adaptation of the compression argument developed by Choi in [4] to prove that $C^*(\mathbb{F}_2)$ is RFD. However, Choi’s argument does not apply directly to A_ε since one cannot arrange for the compressions to satisfy the commutation relation. We instead argue via an auxiliary C^* -algebra, thus employing a method from [3] and [8] which lies behind many results about the structural properties of A_ε .

We define B_ε as the universal C^* -algebra given by the generators $\{u_n\}_{n \in \mathbb{Z}}$ and the relations

$$(1) \quad u_n u_n^* = u_n^* u_n = 1, \quad \|u_{n+1} - u_n\| \leq \varepsilon.$$

Clearly one can define an automorphism α on B_ε by

$$u_n \mapsto u_{n+1},$$

and as seen in [8] one has

$$A_\varepsilon = B_\varepsilon \rtimes_\alpha \mathbb{Z}.$$

There is a faithful conditional expectation $E_\alpha : A_\varepsilon \rightarrow B_\varepsilon$.

Our strategy will be to prove that A_ε is RFD by proving that B_ε is RFD in a way which is covariant with α .

3. FINITE DIMENSIONAL REPRESENTATIONS OF B_ε

We start out by finding a new picture of B_ε by generators and relations.

Lemma 3.1. *For any $\varepsilon < 2$, B_ε is isomorphic to the universal C^* -algebra generated by $v_0, \{h_n\}_{n \in \mathbb{Z}}$ subject to the relations*

$$(2) \quad v_0 v_0^* = v_0^* v_0 = 1, \quad h_n = h_n^*, \quad \|h_n\| \leq \frac{2}{\pi} \arcsin(\varepsilon/2).$$

Proof. Let us denote the C^* -algebra generated by v_0 and h_n subject to (2) by B'_ε . We can define a map $\varphi : B_\varepsilon \rightarrow B'_\varepsilon$ by

$$u_n \mapsto \begin{cases} e^{i\pi h_n} \dots e^{i\pi h_1} v_0, & n > 0, \\ v_0, & n = 0, \\ e^{-i\pi h_n} \dots e^{-i\pi h_{-1}} v_0, & n < 0, \end{cases}$$

since the elements to the right of the arrow above satisfy the relations (1). Similarly, the universal property of B'_ε allows for a map $\psi : B'_\varepsilon \rightarrow B_\varepsilon$ defined by

$$v_0 \mapsto u_0, \quad h_n \mapsto \frac{1}{i\pi} \text{Log}(u_n u_{n-1}^*).$$

Clearly φ and ψ are each others’ inverse. □

This characterization can be used to shorten the proof of [8, 2.2], stating that B_ε is homotopic to $C(\mathbb{T})$. To see this, define maps $\varphi : B'_\varepsilon \rightarrow C(\mathbb{T})$ and $\psi : C(\mathbb{T}) \rightarrow B'_\varepsilon$ by the correspondence $v_0 \leftrightarrow [z \mapsto z], h_n \leftrightarrow 0$. Clearly $\varphi\psi = \text{id}_{C(\mathbb{T})}$, and $\chi_t : B'_\varepsilon \rightarrow B'_\varepsilon$ given by

$$\chi_t(v_0) = v_0, \quad \chi_t(h_n) = th_n$$

provides a homotopy from $\text{id}_{B'_\varepsilon}$ to $\psi\varphi$.

In the following proof, we denote by $\text{Alg}(X)$ the smallest $*$ -algebra, not necessarily closed, generated by the set X inside some C^* -algebra.

Proposition 3.2. *For any $\varepsilon < 2$, B_ε is RFD. In fact, for any $0 \neq b \in B_\varepsilon$ there exists $n \in \mathbb{N}$, an automorphism β of \mathbf{M}_n and a representation $\rho : B_\varepsilon \rightarrow \mathbf{M}_n$ with the properties*

$$\rho(b) \neq 0, \quad \beta\rho = \rho\alpha.$$

Proof. For the first claim we use the characterization of B_ε given by the relations in (2) and proceed as in [4]. Fix a faithful non-degenerate representation $\pi : B_\varepsilon \rightarrow \mathbb{B}(\mathcal{H})$, where we may assume that \mathcal{H} is a separable Hilbert space.

Let P_m be a sequence of projections, with $\text{rank}(P_m) = m$, converging strongly to the unit of $\mathbb{B}(\mathcal{H})$, and abbreviate

$$T_{0,m} = P_m\pi(v_0)P_m, \quad K_{n,m} = P_m\pi(h_n)P_m.$$

Now note that for each m the collection of elements $\{V_{0,m}, H_{n,m} : n \in \mathbb{Z}\}$ defined by

$$V_{0,m} = \begin{bmatrix} T_{0,m} & \sqrt{P_m - T_{0,m}T_{0,m}^*} \\ \sqrt{P_m - T_{0,m}^*T_{0,m}} & -T_{0,m}^* \end{bmatrix},$$

$$H_{n,m} = \begin{bmatrix} K_{n,m} & 0 \\ 0 & K_{n,m} \end{bmatrix}$$

satisfies (2) in $\mathbf{M}_2(P_m\mathbb{B}(\mathcal{H})P_m) \simeq \mathbf{M}_{2m}$. Consequently we get representations $\pi_m : B_\varepsilon \rightarrow \mathbf{M}_{2m}$. We are going to check, following [4], that

$$\underline{\pi} : B_\varepsilon \rightarrow \prod_{m=1}^\infty \mathbf{M}_{2m}, \quad \underline{\pi}(b) = (\pi_m(b))_{m=1}^\infty$$

is an isometry. It suffices to check that $\|\underline{\pi}(x)\| \geq \|x\| - \eta$ for any $\eta > 0$ and any $x \in \text{Alg}(\{v_0, h_{-N}, \dots, h_N\})$. Fix η, N and x and write

$$x = F(v_0, h_{-N}, \dots, h_N)$$

where F is some finite linear combination of finite words in $2N + 2$ variables and their adjoints. Since, when m goes to infinity,

$$V_{0,m} \rightarrow \begin{bmatrix} \pi(v_0) & 0 \\ 0 & -\pi(v_0)^* \end{bmatrix}, \quad H_{n,m} \rightarrow \begin{bmatrix} \pi(h_n) & 0 \\ 0 & \pi(h_n) \end{bmatrix},$$

strongly in the unit ball of $\mathbf{M}_2(\mathbb{B}(\mathcal{H}))$ we conclude that

$$\begin{aligned} & \limsup_{m \rightarrow \infty} \|F(V_{0,m}, H_{-N,m}, \dots, H_{N,m})\| \\ & \geq \left\| \lim_{m \rightarrow \infty} F(V_{0,m}, H_{-N,m}, \dots, H_{N,m}) \right\| \\ & = \left\| \begin{bmatrix} \pi(F(v_0, h_{-N}, \dots, h_N)) & 0 \\ 0 & \pi(F(-v_0^*, h_{-N}, \dots, h_N)) \end{bmatrix} \right\| \\ & \geq \|\pi(x)\| = \|x\|. \end{aligned}$$

We can hence find m such that

$$\|\underline{\pi}(x)\| \geq \|\pi_m(x)\| = \|F(V_{0,m}, H_{-N,m}, \dots, H_{N,m})\| \geq \|x\| - \eta.$$

For the second claim, we go back to the original presentation (1) of B_ε by unitary generators only. For a given $b \in B_\varepsilon$ with $\|b\| = 1$ we fix, using the first part of the proposition, a finite dimensional representation $\pi : B_\varepsilon \rightarrow \mathbf{M}_m$ with $\|\pi(b)\| > \frac{3}{4}$. We also fix c and $N \in \mathbb{N}$ such that

$$\|b - c\| < \frac{1}{4}, \quad c \in \text{Alg}(u_{-N}, \dots, u_N).$$

Choose $M > 0$ and unitaries $v_0^\pm, \dots, v_M^\pm \in \mathbf{M}_m$ with the properties

$$\|v_{n+1}^\pm - v_n^\pm\| \leq \varepsilon, \quad v_0^\pm = \pi(u_{\pm N}), \quad v_M^\pm = 1.$$

There is then exactly one representation $\pi' : B_\varepsilon \rightarrow \mathbf{M}_m$ which is $2(N + M)$ -periodic in the sense that $\pi'(u_n) = \pi'(u_{n+2(N+M)})$ and satisfies

$$\pi'(u_n) = \begin{cases} v_{-N-n}^-, & -M - N \leq n < -N, \\ \pi(u_n), & -N \leq n \leq N, \\ v_{n-N}^+, & N < n \leq N + M. \end{cases}$$

Note that $\pi'(c) = \pi(c)$; in particular $\|\pi'(c)\| \geq \frac{1}{2}$.

Now let $n = 2(N + M)m$. With β defined as the backward cyclic shift in block form (with period $2(N + M)$) we may define a covariant representation ρ of B_ε on \mathbf{M}_n by

$$u_i \mapsto \begin{bmatrix} \pi'(u_i) & & & \\ & \pi'(u_{i+1}) & & \\ & & \ddots & \\ & & & \pi'(u_{i+2(N+M)-1}) \end{bmatrix}.$$

We have

$$\|\rho(b)\| \geq \|\pi'(b)\| \geq \|\pi'(c)\| - \frac{1}{4} > 0.$$

□

4. FINITE DIMENSIONAL REPRESENTATIONS OF A_ε

Theorem 4.1. *For any $\varepsilon > 0$, let A_ε be the universal C^* -algebra generated by a pair of unitaries subject to the relation $\|uv - vu\| \leq \varepsilon$. Then A_ε is residually finite dimensional in the sense that it admits a separating family of finite dimensional representations.*

Proof. We may assume that $0 < \varepsilon < 2$. Let $0 \neq a \in A_\varepsilon$. Then also $b = E_\alpha(a^*a)$ is nonzero, for the conditional expectation is faithful. Choose n, ρ and β as in Proposition 3.2 and define

$$\pi : A_\varepsilon = B_\varepsilon \rtimes_\alpha \mathbb{Z} \rightarrow \mathbf{M}_n \rtimes_\beta \mathbb{Z}$$

as the extension to the crossed product of the covariant $*$ -homomorphism ρ . We then have, with E_β the conditional expectation from $\mathbf{M}_n \rtimes_\beta \mathbb{Z}$ to \mathbf{M}_n ,

$$E_\beta(\pi(a^*a)) = \pi(E_\alpha(a^*a)) = \rho(b) \neq 0,$$

so $\pi(a) \neq 0$.

Note finally that since β is inner,

$$\mathbf{M}_n \rtimes_\beta \mathbb{Z} = \mathbf{M}_n \rtimes_{\text{id}} \mathbb{Z} \simeq C(\mathbb{T}) \otimes \mathbf{M}_n.$$

Therefore we may compose π with an evaluation map of $C(\mathbb{T}) \otimes \mathbf{M}_n$ to exhibit an n -dimensional representation which does not vanish on a . □

Linear algebra tells us that \mathbf{M}_n has a faithful tracial state and has the property that every matrix x which is *hyponormal* is the sense that

$$x^*x \geq xx^*$$

is in fact normal. As in [4], we may conclude:

Corollary 4.2. *For any ε , A_ε has a faithful tracial state, and any hyponormal operator in A_ε is normal.*

Proof. Such properties clearly pass from matrices to sums of the form $\prod_{n \in \mathbb{N}} \mathbf{M}_{m_n}$, and from these sums to any of their subalgebras. By the theorem, A_ε is one such. \square

REFERENCES

- [1] R. J. Archbold, *On residually finite-dimensional C^* -algebras*, Proc. Amer. Math. Soc. **123** (1995), no. 9, 2935–2937. MR **95m**:46089
- [2] B. Blackadar and E. Kirchberg, *Generalized inductive limits of finite-dimensional C^* -algebras*, Math. Ann. **307** (1997), no. 3, 343–380. MR **98c**:46112
- [3] C. Cerri, *Non-commutative deformations of $C(\mathbb{T}^2)$ and K -theory*, Internat. J. Math. **8** (1997), no. 5, 555–571. MR **98j**:46082
- [4] M.D. Choi, *The full C^* -algebra of the free group on two generators*, Pacific J. Math. **87** (1980), no. 1, 41–48. MR **82b**:46069
- [5] M. Dădărlat, *Nonnuclear subalgebras of AF algebras*, Amer. J. Math. **122** (2000), no. 3, 581–597. CMP 2000:13
- [6] ———, *On the approximation of quasidiagonal C^* -algebras*, J. Funct. Anal. **167** (1999), no. 1, 69–78. MR **2000f**:46069
- [7] G.A. Elliott, R. Exel, and T.A. Loring, *The soft torus. III. The flip*, J. Operator Theory **26** (1991), no. 2, 333–344. MR **94f**:46086
- [8] R. Exel, *The soft torus and applications to almost commuting matrices*, Pacific J. Math. **160** (1993), 207–217. MR **94f**:46091
- [9] R. Exel, *The soft torus: a variational analysis of commutator norms*, J. Funct. Anal. **126** (1994), no. 2, 259–273. MR **95i**:46085
- [10] R. Exel and T.A. Loring, *Invariants of almost commuting unitaries*, J. Funct. Anal. **95** (1991), 364–376. MR **92a**:46083
- [11] ———, *Finite-dimensional representations of free product C^* -algebras*, Internat. J. Math. **3** (1992), 469–476. MR **93f**:46091
- [12] K. R. Goodearl and P. Menal, *Free and residually finite-dimensional C^* -algebras*, J. Funct. Anal. **90** (1990), no. 2, 391–410. MR **91f**:46078

MATEMATISK AFDELING, KØBENHAVNS UNIVERSITET, UNIVERSITETSPARKEN 5, DK-2100 COPENHAGEN Ø, DENMARK

E-mail address: eilers@math.ku.dk

URL: <http://www.math.ku.dk/~eilers>

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DE SANTA CATARINA, 88040-900 FLORIANÓPOLIS SC, BRAZIL

E-mail address: exel@mtm.ufsc.br

URL: <http://www.mtm.ufsc.br/~exel>