

STRONGLY REPRESENTABLE ATOM STRUCTURES OF RELATION ALGEBRAS

ROBIN HIRSCH AND IAN HODKINSON

(Communicated by Carl G. Jockusch, Jr.)

ABSTRACT. A relation algebra atom structure α is said to be *strongly representable* if all atomic relation algebras with that atom structure are representable. This is equivalent to saying that the complex algebra $\mathfrak{Cm}\alpha$ is a representable relation algebra. We show that the class of all strongly representable relation algebra atom structures is not closed under ultraproducts and is therefore not elementary. This answers a question of Maddux (1982).

Our proof is based on the following construction. From an arbitrary undirected, loop-free graph Γ , we construct a relation algebra atom structure $\alpha(\Gamma)$ and prove, for infinite Γ , that $\alpha(\Gamma)$ is strongly representable if and only if the chromatic number of Γ is infinite. A construction of Erdős shows that there are graphs Γ_r ($r < \omega$) with infinite chromatic number, with a non-principal ultraproduct $\prod_D \Gamma_r$ whose chromatic number is just two. It follows that $\alpha(\Gamma_r)$ is strongly representable (each $r < \omega$) but $\prod_D(\alpha(\Gamma_r))$ is not.

1. INTRODUCTION

Representability is difficult to characterise for relation algebras. Monk proved that no finite set of first-order sentences can define the class RRA of representable relation algebras [21]. Various strengthenings of this result have been obtained [13], [1], [25]. Furthermore, the representability problem is known to be undecidable for finite relation algebras [11]. On the other hand, representability is quite easily characterised for boolean algebras: every field of sets is a boolean algebra, and by Stone's theorem, every boolean algebra can be represented as a field of sets.

For atomic relation algebras \mathcal{A} (where every non-zero element is above some minimal, non-zero element or *atom*) the non-boolean operations (identity, converse and composition) are determined by the *atom structure* $At\mathcal{A}$ of the relation algebra. This atom structure tells which atoms lie under the identity, which atoms are converses of each other, and what the composition of any two atoms is. It is possible to recover the behaviour of arbitrary elements of the relation algebra from the atom structure, using the fact that converse and composition are completely additive. The only thing the atom structure does not determine is the boolean

Received by the editors October 20, 2000 and, in revised form, November 27, 2000.

2000 *Mathematics Subject Classification*. Primary 03G15; Secondary 03C05, 05C80.

Key words and phrases. Elementary class, complex algebra, relation algebra, representation.

This research was partially supported by UK EPSRC grants GR/L85961, GR/K54946, GR/L85978. Thanks to Rob Goldblatt for valuable additions and suggestions, and Imre Leader for a helpful discussion about the graph-theoretic aspects. Thanks also to the referee for suggesting useful improvements to the paper.

structure: it is possible to have two atomic relation algebras with the same atom structure but certain suprema of sets of atoms are present in one yet not in the other.

The fact that representability is so difficult to pin down for relation algebras but so easy with boolean algebras, together with the informal equation ‘relation algebra = boolean algebra + atom structure’ just established, might lead one to conclude that the atom structure alone determines whether an atomic relation algebra is representable or not. However, this turns out to be false. In [12], two atomic relation algebras with the same atom structure were constructed: one is representable while the other is not. (See §6.1 below for a simple proof.)

So there are two notions of representability for relation algebra atom structures: an atom structure is *weakly representable* if at least one atomic representable relation algebra has that atom structure, and it is *strongly representable* if every atomic relation algebra with that atom structure is representable. Since representability is preserved under subalgebras, and every atomic relation algebra embeds into the complex algebra of its atom structure (the relation algebra with this atom structure whose domain is the full power set of the set of atoms), it is quite easy to see that an atom structure is strongly representable if and only if its complex algebra is representable. Similarly, there is a minimal atomic relation algebra with a given atom structure, called the *term algebra*: it is the subalgebra of the complex algebra generated by the atoms. An atom structure is weakly representable if and only if its term algebra is representable.

By analogy with RRA, it is of interest to provide intrinsic characterisations of the classes of weakly and strongly representable atom structures. The class WRAS of weakly representable atom structures turns out to be elementary: in [24], Venema showed how to take an equational axiomatisation of RRA (e.g., from [15]) and translate the equations into first-order sentences in the language of atom structures. The resulting sentences hold in an atom structure if and only if the equations hold in its term algebra. This provides a first-order axiomatisation of WRAS (or, in another common notation, ‘*At RRA*’, the class of atom structures of atomic, representable relation algebras).¹ It was remarked in [12] that no finite first-order axiomatisation exists, nor one by a sentence of the finite-variable infinitary logic $L_{\infty\omega}^{\omega}$.

But the class of strongly representable atom structures (SRAS, or ‘*Str RRA*’) seems harder to handle, and indeed, not much seems to be known about this class. The following is known. It is easily seen that a finite atom structure is weakly representable if and only if it is strongly representable. Again, SRAS is not finitely axiomatisable or $L_{\infty\omega}^{\omega}$ -axiomatisable [12]. There is also a connection between SRAS and the class CRAS of *completely representable atom structures* — the atom structures of relation algebras that have *complete* representations, respecting arbitrary suprema wherever they exist in the algebra. Any such relation algebra is atomic and has an ‘atomic’ representation [9, Theorem 7]; such a representation induces a representation of the complex algebra of its atom structure, so the atom structure

¹Venema’s argument was given for an arbitrary completely additive variety \mathbf{V} of boolean algebras with operators. The resulting axiomatisation of *At V* is recursive if the initial equational axiomatisation of \mathbf{V} is recursive.

It is historically of interest that Lyndon followed approximately the reverse procedure to get his equational axiomatisation of RRA [15]. He started from the ‘Lyndon conditions’ of [14] (see §6.3), and, using what he called ‘polarisation’, translated them into equational conditions which, when applied to an algebra, assert roughly that its canonical extension satisfies the Lyndon conditions.

must be strongly representable. However, examples of Maddux [16, Examples 23, p. 154ff] yield strongly representable atom structures that are not completely representable. (See §6.2 for another example.) Using the result of [12] mentioned above, we conclude that $\text{CRAS} \subset \text{SRAS} \subset \text{WRAS}$.

Maddux asked [17, Problem 2.3] whether SRAS is an elementary class. In universal algebra, the standard way of proving that a class is or is not elementary is by the following theorem (see [3, Theorems 4.1.12 and 6.1.15] or [23]).

Theorem 1 (Keisler–Shelah). *A class of structures in some first-order signature is elementary if and only if the class is closed under isomorphism, ultraroots, and ultraproducts.*

So to prove that a class is non-elementary, it suffices to show that it is not closed under either or both ultraproducts and ultraroots. Conversely, it will immediately be asked about any non-elementary class whether it fails to be closed under ultraproducts, ultraroots, or both. Consider for example the class CRAS . In [9, Theorem 24], an atomic relation algebra was constructed that has no complete representation but is elementarily equivalent to one with a complete representation. This proved that the class of relation algebras with complete representations is not elementary. It is not too hard to show that this class is closed under ultraproducts; this can be done quite easily by showing that it is a PC_Δ or *pseudo-elementary* class [3, Exercise 4.1.17, Corollary 6.1.16]. Hence, it is not closed under ultraroots. Since the atom structure of an atomic relation algebra is first-order interpretable in the algebra, the operation of taking the atom structure commutes with taking ultraroots. Hence, CRAS is not closed under ultraroots either, so it is not elementary.

However the reader who wishes to prove that SRAS is not elementary by that method will not get very far.

Theorem 2. *SRAS is closed under ultraroots.²*

Proof. Suppose that α is an atom structure that is not strongly representable. So there is an atomic relation algebra \mathcal{A} with atom structure α that is not representable. If D is any ultrafilter on a set I , then the ultrapower \mathcal{A}^I/D is an atomic relation algebra with atom structure α^I/D . As \mathcal{A} is elementarily equivalent to \mathcal{A}^I/D , and RRA is a variety, \mathcal{A}^I/D is not representable. Hence $\alpha^I/D = \text{At}(\mathcal{A}^I/D)$ is not strongly representable. \square

So we might hope to prove that SRAS is an elementary class by showing that it is closed under ultraproducts, as follows. Let α_i ($i \in I$) be strongly representable atom structures. This means that the complex algebras $\mathfrak{Cm}(\alpha_i)$ are representable. Since RRA is elementary, it follows that any ultraproduct $\prod_D \mathfrak{Cm}(\alpha_i)$ is also representable. However, the proof may not be completed, because $\prod_D \mathfrak{Cm}(\alpha_i)$ may not be isomorphic to $\mathfrak{Cm}(\prod_D \alpha_i)$ so we cannot conclude that $\prod_D \alpha_i$ is strongly representable.

In this paper, we provide a negative answer to Maddux’s question, proving (in Corollary 13) the following:

Theorem 3. *SRAS is not an elementary class.*

²A generalisation of this result to arbitrary varieties of Boolean algebras with operators is proved in [7, Theorem 3.8.1(1)].

We proceed by constructing a sequence of strongly representable atom structures α_r ($r < \omega$) whose ultraproduct $\prod_D \alpha_r$ is not strongly representable. The construction makes use of a result of Erdős [5], which provides finite graphs of arbitrarily high chromatic number and girth. By taking disjoint unions of them, we obtain an infinite graph Γ_r , for each $r < \omega$, that cannot be coloured by any finite number of colours but whose ‘small’ subgraphs of size $< r$ have no cycles and so can be coloured by just two colours. It follows that a non-principal ultraproduct of the Γ_r has no cycles at all and is therefore 2-colourable.

From a graph Γ , we construct an atom structure $\alpha(\Gamma)$ and show, for infinite Γ , that $\alpha(\Gamma)$ is strongly representable if and only if there is no finite colouring of Γ . So for each $r < \omega$, $\alpha(\Gamma_r)$ is strongly representable but $\prod_D \alpha(\Gamma_r)$ is not, for any non-principal ultrafilter D over ω .

As is often the case, the use of ultraproducts can be easily replaced by first-order compactness.

By [7, Theorem 3.8.4], SRAS is elementary iff it is closed under elementary equivalence, iff it is closed under ultrapowers, iff it is closed under ultraproducts. Hence, SRAS has none of these closure properties.

We assume a basic knowledge of relation algebras, as can be found in [19] for example. To make the paper self-contained, we recall some elementary definitions and results in section 3.

2. GRAPHS AND COLOURINGS

Let $\Gamma = (V, E)$ be an undirected graph ($V \neq \emptyset$ is the set of vertices or nodes, and E , the set of edges, is an irreflexive, symmetric binary relation on V). Let C be a non-empty set (of ‘colours’). $X \subseteq V$ is said to be an *independent set* if $(x, y) \notin E$ for all $x, y \in X$. A function $f : V \rightarrow C$ is called a *C-colouring* of Γ if $(v, w) \in E \Rightarrow f(v) \neq f(w)$. The *chromatic number* of Γ , denoted $\chi(\Gamma)$, is the size of the smallest finite set C such that there is a C -colouring of Γ , if such a set exists, and $\chi(\Gamma) = \infty$ otherwise. A *cycle* in Γ is a finite sequence $\gamma = \langle v_0, v_1, \dots, v_{k-1} \rangle$ of distinct nodes (some $k \geq 3$) such that $(v_0, v_1), \dots, (v_{k-2}, v_{k-1}), (v_{k-1}, v_0) \in E$. The *length* of such a cycle is k . The *girth* of Γ , denoted $g(\Gamma)$, is the length of the shortest cycle in Γ if Γ has any cycles, and $g(\Gamma) = \infty$ if not.

Theorem 4 (Erdős, [5]). *For all $r < \omega$, there exists a finite graph G_r with $\chi(G_r) > r$ and $g(G_r) > r$.*

A recent presentation of Erdős’ proof of this theorem can be found in [4, Theorem 11.2.2]. Erdős’ proof was radical and was one of the earliest results using the probabilistic method. The idea is to construct a graph G on n vertices by randomly choosing to include an edge between two distinct vertices with probability p and repeating this random choice independently for each pair of distinct vertices in the graph. For $p > \frac{6 \ln n}{n}$ the probability that G contains an independent set of size $\frac{n}{2r}$ tends to zero as n tends to infinity.

If $p < n^{\frac{1}{r}-1}$, then although there may be cycles of length r or less (short cycles) in G , we find that the probability that there are at least $\frac{n}{2}$ short cycles also tends to zero as n tends to infinity. For large n , it is possible to find a value of p with $\frac{6 \ln n}{n} < p < n^{\frac{1}{r}-1}$. With such a value for p , for sufficiently large n , the probability that G has no independent set of size $\frac{n}{2r}$ and fewer than $\frac{n}{2}$ short cycles is strictly positive. It follows that a graph G with both properties exists.

From each of the short cycles of such a graph G we delete a single node to obtain a graph H . Clearly, H has no short cycles, so $g(H) > r$. Also, $|H| \geq \frac{n}{2}$ and H has no independent set of size $\frac{n}{2r}$. But for any r -colouring of H there must be a set of at least $\frac{|H|}{r} \geq \frac{n}{2r}$ nodes, all with the same colour. There are no independent sets in H of this size, so there cannot be such an r -colouring after all. Hence $\chi(H) > r$.

Definition 5. For $r < \omega$, let Γ_r be a disjoint union $\bigcup_{r \leq s < \omega} G_s$, where G_s is a graph, as in Theorem 4, with $\chi(G_s) > s$ and $g(G_s) > s$.

Corollary 6. For all $r < \omega$, Γ_r is an infinite graph with $g(\Gamma_r) > r$ and $\chi(\Gamma_r) = \infty$. Furthermore, if D is any non-principal ultrafilter over ω , we have $\chi(\prod_D \Gamma_r) = 2$.

Proof. The first part of the corollary is clear from the definition of Γ_r . For the second part, since the girth of Γ_r is more than r , Γ_r does not satisfy the following sentence σ_i :

$$\exists x_0 \dots \exists x_{i-1} \left(\bigwedge_{j < k < i} x_j \neq x_k \wedge \bigwedge_{j < i-1} E(x_j, x_{j+1}) \wedge E(x_{i-1}, x_0) \right)$$

for any i with $3 \leq i \leq r$. Hence $\Gamma_r \models \neg \sigma_i$, for $3 \leq i \leq r$. So let Γ be any non-principal ultraproduct of the Γ_r . By Loś' theorem, $\Gamma \models \neg \sigma_i$ for all finite $i \geq 3$. Thus, Γ has no cycles, which implies that Γ can be coloured with just two colours. □

Corollary 7. The class $\mathbf{N} = \{\Gamma : \chi(\Gamma) = \infty\}$ is not elementary.

Proof. By Corollary 6, $\Gamma_r \in \mathbf{N}$, for $r < \omega$, and $\prod_D \Gamma_r \notin \mathbf{N}$. This shows that \mathbf{N} is not closed under ultraproducts and cannot be elementary. □

It is easily seen that \mathbf{N} is closed under elementary equivalence, so (equivalently) it is closed under ultraroots and ultrapowers. The complement $\{\text{graphs } \Gamma : \chi(\Gamma) < \infty\}$ of \mathbf{N} is easily seen not to be closed under ultraproducts (e.g., of complete graphs of arbitrarily large finite size), so it is not elementary either.

3. ATOM STRUCTURES, NETWORKS, AND GAMES

We recall some elementary definitions and results on relation algebra atom structures. This material originated in [14] and appeared in the form given below in [17, Theorem 2.2] and [19, Theorem 37]. A *relation algebra atom structure* $\alpha = \langle A, Id, \smile, C \rangle$ consists of a non-empty set A (of atoms), a unary predicate Id over A (the identity atoms), a unary function $\smile : A \rightarrow A$, and a ternary relation C satisfying, for all $a, b, c, d, g \in A$,

Identity: $a = b \iff \exists e \in A (Id(e) \wedge C(a, e, b))$,

Peircean law: $C(a, b, c) \Rightarrow (C(\check{a}, c, b) \wedge C(c, \check{b}, a))$,

Associativity: $C(a, b, c) \wedge C(c, d, g) \Rightarrow \exists f \in A (C(a, f, g) \wedge C(b, d, f))$.

If $\mathcal{A} = \langle A, 0, 1, +, -, 1', \smile, ; \rangle$ is any atomic relation algebra, we can extract an atom structure

$$At(\mathcal{A}) = \langle \{a \in A : \forall x (x < a \leftrightarrow x = 0)\}, Id, \smile, C \rangle$$

whose domain is the set of atoms of \mathcal{A} , the identity $Id = \{\text{atoms } e : e \leq 1'\}$, converse is the restriction of \smile to the atoms of \mathcal{A} (the converse of a relation algebra

atom can be proved to be an atom), and the ternary relation C is defined by $C(a, b, c) \iff a ; b \geq c$, for all atoms a, b, c .

Conversely, from an atom structure $\alpha = \langle A, Id, \checkmark, C \rangle$ it is possible to construct a relation algebra $\mathfrak{Cm}(\alpha)$ whose domain is the power set of A , with boolean operations \cup and complement in A , and where the non-boolean operators are defined by

$$\begin{aligned} 1' &= Id, \\ \checkmark &= \{\check{a} : a \in r\}, \\ r ; s &= \{c : \exists a \in r, \exists b \in s, C(a, b, c)\}. \end{aligned}$$

$\mathfrak{Cm}(\alpha)$ is called the (full) complex algebra over α . In the complex algebra, it is convenient to identify a set with just one atom in it with the atom itself. Thus, we write $a_1 ; a_2$ instead of the more accurate $\{a_1\} ; \{a_2\}$, for $a_1, a_2 \in A$.

The set of forbidden triples of α is $(A \times A \times A) \setminus C$. Listing the forbidden triples is often a useful way of specifying the composition part of an atom structure.

An atom structure α is strongly representable if every atomic relation algebra whose atom structure is isomorphic to α is representable. Equivalently, $\mathfrak{Cm}(\alpha)$ is representable.

Let \mathcal{A} be a relation algebra. An \mathcal{A} -labelled graph $N = (N_1, N_2)$ consists of a set N_1 of nodes and a function $N_2 : N_1 \times N_1 \rightarrow \mathcal{A}$. An \mathcal{A} -labelled graph $N = (N_1, N_2)$ is called an (\mathcal{A} -)network [10, §3.4] if for all nodes $x, y, z \in N_1$:

- $N_2(x, x) \leq 1'$,
- $N_2(x, y) \cdot (N_2(x, z) ; N_2(z, y)) \neq 0$.

Often we drop the subscripts and use N to denote a labelled graph or network, its set of nodes, and the labelling function, distinguishing the cases by context. So we write $n \in N$ to mean that n is a node of N , and $|N|$ for the cardinality of the set of nodes. In case of possible confusion we may write $nodes(N)$ for the set N_1 of nodes. Let M and N be \mathcal{A} -labelled graphs. We say that M is a refinement of N , and we write $M \supseteq N$, if $nodes(M) \supseteq nodes(N)$ and for all $x, y \in nodes(N)$ we have $M(x, y) \leq N(x, y)$. The idea is that M carries more information than N .

To test if a relation algebra \mathcal{A} is representable, we define a two-player game $G_k(\mathcal{A})$ (for any $k < \omega$) as in [10, §9.1], played over labelled graphs. The players are called \forall and \exists . They construct a sequence

$$N_0 \subseteq N_1 \subseteq N_2 \subseteq \dots \subseteq N_k$$

of labelled graphs as follows.

Initial round: In the initial round (round 0), \forall picks any non-zero element a of \mathcal{A} , and \exists must respond with the two-node network N_0 with nodes x_0, y_0 , say, and labelling defined by $N_0(x_0, y_0) = a$, $N_0(y_0, x_0) = \check{a}$, and $N_0(x_0, x_0) = N_0(y_0, y_0) = 1'$.

In round $t + 1$, for $t < k$, suppose N_t has just been played. \forall picks nodes $x, y \in N_t$ and elements $a, b \in \mathcal{A}$. \exists responds by playing a labelled graph $N_{t+1} \supseteq N_t$ such that $nodes(N_{t+1}) = nodes(N_t) \cup \{z\}$ for some new $z \notin N_t$ and satisfying either of the following two conditions, at her choice:

Reject: $N_{t+1}(x, y) = N_t(x, y) \cdot -(a ; b)$, $N_{t+1}(p, q) = N_t(p, q)$ for all $p, q \in N_t$ with $(p, q) \neq (x, y)$, $N_{t+1}(z, z) = 1'$, and for all edges (p, q) of N_{t+1} not yet mentioned, $N_{t+1}(p, q) = 1$.

Accept: $N_{t+1}(x, y) = N_t(x, y) \cdot (a; b)$, $N_{t+1}(p, q) = N_t(p, q)$ for all $p, q \in N_t$ with $(p, q) \neq (x, y)$, $N_{t+1}(x, z) = a$, $N_{t+1}(z, z) = 1'$, $N_{t+1}(z, y) = b$, and for all edges (p, q) of N_{t+1} not yet mentioned, $N_{t+1}(p, q) = 1$.

If for any $t \leq k$, the labelled graph N_t is not a network (it fails one of the two network conditions), then \forall wins. If every labelled graph in the play is a network, then \exists wins. It can be seen by the definition of the game that the labelled graphs all satisfy the first network condition. So \forall wins if, in some round, a graph N_t is played that contains nodes x, y, z such that $(N_t(x, z); N_t(z, y)) \cdot N_t(x, y) = 0$. Such a triangle (x, y, z) is said to be *inconsistent*. If $(N_t(x, z); N_t(z, y)) \cdot N_t(x, y) \neq 0$, we say that the triangle (x, y, z) is *consistent*.

Theorem 8. *Let \mathcal{A} be a relation algebra. \mathcal{A} is representable if and only if \exists has a winning strategy in each game $G_k(\mathcal{A})$, for $k < \omega$.*

Proof. [10, Proposition 23] proves this for countable \mathcal{A} . For arbitrary \mathcal{A} , [10, Theorem 24(2)] shows that \exists 's having a winning strategy in $G_k(\mathcal{A})$ is an elementary property, defined by a first-order sentence ϕ_k . Also, RRA is an elementary class (a variety). Hence, if \mathcal{A} is any relation algebra and \mathcal{C} a countable elementary subalgebra of \mathcal{A} , the following are equivalent: $\mathcal{A} \in \text{RRA}$; $\mathcal{C} \in \text{RRA}$; \exists has a winning strategy in $G_k(\mathcal{C})$ for all k ; $\mathcal{C} \models \phi_k$ for all k ; $\mathcal{A} \models \phi_k$ for all k ; \exists has a winning strategy in $G_k(\mathcal{A})$ for all k . □

Lemma 9. *Let $k < \omega$ and let \mathcal{A} be a relation algebra. If \forall has a winning strategy in the game $G_k(\mathcal{A})$, then there is a non-zero element $a_0 \in \mathcal{A}$ and a finite set W of pairs of elements of \mathcal{A} such that his winning strategy selects a_0 in round 0 and in later rounds only ever directs him to choose elements from W .*

Proof. Let σ be a fixed (deterministic) winning strategy for \forall . σ determines his initial choice a_0 . In each round, \exists has at most two choices for her moves. Therefore, there are at most 2^k possible plays of $G_k(\mathcal{A})$ in which \forall uses σ . In each round, \forall is directed to choose a pair of elements in his moves. So σ can direct him to choose at most $k \times 2^k$ different pairs. □

4. THE CONSTRUCTION

Let Γ be a graph. We define a relation algebra atom structure $\alpha(\Gamma)$ with atoms $1'$ (the sole identity atom) and r_i, b_i, g_i for $i \in \Gamma$. The non-identity atoms r_i, b_i, g_i are regarded as coloured red, blue, and green, respectively. All atoms are self-converse (i.e., $\check{a} = a$). To define composition on atoms, we list the *forbidden triples* of atoms (a, b, c) : those such that $\neg C(a, b, c)$, or equivalently, $(a; b) \cdot c = 0$. The forbidden triples are:

- (A) $(1', a, b), (a, 1', b), (a, b, 1')$ where $a \neq b$,
- (B) $(r_i, r_j, r_k), (b_i, b_j, b_k), (g_i, g_j, g_k)$ where $\{i, j, k\} \subseteq \Gamma$ is an independent set.

It is easily checked that $\alpha(\Gamma)$ meets the conditions of being a relation algebra atom structure. Therefore, $\mathfrak{Cm}(\alpha(\Gamma))$ is a relation algebra, known colloquially as a kind of ‘Monk algebra’ (cf. [22]).

For a graph Γ and $X \subseteq \Gamma$, we define $R_X, B_X, G_X \in \mathfrak{Cm}(\alpha(\Gamma))$ to be the sets of all red, blue, and green atoms, respectively, with indices in X : that is, $R_X = \{r_i : i \in X\}$, etc. A non-zero element a of $\mathfrak{Cm}(\alpha(\Gamma))$ is said to be *monochromatic* if $a \leq 1'$, $a \leq R_\Gamma$, $a \leq B_\Gamma$, or $a \leq G_\Gamma$.

Theorem 10. *If Γ is infinite and $\chi(\Gamma) < \infty$, then $\alpha(\Gamma)$ is not strongly representable.*

Proof. We must show that $\mathfrak{Cm}(\alpha(\Gamma))$ is not representable. Assume for contradiction that $h : \mathfrak{Cm}(\alpha(\Gamma)) \rightarrow \mathcal{A}$ is an isomorphism from $\mathfrak{Cm}(\alpha(\Gamma))$ into a proper relation algebra \mathcal{A} with base set X . We may assume that $h(1) = X \times X$.³ Each $h(a)$ ($a \in \mathfrak{Cm}(\alpha(\Gamma))$) is a binary relation on X , and h respects the relation algebra operations. In particular,

- (1) if $a, b \in \mathfrak{Cm}(\alpha(\Gamma))$, then $h(a + b) = h(a) \cup h(b)$,
- (2) if $x, y \in X$ and $(x, y) \in h(1')$, then $x = y$,
- (3) if $x_0, x_1, x_2 \in X$, $a, b, c \in \mathfrak{Cm}(\alpha(\Gamma))$, and $(x_0, x_1) \in h(a)$, $(x_1, x_2) \in h(b)$, and $(x_0, x_2) \in h(c)$, then $(a ; b) \cdot c \neq 0$.

We know that Γ has a finite colouring, so partition the nodes of Γ into sets C_j ($j < n$ for some finite n) such that there are no edges within any C_j . Let $\Pi = \{1', R_{C_j}, B_{C_j}, G_{C_j} : j < n\}$. Then $\sum \Pi = 1$ in $\mathfrak{Cm}(\alpha(\Gamma))$. As Π is finite, repeated use of (1) shows that for any $x, y \in X$ there is $a \in \Pi$ with $(x, y) \in h(a)$.

Since $\mathfrak{Cm}(\alpha(\Gamma))$ is infinite, X must also be infinite. By Ramsey's theorem, there are distinct $x_i \in X$ ($i < \omega$) and $a \in \Pi$ such that $(x_i, x_j) \in h(a)$ for all $i < j < \omega$. By (2) above, $a \neq 1'$. By (3), $(a ; a) \cdot a \neq 0$. But because the C_j are independent sets and a is monochromatic, it is easily checked using rule (B) of the definition of $\alpha(\Gamma)$ that $(a ; a) \cdot a = 0$. This contradiction completes the proof that $\mathfrak{Cm}(\alpha(\Gamma))$ is not representable. \square

Theorem 11. *If $\chi(\Gamma) = \infty$, then $\alpha(\Gamma)$ is strongly representable.*

Proof. We show that $\alpha(\Gamma)$ is strongly representable by showing that \exists has a winning strategy in $G_k(\mathfrak{Cm}(\alpha(\Gamma)))$ for any finite k .⁴ Strong representability then follows by Theorem 8.

Write α for $\alpha(\Gamma)$. Suppose, for contradiction, that \exists does not have a winning strategy in $G_k(\mathfrak{Cm} \alpha)$, for some finite k . Since finite-length games are determined [6], it follows that \forall must have a winning strategy in this game.⁵ By Lemma 9, there is non-zero $a_0 \in \mathfrak{Cm} \alpha$ and a finite set W of pairs of elements of $\mathfrak{Cm} \alpha$ such that his winning strategy picks a_0 in round 0 and in later rounds only ever directs him to choose pairs of elements in W .

We define a finite boolean subalgebra \mathcal{B} of the boolean reduct $bool(\mathfrak{Cm} \alpha)$ of $\mathfrak{Cm} \alpha$ in three steps. Start with the finite set

$$B_0 = \{1', a_0, R_\Gamma, B_\Gamma, G_\Gamma\} \cup \{a, b, a ; b : (a, b) \in W\}.$$

First let B_1 be the finite boolean subalgebra of $bool(\mathfrak{Cm} \alpha)$ generated by B_0 (the set of all unions of intersections of members of B_0 and complements of members of B_0).

³Note that $\mathfrak{Cm}(\alpha(\Gamma))$ is a *simple* relation algebra. Hence, if it has any representation, then it has a *square* representation h satisfying $h(1) = X \times X$, where X is the base of the representation.

⁴ $\alpha(\Gamma)$ need not be *completely* representable, nor need it satisfy the Lyndon conditions of [14] (see §§6.2–6.3 for specific counter-examples and Theorem 14 for links between these notions). In [10, §5], we defined a game $G_k^{at}(\mathcal{A})$, for an atomic relation algebra \mathcal{A} . This game is similar to $G_k(\mathcal{A})$ but is played over atomic networks (networks in which each edge is labelled by an atom of \mathcal{A}). If \mathcal{A} has a complete representation, then \exists has a winning strategy in $G_k^{at}(\mathcal{A})$ for all finite k (though the converse is false), and the latter condition is equivalent to the Lyndon conditions. So G_k^{at} , though easier to handle than G_k , is not appropriate for Theorem 11.

⁵It is more convenient for us to deduce a contradiction from a winning strategy for \forall in this finite-length game than to prove directly that \exists has a winning strategy, because we can then apply Lemma 9 and restrict our attention to just a finite set of elements of the complex algebra.

Second, let $B_2 = B_1 \cup \{R_X, B_X, G_X : R_X \in B_1 \text{ or } B_X \in B_1 \text{ or } G_X \in B_1\}$ (copy monochromatic elements to all colours). Finally, let \mathcal{B} be the boolean subalgebra of $bool(\mathfrak{Cm} \alpha)$ generated by B_2 . Observe that

- $1', a_0 \in \mathcal{B}$, and if $(a, b) \in W$, then $a, b, (a; b) \in \mathcal{B}$,
- each atom of \mathcal{B} is monochromatic,
- for any $X \subseteq \Gamma$, the following are equivalent: $R_X \in \mathcal{B}, B_X \in \mathcal{B}, G_X \in \mathcal{B}$.

Write $At \mathcal{B}$ for the set of atoms of \mathcal{B} . A subset $X \subseteq \Gamma$ is said to be *small* if $R_X \in At \mathcal{B}$. Clearly, the small sets form a finite partition of Γ . Since there is no finite colouring of Γ , there must be some small set $S \subseteq \Gamma$ such that (i, j) is an edge of Γ for some $i, j \in S$. Then by rule (B) of the definition of $\alpha(\Gamma)$, (r_i, r_j, r_l) , (r_i, r_j, b_l) , and (r_i, r_j, g_l) are not forbidden triples of atoms of α for any $l \in \Gamma$. The same holds with b_i, b_j and with g_i, g_j . Hence, by definition of complex algebras,

$$(3) \quad C_S; C_S = 1 \text{ in } \mathfrak{Cm} \alpha, \text{ whenever } C \text{ is } R, B, \text{ or } G.$$

To obtain a contradiction, we will show how \exists can win $G_k(\mathfrak{Cm} \alpha)$ when \forall uses his supposedly winning strategy. To help her win, she will maintain an auxiliary network M_t (for each $t \leq k$) satisfying the following conditions:

- $M_t(e) \in At \mathcal{B}$ for each edge e of M_t ,
- if $x, y \in M_t$, then $M_t(x, y) = 1'$ if and only if $x = y$,
- there is a map $\prime : nodes(N_t) \rightarrow nodes(M_t)$ such that $M_t(x', y') \leq N_t(x, y)$ for all $x, y \in nodes(N_t)$,

where N_t is the labelled graph in play after round t .

For the initial move ($t = 0$), \forall chooses the non-zero element $a_0 \in \mathcal{B}$. \exists must respond with a two-node network N_0 with distinct nodes x_0, y_0 , say, with $N_0(x_0, y_0) = N_0(y_0, x_0) = a_0$ ($= \check{a}_0$) and $N_0(x_0, x_0) = N_0(y_0, y_0) = 1'$. To define M_0 , she chooses $a^* \in At \mathcal{B}$ with $a^* \leq a_0$. If $a^* = 1'$, she defines M_0 to be the network with a single node, say x_0 , with $M_0(x_0, x_0) = 1'$; she defines $x'_0 = y'_0 = x_0$. Otherwise, $a^* \leq -1'$, and \exists defines M_0 to have nodes x_0, y_0 only, with $M_0(x_0, y_0) = M_0(y_0, x_0) = a^*$, and $M_0(x_0, x_0) = M_0(y_0, y_0) = 1'$. She lets $x'_0 = x_0$ and $y'_0 = y_0$. This all meets the conditions above.

Let $t < k$ and assume inductively that \exists has defined M_t for N_t . We will show how \exists can respond in the next round of the game and how she can construct a new network M_{t+1} meeting the above conditions with respect to N_{t+1} . \forall plays the round by choosing nodes $x, y \in N_t$ and elements $a, b \in \mathfrak{Cm} \alpha$ according to his strategy. Note that $a, b, a; b \in \mathcal{B}$. \exists uses the label $M_t(x', y')$ to calculate her response. If $M_t(x', y') \leq -(a; b)$, then she rejects his move and plays N_{t+1} as specified in the definition in section 3 of the game G_k , with a new node z . She defines $M_{t+1} = M_t$, and defines z' to be any node of M_t (for definiteness, say $z' = x$). This preserves the requirements on M, N .

Otherwise, since $M_t(x', y') \in At \mathcal{B}$, we have $M_t(x', y') \leq a; b$. \exists then accepts \forall 's move by playing a labelled graph N_{t+1} with a new node z and labelling it as specified earlier. She now defines M_{t+1} meeting the conditions, as follows.

If there is already a node $p \in M_t$ with $M_t(x', p) \leq a$ and $M_t(p, y') \leq b$, then \exists can set $M_{t+1} = M_t$ and $z' = p$. This meets the required conditions on N_{t+1}, M_{t+1} .

Otherwise, since $a; b \geq M_t(x', y') \in At \mathcal{B}$ and $a, b \in \mathcal{B}$, by additivity of composition in $\mathfrak{Cm} \alpha$ there must be $a^*, b^* \in At \mathcal{B}$ with $a^* \leq a, b^* \leq b$, and $(a^*; b^*) \cdot M_t(x', y') \neq 0$. \exists chooses such a^*, b^* ; it can be checked using standard relation algebra properties that $a^*, b^* \neq 1'$, and that if $x' = y'$, then $a^* = b^*$. Note that a^*, b^*

are monochromatic. Choose C from R, B, G of different colour from the colours of (atoms in) a^*, b^* . \exists now refines M_t to $M_{t+1} \supseteq M_t$ by adding a new node, defining z' to be that new node, and labelling edges of M_{t+1} with atoms of \mathcal{B} as follows:

- $M_{t+1}(p, q) = M_t(p, q)$ for all $p, q \in M_t$.
- $M_{t+1}(x', z') = M_{t+1}(z', x') = a^*$, $M_{t+1}(z', z') = 1'$, and $M_{t+1}(z', y') = M_{t+1}(y', z') = b^*$. This is well-defined if $x' = y'$, as $a^* = b^*$ in that case.
- For $p \in nodes(M_t) \setminus \{x', y'\}$, \exists labels $M_{t+1}(p, z') = M_{t+1}(z', p) = C_S$.

We claim that every triangle (p, q, r) of nodes of the labelled graph M_{t+1} is consistent — $M_{t+1}(p, q) \cdot (M_{t+1}(p, r); M_{t+1}(r, q)) \neq 0$ in $\mathfrak{Cm} \alpha$. If $p, q, r \in M_t$, this is clear inductively, if $|\{p, q, r\}| < 3$, it is trivial, and by definition of α and $\mathfrak{Cm} \alpha$, the order of p, q, r is not significant, so we need only consider triangles of distinct nodes of the form (p, q, z') for $p, q \in M_t$. We already know (x', y', z') to be consistent, since $(a^*; b^*) \cdot M_t(x', y') \neq 0$. The labels in any triangle of the form (x', q, z') with $q \neq x', y'$ are c, C_S, a^* for some $c \in At \mathcal{B} \setminus \{1'\}$; because C is not the colour of a^* , the triangle must be consistent. The case of (y', q, z') is similar. Finally, if (p, q, z') is a triangle with $p, q \in M_t \setminus \{x, y\}$, we have $M_{t+1}(p, z') = M_{t+1}(q, z') = C_S$ and $M_{t+1}(p, q) = c$ for some $c \in At \mathcal{B}$. By (3) above, $c \leq C_S; C_S$, whence (p, q, z') is consistent. This proves the claim.

By the claim, M_{t+1} is a network. We have arranged that every edge of M_{t+1} is labelled by an element of $At \mathcal{B}$ beneath the corresponding label in N_{t+1} . Thus, the inductive conditions on M_t are maintained for another round.

It follows that each labelled graph N_t ($t \leq k$) is a network. To see this, let $x, y, z \in N_t$. Then $N_t(x, z); N_t(z, y) \cdot N_t(x, y) \geq M_t(x', z'); M_t(z', y') \cdot M_t(x', y') \neq 0$. Thus, \exists never loses, which is a contradiction. □

The proof yields the following corollary:

Corollary 12. *Let Γ be an infinite graph and \mathcal{A} a subalgebra of $\mathfrak{Cm}(\alpha(\Gamma))$ such that $\alpha(\Gamma) \subseteq \mathcal{A}$ and $R_\Gamma, B_\Gamma, G_\Gamma \in \mathcal{A}$. Suppose that whenever E_0, \dots, E_n is a finite partition of Γ , C is R, B , or G , and $C_{E_i} \in \mathcal{A}$ for all $i \leq n$, then some E_i is not independent. Then \mathcal{A} is representable.*

5. SRAS IS NOT ELEMENTARY

Theorem 3 follows from:

Corollary 13. *SRAS is not closed under ultraproducts, so it is not an elementary class.*

Proof. For $r < \omega$ let Γ_r be an infinite graph, as in Definition 5, with $g(\Gamma_r) > r$ and $\chi(\Gamma_r) = \infty$ (Corollary 6). Then the atom structure $\alpha(\Gamma_r)$ is strongly representable by Theorem 11. Consider a non-principal ultraproduct $\prod_D \alpha(\Gamma_r)$ over ω . It is easily seen that this is isomorphic to $\alpha(\prod_D \Gamma_r)$. By Corollary 6, $\prod_D \Gamma_r$ is two-colourable, so by Theorem 10, $\alpha(\prod_D \Gamma_r)$ is not strongly representable. Hence SRAS is not closed under ultraproducts, and, by Theorem 1, not elementary. □

6. INCLUSIONS BETWEEN CLASSES

We end with some remarks and open questions concerning classes of atom structures related to SRAS.

6.1. Weakly but not strongly representable atom structures. Since $\text{SRAS} \subseteq \text{WRAS}$ and WRAS is elementary, Corollary 13 gives another proof of a result in [12, §6.2] that $\text{SRAS} \neq \text{WRAS}$. Indeed, regarding the integers \mathbb{Z} as a graph with edges between consecutive integers, we claim that $\alpha(\mathbb{Z}) \in \text{WRAS} \setminus \text{SRAS}$ (see §4 for $\alpha(\mathbb{Z})$). For this, first note that $\alpha(\mathbb{Z}) \notin \text{SRAS}$ by Theorem 10, since $\chi(\mathbb{Z}) = 2$. Now, for the other half, observe that the term algebra τ of $\alpha(\mathbb{Z})$ is the subalgebra of $\mathfrak{Cm}(\alpha(\mathbb{Z}))$ consisting of all elements a such that for every colour C (R, B, or G), $a \cap C_{\mathbb{Z}}$ is finite or co-finite in $C_{\mathbb{Z}}$. So if E_0, \dots, E_n forms a finite partition of \mathbb{Z} , and $C_{E_0}, \dots, C_{E_n} \in \tau$, then one of the E_i is cofinite in \mathbb{Z} and so contains an edge of \mathbb{Z} . By Corollary 12, τ is representable, so $\alpha(\mathbb{Z}) \in \text{WRAS}$.⁶

[12] also gives ‘ n -dimensional’ results of this kind, for the variety RCA_n of representable n -dimensional cylindric algebras. It is shown that $\text{Str RCA}_n \subset \text{At RCA}_n$ for all finite $n \geq 3$ (see §1 above for the notations Str , At). See Problem 1 below. The argument of [12, §6.2] also works for the varieties RA_n and $\text{S}\mathfrak{R}\mathfrak{a}\text{CA}_n$ of relation algebras (see [18, 20] for information about these), showing that $\text{Str RA}_n \subset \text{At RA}_n$ and $\text{Str S}\mathfrak{R}\mathfrak{a}\text{CA}_n \subset \text{At S}\mathfrak{R}\mathfrak{a}\text{CA}_n$ for finite $n \geq 7$. A slightly modified argument also covers the case $n = 6$; for $n \leq 4$ it is false; and the case $n = 5$ remains open. It follows that these varieties are not closed under completions, and are not Sahlqvist-axiomatisable [25].

6.2. Completely representable atom structures. We mentioned in section 1 that atom structures devised by Maddux [16, Examples 23] can be adapted to show that $\text{CRAS} \subset \text{SRAS}$. This can also be proved using Corollary 13. As we said in section 1, $\text{CRAS} \subseteq \text{SRAS}$, and CRAS is pseudo-elementary and so closed under ultraproducts. Since SRAS is not closed under ultraproducts, the inclusion is strict.

Indeed, we can exhibit a specific example: if Γ is a graph with $g(\Gamma) > 15$ and $\chi(\Gamma) = \infty$ (Corollary 6), then $\alpha(\Gamma) \in \text{SRAS} \setminus \text{CRAS}$. We can see this as follows. Theorem 11 shows that $\alpha(\Gamma) \in \text{SRAS}$. To see that $\alpha(\Gamma) \notin \text{CRAS}$, suppose otherwise, for contradiction. A complete representation of a relation algebra with atom structure $\alpha(\Gamma)$ yields an infinite set X and a map $\lambda : X \times X \rightarrow \alpha(\Gamma)$ such that for all $x, y, z \in X$ we have $\lambda(x, y) = 1' \iff x = y$, $\lambda(x, y) = \lambda(y, x)$, and $(\lambda(x, y), \lambda(y, z), \lambda(x, z))$ is a consistent triple in $\alpha(\Gamma)$ (cf. [9, Theorem 7]). By Ramsey’s theorem we may take distinct $x_i \in X$ ($i < 6$) such that for some colour c (i.e., r, b, or g), $\lambda(x_i, x_j) = c_{a_{ij}}$ for some $a_{ij} \in \Gamma$ whenever $i, j < 6$ are distinct. Let Δ be the induced subgraph of Γ with nodes $\{a_{ij} : i < j < 6\}$. Since $|\Delta| \leq 15 < g(\Gamma)$, Δ is 2-colourable and we can partition its nodes into sets D_0, D_1 , neither of which contain any edges of Γ . Since any 2-colouring of the edges of a complete graph of size 6 has a monochromatic triangle, there are distinct $i, j, k < 6$ and $d < 2$ such that $a_{ij}, a_{jk}, a_{ik} \in D_d$. So $\{a_{ij}, a_{jk}, a_{ik}\}$ is an independent set in Γ . But then, $(\lambda(x_i, x_j), \lambda(x_j, x_k), \lambda(x_i, x_k)) = (c_{a_{ij}}, c_{a_{jk}}, c_{a_{ik}})$ is inconsistent, a contradiction.

6.3. Lyndon conditions. Let LC be the class of atom structures of atomic relation algebras satisfying the ‘Lyndon conditions’ (first-order sentences given in [14] and axiomatising the representable relation algebras over finite relation algebras). All quantifiers in the Lyndon conditions are relativised to atoms, so they are essentially first-order conditions on relation algebra atom structures. Hence, LC is elementary. Now an arbitrary atomic relation algebra is elementarily equivalent to a completely

⁶Alternatively, show that \exists has a winning strategy in the game $G_{\omega}^{at}(\tau^+)$ of [10, §5], where τ^+ is the canonical extension of τ .

representable relation algebra (so is itself representable) iff it satisfies all the Lyndon conditions [9, Theorem 20]. Hence, $\text{CRAS} \subseteq \text{LC} \subseteq \text{SRAS}$. Since LC is elementary, and, as seen in section 1 and Corollary 13, CRAS and SRAS are not, both inclusions are strict.

An example of an atom structure in $\text{LC} \setminus \text{CRAS}$ is given in [9, Theorem 21]. For an explicit example of an atom structure in $\text{SRAS} \setminus \text{LC}$, take a graph Γ with $\chi(\Gamma) = \infty$ and $g(\Gamma) > 15$. If n is large enough so that any 3-colouring of the edges of a graph of size n must contain six nodes such that all the edges between the six have the same colour, then the argument of §6.2 shows that $\alpha(\Gamma)$ fails the n th Lyndon condition. By Theorem 11, $\alpha(\Gamma) \in \text{SRAS} \setminus \text{LC}$.

6.4. First-order algebras. For a relation algebra atom structure α , the *first-order algebra over α* is the subalgebra of $\mathfrak{Cm} \alpha$ consisting of all sets of atoms that are first-order definable with parameters in α — that is, sets of the form $\{a \in \alpha : \alpha \models \varphi(a, \bar{b})\}$ for some first-order formula $\varphi(x, \bar{y})$ of the signature of α and some tuple \bar{b} of elements of α . It can be checked that these sets indeed form a subalgebra of $\mathfrak{Cm} \alpha$.

Let FO denote the class of relation algebra atom structures whose first-order algebra is representable.⁷ Evidently, $\text{FO} \subseteq \text{SRAS} \subseteq \text{WRAS}$. FO can be shown to be elementary by the argument of [24], or by using Theorem 1; so by Corollary 13, the first inclusion is proper. Indeed, an Ehrenfeucht–Fraïssé game argument will establish that the first-order algebra of the atom structure $\alpha(\mathbb{Z})$ of §6.1 is the same as its term algebra, so that $\alpha(\mathbb{Z}) \in \text{FO} \setminus \text{SRAS}$. To see that the second inclusion is also proper, let Γ consist of ω disjoint copies of the three-node graph with nodes 0, 1, 2, and edges 0–1 and 1–2 only. As Γ has a first-order-definable 2-colouring (nodes can be coloured according to their degree: 1 or 2), and Γ is interpretable in $\alpha(\Gamma)$, the argument of Theorem 10 shows that the first-order algebra over $\alpha(\Gamma)$ is not representable, so $\alpha(\Gamma) \notin \text{FO}$. However, the term algebra over $\alpha(\Gamma)$ consists of all subsets S of $\alpha(\Gamma)$ such that each of $R_\Gamma \cap S$, $B_\Gamma \cap S$ and $G_\Gamma \cap S$ is finite or cofinite in $R_\Gamma, B_\Gamma, G_\Gamma$, respectively. The term algebra can now be shown to be representable by an argument similar to that of §6.1. So $\alpha(\Gamma) \in \text{WRAS}$.

6.5. Summary of inclusions. We conclude from these remarks that

Theorem 14. $\text{CRAS} \subset \underline{\text{LC}} \subset \text{SRAS} \subset \underline{\text{FO}} \subset \underline{\text{WRAS}}$, the elementary classes being underlined.

We end with a problem.

Problem 1. Let $3 \leq n$. Can the results of the present paper be extended to prove that the class StrRCA_n of strongly representable n -dimensional cylindric algebra atom structures is not elementary? (Cf. [9, Theorem 34] for a proof that the class of completely representable n -dimensional cylindric algebras is not elementary for $n \geq 3$.) Note from [8, Theorem 3.2.11] that with infinitely many dimensions, every semi-simple n -dimensional cylindric algebra is representable, but there is no discriminator so not every cylindric algebra is semi-simple.

REFERENCES

- [1] H. Andréka, *Complexity of equations valid in algebras of relations, Part I: Strong non-finitizability*, Annals of Pure and Applied Logic **89** (1997), 149–209. MR **99b**:03078

⁷We thank Rob Goldblatt for suggesting this example.

- [2] H. Andréka, J. D. Monk, and I. Németi (eds.), *Algebraic logic*, Colloq. Math. Soc. J. Bolyai, no. 54, North-Holland, Amsterdam, 1991, Proc. Conf. Budapest, 1988. MR **92m**:03003
- [3] C. C. Chang and H. J. Keisler, *Model theory*, 3rd ed., North-Holland, Amsterdam, 1990. MR **91c**:03026
- [4] R. Diestel, *Graph theory*, Graduate Texts in Mathematics, vol. 173, Springer-Verlag, Berlin, 1997. CMP 97:12
- [5] P. Erdős, *Graph theory and probability*, Canad. J. Math. **11** (1959), 34–38. MR **21**:876
- [6] D. Gale and F. Stewart, *Infinite games with perfect information*, Contributions to the theory of games II, Annals of mathematical studies **28** (1953), 291–296. MR **14**:999b
- [7] R. Goldblatt, *Varieties of complex algebras*, Annals of Pure and Applied Logic **44** (1989), 173–242. MR **91d**:08005
- [8] L. Henkin, J. D. Monk, and A. Tarski, *Cylindric algebras part II*, North-Holland, 1985. MR **86m**:03095b
- [9] R. Hirsch and I. Hodkinson, *Complete representations in algebraic logic*, Journal of Symbolic Logic **62** (1997), 816–847. MR **98m**:03123
- [10] ———, *Step by step — building representations in algebraic logic*, Journal of Symbolic Logic **62** (1997), 225–279. MR **98g**:03145
- [11] ———, *Representability is not decidable for finite relation algebras*, Trans. Amer. Math. Soc. **353** (2001), 1403–1425. CMP 2001:06
- [12] I. Hodkinson, *Atom structures of cylindric algebras and relation algebras*, Annals of Pure and Applied Logic **89** (1997), 117–148. MR **99c**:03103
- [13] B. Jónsson, *The theory of binary relations*, In Andréka et. al. [2], pp. 245–292. MR **93b**:03114
- [14] R. Lyndon, *The representation of relational algebras*, Annals of Mathematics **51** (1950), 707–729. MR **12**:237a
- [15] ———, *The representation of relation algebras, II*, Annals of Mathematics **63** (1956), 294–307. MR **18**:106f
- [16] R. Maddux, *Topics in relation algebra*, Ph.D. thesis, University of California, Berkeley, 1978.
- [17] ———, *Some varieties containing relation algebras*, Trans. Amer. Math. Soc. **272** (1982), 501–526. MR **84a**:03079
- [18] ———, *A sequent calculus for relation algebras*, Annals of Pure and Applied Logic **25** (1983), 73–101. MR **85h**:03067
- [19] ———, *Introductory course on relation algebras, finite-dimensional cylindric algebras, and their interconnections*, In Andréka et. al. [2], pp. 361–392. MR **93c**:03082
- [20] ———, *Non-finite axiomatizability results for cylindric and relation algebras*, Journal of Symbolic Logic **54** (1989), 951–974. MR **90f**:03099
- [21] J. D. Monk, *On representable relation algebras*, Michigan Mathematics Journal **11** (1964), 207–210. MR **30**:3016
- [22] ———, *Nonfinitizability of classes of representable cylindric algebras*, Journal of Symbolic Logic **34** (1969), 331–343. MR **41**:1517
- [23] S. Shelah, *Every two elementarily equivalent models have isomorphic ultrapowers*, Israel J. Math. **10** (1971), 224–233. MR **45**:6608
- [24] Y. Venema, *Atom structures*, Advances in Modal Logic '96 (M. Kracht, M. de Rijke, H. Wansing, and M. Zakharyashev, eds.), CSLI Publications, Stanford, 1997, pp. 291–305. CMP 99:12
- [25] ———, *Atom structures and Sahlqvist equations*, Algebra Universalis **38** (1997), 185–199. MR **99g**:06019

DEPARTMENT OF COMPUTER SCIENCE, UNIVERSITY COLLEGE, GOWER STREET, LONDON WC1E 6BT, UNITED KINGDOM

E-mail address: r.hirsch@cs.ucl.ac.uk

URL: <http://www.cs.ucl.ac.uk/staff/R.Hirsch/>

DEPARTMENT OF COMPUTING, IMPERIAL COLLEGE, QUEEN'S GATE, LONDON SW7 2BZ, UNITED KINGDOM

E-mail address: imh@doc.ic.ac.uk

URL: <http://www.doc.ic.ac.uk/~imh/>