

SUBSPACES WITH NORMALIZED TIGHT FRAME WAVELETS IN \mathbb{R}

XINGDE DAI, YUANAN DIAO, AND QING GU

(Communicated by David R. Larson)

ABSTRACT. In this paper we investigate the subspaces of $L^2(\mathbb{R})$ which have normalized tight frame wavelets that are defined by set functions on some measurable subsets of \mathbb{R} called Bessel sets. We show that a subspace admitting such a normalized tight frame wavelet falls into a class of subspaces called reducing subspaces. We also consider the subspaces of $L^2(\mathbb{R})$ that are generated by a Bessel set E in a special way. We present some results concerning the relation between a Bessel set E and the corresponding subspace of $L^2(\mathbb{R})$ which either has a normalized tight frame wavelet defined by the set function on E or is generated by E .

§1. INTRODUCTION AND PRELIMINARIES

The concept of frames first appeared in the late 40's and early 50's ([1], [10], [11]). There has recently been renewed interest in frames due to the development of wavelet theory since wavelets and frames are tightly related ([4], [5], [6] and [9]). A sequence of elements $\{x_j\}$ in \mathcal{H} is called a *Bessel sequence* if there exists a positive constant B such that $\sum_j |\langle x, x_j \rangle|^2 \leq B\|x\|^2$ for all $x \in \mathcal{H}$. If in addition, there exists a positive constant A such that $A\|x\|^2 \leq \sum_j |\langle x, x_j \rangle|^2$ for all $x \in \mathcal{H}$, then $\{x_j\}$ is called a frame for \mathcal{H} . The supremum of all such numbers A and the infimum of all such numbers B are called the *frame bounds* of the frame and denoted by A_0 and B_0 respectively. The frame is called a *tight* frame when $A_0 = B_0$ and is called a *normalized tight* frame when $A_0 = B_0 = 1$. The kind of frames that are related to wavelets and of special interest to mathematicians are of the form $\{\psi_{j,k}(x) = 2^{\frac{j}{2}}\psi(2^j x - k) : j, k \in \mathbb{Z}\}$, where $\psi \in L^2(\mathbb{R})$. The function ψ is called a normalized tight frame wavelet (NTF wavelet) if this frame is a normalized tight frame (NTF). Similarly, ψ is called a tight frame wavelet (TF wavelet) if this frame is a tight frame. From an operator theoretic point of view, NTF wavelets are simply the frame vectors for the unitary system $\mathcal{U} = \{D^n T^l : l, n \in \mathbb{Z}\}$, where D and T are the unitary dilation and translation operators on $L^2(\mathbb{R})$ (see [7]) defined by $(Df)(x) = \sqrt{2}f(2x)$ and $(Tf)(x) = f(x-1)$ for any $f \in L^2(\mathbb{R})$. A characterization of the NTF wavelets is obtained in [8].

In this paper, we are mainly interested in the case when ψ is defined by $\hat{\psi} = \frac{1}{\sqrt{2\pi}}\chi_E$, where E is a Lebesgue measurable set with finite measure. We will call E a *normalized tight frame wavelet set* (NTF wavelet set) when ψ is an NTF wavelet.

Received by the editors June 26, 2000 and, in revised form, November 21, 2000.
1991 *Mathematics Subject Classification*. Primary 46N99, 46B28.

©2001 American Mathematical Society

A *tight frame wavelet set* (TF wavelet set) and a *frame wavelet set* are defined similarly. Likewise, E is called a *Bessel set* if $\{\psi_{j,k}(x) : j, k \in \mathbb{Z}\}$ is a Bessel sequence. All these concepts can be applied to the subspaces of $L^2(\mathbb{R})$ (to be formally defined in the next section) and many questions can be asked. A closed subspace X of $L^2(\mathbb{R})$ is called a reducing subspace of D and T (or simply *reducing subspaces* in this paper) if $TX = X$ and $DX = X$. In [2], it is shown that the reducing subspaces are pertinent to the study of wavelets in subspace. It is shown there that such a subspace always has an *orthonormal wavelet*. It is natural to ask the question about the existence of NTF wavelets for an arbitrary subspace of $L^2(\mathbb{R})$. One may even try to characterize all the NTF wavelets for a given subspace of $L^2(\mathbb{R})$. As a first step, perhaps it will shed some light on the whole picture by looking at the NTF wavelets which are generated by some subsets of \mathbb{R} .

In this paper, we show that if X is a reducing subspace with an NTF wavelet ψ whose Fourier transform is $\frac{1}{\sqrt{2\pi}}\chi_E$, then E can be completely characterized (Theorem 1). On the other hand, we also show that if X admits an NTF wavelet ψ whose Fourier transform is $\frac{1}{\sqrt{2\pi}}\chi_E$ for a Bessel set E , then X must be a reducing subspace (Theorem 3). Furthermore, we prove that a maximal reducing subspace exists in the subspace X_E generated (see Definition 3) by a Bessel set E and this maximal reducing subspace is completely characterized (Theorem 2).

We will use μ to denote the Lebesgue measure. The inner product of two functions f and g in $L^2(\mathbb{R})$ is denoted by $\langle f, g \rangle$. The Fourier transform of a function f in $L^2(\mathbb{R})$ is denoted by $\mathcal{F}(f)$ or simply \hat{f} . Define $\hat{T} = \mathcal{F}T\mathcal{F}^{-1}$, $\hat{D} = \mathcal{F}D\mathcal{F}^{-1}$. We have $\hat{D} = D^{-1} = D^*$ and $\hat{T}(f) = e^{-is}f$ for $f \in L^2(\mathbb{R})$.

Let us first introduce some set theoretic notations.

Let E be a set. Define $\tau(E) = \bigcup_{n \in \mathbb{Z}} (E \cap [2n\pi, 2(n+1)\pi) - 2n\pi)$. If this is a disjoint union, we say that E is 2π -translation equivalent to $\tau(E)$, which is always a subset of $[0, 2\pi)$. If E and F are 2π -translation equivalent to the same subset in $[0, 2\pi)$, then we say E and F are 2π -translation equivalent. This defines an equivalence relation and is denoted by $\overset{\sim}{\sim}$. It is clear that $\mu(E) \geq \mu(\tau(E))$. The equality holds if and only if $E \overset{\sim}{\sim} \tau(E)$. If $E \overset{\sim}{\sim} F$, then $\mu(E) = \mu(F)$. Two points $x, y \in E$ are said to be 2π -translation equivalent in E if $x - y = 2m\pi$ for some integer m . The 2π -translation redundancy index of a point x in E is the number of elements in its equivalence class. We write $E(\tau, k)$ for the set of all points in E with 2π -translation redundancy index k . Of course, $E(\tau, k)$ could be an empty set, a proper subset of E , or the set E itself. For $k \neq m$, $E(\tau, k) \cap E(\tau, m) = \emptyset$, so $E = E(\tau, \infty) \cup (\bigcup_{n \in \mathbb{N}} E(\tau, n))$ is a disjoint union.

Similarly, let E be a set and define

$$\delta(E) = \bigcup_{n \in \mathbb{Z}} 2^{-n}(E \cap ([-2^{n+1}\pi, -2^n\pi) \cup [2^n\pi, 2^{n+1}\pi))).$$

If this union is disjoint, we say that E is 2 -dilation equivalent to $\delta(E)$, which is a subset of $[-2\pi, -\pi) \cup [\pi, 2\pi)$, the *Littlewood-Paley wavelet set*. If E and F are 2 -dilation equivalent to the same subset in $[-2\pi, -\pi) \cup [\pi, 2\pi)$, then we say E and F are 2 -dilation equivalent. This also defines an equivalence relation and is denoted by $\overset{\delta}{\sim}$. Two non-zero points $x, y \in E$ are said to be 2 -dilation equivalent if $\log_2 \frac{x}{y} \in \mathbb{Z}$. The 2 -dilation redundancy index of a point x in E is the number of elements in its equivalence class. The set of all points in E with 2 -dilation

redundancy index k is denoted by $E(\delta, k)$. For $k \neq m$, $E(\delta, k) \cap E(\delta, m) = \emptyset$ and $E = E(\delta, \infty) \cup (\bigcup_{n \in \mathbb{N}} E(\delta, n))$ is a disjoint union.

When E is a measurable set, one can prove that $E(\tau, m)$ and $E(\delta, m)$ are also measurable for any $m \geq 1$. Furthermore, each $E(\tau, m)$ can be decomposed into m disjoint measurable subsets $E^{(j)}(\tau, m)$ ($j = 1, 2, \dots, m$) such that each $E^{(j)}(\tau, m)$ is 2π -translation equivalent to a subset of $[0, 2\pi)$. Similarly, each $E(\delta, m)$ can be decomposed into m disjoint measurable subsets $E^{(j)}(\delta, m)$ ($j = 1, 2, \dots, m$) such that each $E^{(j)}(\delta, m)$ is 2-dilation equivalent to a subset of $[-2\pi, -\pi) \cup [\pi, 2\pi)$. The proof of these facts is elementary and is left to our reader. Notice that the decompositions of $E(\tau, m)$ and $E(\delta, m)$ here are not unique, but we will choose one and stay with it throughout the paper.

Definition 1. A measurable set E of finite measure is called a *basic set* if there exists $M > 0$ such that $\mu(E(\tau, m)) = \mu(E(\delta, m)) = 0$ for all $m > M$.

Let X be a subspace of $L^2(\mathbb{R})$. A function $\psi \in X$ is an *orthonormal wavelet* for X if the set $\{D^n T^\ell \psi : n, \ell \in \mathbb{Z}\}$ forms an orthonormal basis for X . A measurable set $E \subset \mathbb{R}$ is a *wavelet set* for X if the Fourier inverse transform of $\frac{1}{\sqrt{2\pi}}\chi_E$, namely $\mathcal{F}^{-1}(\frac{1}{\sqrt{2\pi}}\chi_E)$, is an orthonormal wavelet for the subspace X . It is shown in [2] that X is a reducing subspace if and only if there exists a subset Ω of \mathbb{R} with the property $\Omega = 2\Omega$ such that $\widehat{X} = L^2(\mathbb{R}) \cdot \chi_\Omega$, where \widehat{X} is the set of all Fourier transforms of elements in X (which is itself a subspace of $L^2(\mathbb{R})$).

Definition 2. Let X be a closed subspace of $L^2(\mathbb{R})$ and let ψ be a function in $L^2(\mathbb{R})$. Then ψ is called an NTF wavelet for X if $D^n T^\ell \psi \in X$ for each $n, \ell \in \mathbb{Z}$ and

$$(1) \quad f = \sum_{n, \ell \in \mathbb{Z}} \langle f, D^n T^\ell \psi \rangle D^n T^\ell \psi, \forall f \in X.$$

A measurable set E is called an NTF wavelet set for X if the function ψ_E defined by $\widehat{\psi}_E = \frac{1}{\sqrt{2\pi}}\chi_E$ is an NTF wavelet for X .

Remark. The convergence in (1) is unconditional in norm topology of $L^2(\mathbb{R})$. The same is true for all other similar occasions unless otherwise specified.

Let E be a Lebesgue measurable set with finite measure. Let ψ be the function defined by $\widehat{\psi}(s) = \frac{1}{\sqrt{2\pi}}\chi_E$. Consider the formal sum

$$(2) \quad G_E(f) = \sum_{n, \ell \in \mathbb{Z}} \langle f, D^n T^\ell \psi \rangle D^n T^\ell \psi,$$

where f is any function in $L^2(\mathbb{R})$. If $G_E(f)$ converges (in $L^2(\mathbb{R})$ norm) for all $f \in L^2(\mathbb{R})$, then it defines a bounded linear operator on $L^2(\mathbb{R})$ (by the Banach-Steinhaus Theorem). Apparently, when this is the case, E has to be a Bessel set. By definition, a frame wavelet set is always a Bessel set but not vice versa. The following proposition links the Bessel sets and basic sets.

Proposition 1. *Let E be a measurable set of finite measure. Then the following statements are equivalent:*

- (i) E is a Bessel set.
- (ii) E is a basic set.
- (iii) G_E is a bounded linear operator.

The proof is omitted since (iii) \implies (i) is trivial, (i) \implies (ii) and (ii) \implies (iii) can be found in Theorem 1 and its proof in [3].

Denote the formal sums

$$\sum_{\ell \in \mathbb{Z}} \langle \widehat{f}, \widehat{D}^k \widehat{T}^\ell \frac{1}{\sqrt{2\pi}} \chi_E \rangle \widehat{D}^k \widehat{T}^\ell \frac{1}{\sqrt{2\pi}} \chi_E$$

and

$$\sum_{k, \ell \in \mathbb{Z}} \langle \widehat{f}, \widehat{D}^k \widehat{T}^\ell \frac{1}{\sqrt{2\pi}} \chi_E \rangle \widehat{D}^k \widehat{T}^\ell \frac{1}{\sqrt{2\pi}} \chi_E$$

by $H_E^k(\widehat{f})$ and $H_E(\widehat{f})$ respectively. Notice that $A\|f\|^2 \leq \langle G_E(f), f \rangle \leq B\|f\|^2$ is equivalent to $A\|\widehat{f}\|^2 \leq \langle H_E(\widehat{f}), \widehat{f} \rangle \leq B\|\widehat{f}\|^2$.

Definition 3. Let E be a Bessel set. Then $X_E = \{f \in L^2(\mathbb{R}) : f = G_E(f)\}$ is called the subspace generated by E .

Note. In general, the function ψ defined by $\widehat{\psi} = \frac{1}{\sqrt{2\pi}} \chi_E$ may not belong to X_E so E may not be an NTF wavelet set for X_E . But if E is an NTF wavelet set for a closed subspace X , then $X = X_E$ by the definitions. What we are interested in is that under what conditions will E be an NTF wavelet set of X_E (or any other subspace X)? Furthermore, under what conditions will X_E be a reducing subspace? Some partial answers to these questions are obtained in this paper.

Let E be a Lebesgue measurable set in \mathbb{R} with finite positive measure. For any $f \in L^2(\mathbb{R})$, let $\widehat{f}_{m_j}^k$ be the $2^{k+1}\pi$ periodic extension of $\widehat{f} \cdot \chi_{2^k E(j)(\tau, m)}$ over \mathbb{R} . The following proposition is the main tool we need in proving our results. Its proof can be found in [3].

Proposition 2. Let E be a Bessel set (so it is also a basic set). Let M be as defined in Definition 1. Then $H_E^k(\widehat{f})$ converges to $\sum_{m=1}^M \sum_{j=1}^m \widehat{f}_{m_j}^k \cdot \chi_{2^k E}$ and $H_E(\widehat{f})$ converges to $\sum_{k \in \mathbb{Z}} \sum_{m=1}^M \sum_{j=1}^m \widehat{f}_{m_j}^k \cdot \chi_{2^k E}$ for all $f \in L^2(\mathbb{R})$. Here the convergence is unconditional under the norm of $L^2(\mathbb{R})$.

§2. THEOREMS AND PROOFS

We begin this section with the following theorem, which is a natural generalization of the characterization (Theorem 5.4, [7]) of normalized tight frame wavelet set for $L^2(\mathbb{R})$ to the reducing subspaces.

Theorem 1. Let X be a reducing subspace so that $\widehat{X} = L^2(\mathbb{R}) \cdot \chi_\Omega$, where $\Omega = 2\Omega$ is a measurable subset in \mathbb{R} . Then a measurable set E is an NTF wavelet set for X if and only if $E = E(\delta, 1) = E(\tau, 1)$ and $\bigcup_{k \in \mathbb{Z}} 2^k E = \Omega$.

Proof. Assume that E satisfies the conditions in Theorem 1. For any $f \in X$, we have $\widehat{f} = \sum_{n \in \mathbb{Z}} \widehat{f} \chi_{2^n E}$ since $\Omega = \bigcup_{n \in \mathbb{Z}} 2^n E$ and the union is a disjoint union. But by Proposition 2, $H_E(\widehat{f})$ also converges in $L^2(\mathbb{R})$ to $\sum_{n \in \mathbb{Z}} H_E^n(\widehat{f}) = \sum_{n \in \mathbb{Z}} \widehat{f} \chi_{2^n E}$ since $\mu(E(\tau, m)) = \mu(E(\delta, m)) = 0$ for all $m \geq 2$.

Now assume that E is an NTF wavelet set for X . Since X is a reducing subspace by the assumption, $\chi_E \in X = L^2(\mathbb{R}) \cdot \chi_\Omega$, where $\Omega = 2\Omega$. So $E \subseteq \Omega$ and hence $E(\tau, m) \subseteq \Omega$. Therefore, $\widehat{f} = \chi_{E(\tau, m)} \in X$ and $\widehat{f} = H_E(\widehat{f})$. By Proposition (2), $H_E^0(\widehat{f}) = m\widehat{f}$. So $\|\widehat{f}\|^2 = \langle H_E(\widehat{f}), \widehat{f} \rangle \geq \langle H_E^0(\widehat{f}), \widehat{f} \rangle = m\|\widehat{f}\|^2$. Therefore $\mu(E(\tau, m)) = 0$ for all $m > 1$ and $E = E(\tau, 1)$. If $\mu(E(\delta, k)) > 0$ for some $k > 1$,

we can find a subset F of E such that $\mu(F) > 0$ and $2^p F \subset E$ for some $p > 0$. A contradiction can be then derived in a similar way by choosing $\hat{f} = \chi_F$. The fact that $\Omega = \bigcup_{n \in \mathbb{Z}} 2^n E$ is rather obvious. \square

Theorem 2. *Let E be a Bessel set. Then:*

(i) *If $E = E(\delta, 1)$, then X_E is a reducing subspace with $\hat{X}_E = L^2(\mathbb{R})\chi_\Omega$, where $\Omega = \bigcup_{n \in \mathbb{Z}} 2^n E(\tau, 1)$.*

(ii) *If $E = E(\tau, 1)$, then X_E is a reducing subspace with $\hat{X}_E = L^2(\mathbb{R})\chi_\Omega$, where $\Omega = \bigcup_{n \in \mathbb{Z}} 2^n E(\delta, 1)$.*

(iii) *There exists a maximal reducing subspace $Y \subset X_E$. Furthermore, $\hat{Y} = L^2(\mathbb{R})\chi_\Omega$, where $\Omega = \bigcup_{n \in \mathbb{Z}} 2^n (E(\delta, 1) \cap E(\tau, 1))$.*

Proof. (i) Let $\hat{f} \in L^2(\mathbb{R}) \cdot \chi_\Omega$, and $F = E(\tau, 1)$. Since $\text{supp}(\hat{f}) \subset \Omega = \bigcup_{n \in \mathbb{Z}} 2^n F$, $\hat{f}_{m_j}^k = 0$ for all $m \geq 2$. Therefore, by Proposition 2, $H_E(\hat{f}) = \sum_{n \in \mathbb{Z}} \hat{f} \cdot \chi_{2^n E} = \hat{f}$. So $f \in X_E$ and it follows that $L^2(\mathbb{R}) \cdot \chi_\Omega \subseteq X_E$.

Now assume $f \in X_E$. By Proposition 2, we have

$$\hat{f} = H_E(\hat{f}) = \sum_{n \in \mathbb{Z}} \sum_{m=1}^M \sum_{j=1}^m \hat{f}_{m_j}^n \chi_{2^n E}$$

where the summation converges in $L^2(\mathbb{R})$ norm and hence converges almost everywhere. Let $s \in 2^n E(\tau, m)$ be any point such that the summation converges at all its $2^{n+1}\pi$ equivalent points (denoted by s_1, s_2, \dots, s_m) in $2^n E(\tau, m)$. Then we have $\hat{f}(s_i) = \sum_{1 \leq j \leq m} \hat{f}(s_j)$ for any $1 \leq i \leq m$ by Proposition 1. This leads to $\hat{f}(s_j) = 0$ if $m > 1$. Since s is arbitrary, $\hat{f} = 0$ on $2^n E(\tau, m)$ for any $n \in \mathbb{Z}$ and any $m > 1$. Therefore, $\text{supp}(\hat{f}) \subset \bigcup_{n \in \mathbb{Z}} 2^n E(\tau, 1)$.

(ii) It is obvious that $L^2(\mathbb{R})\chi_\Omega \subset \hat{X}_E$. Let $F = E \setminus E(\delta, 1)$, and let $\Omega_1 = \bigcup_{n \in \mathbb{Z}} 2^n E(\delta, 1)$, $\Omega_2 = \bigcup_{n \in \mathbb{Z}} 2^n F$. For any $f \in X_E$, $\|f\|^2 = \|\hat{f}\chi_{\Omega_1}\|^2 + \|\hat{f}\chi_{\Omega_2}\|^2$. On the other hand, since $E = E(\tau, 1)$, we have $\hat{f} = \sum_{n \in \mathbb{Z}} \hat{f}\chi_{2^n E}$ by Proposition 2. It follows that $\|\hat{f}\|^2 \geq \int_{\Omega_1} |\hat{f}|^2 ds + 2 \int_{\Omega_2} |\hat{f}|^2 ds$. Thus, $\int_{\Omega_2} |\hat{f}|^2 ds = 0$ and hence the support of \hat{f} is contained in Ω_1 .

(iii) It is obvious that the subspace defined by $\hat{Y} = L^2(\mathbb{R})\chi_\Omega$ is a reducing subspace. We need only to show that if Y_1 is also a reducing subspace contained in X_E , then $Y_1 \subset Y$. Let $f \in Y_1$; then $\hat{f} \in \hat{Y}_1$ and hence $|\hat{f}| \in \hat{Y}_1$. It follows that $|\hat{f}| \in \hat{X}_E$. By Proposition 1 and similar arguments in the above proofs, we get $\text{supp}(\hat{f}) \subset \Omega$. Hence $\hat{f} \in \hat{Y}$. Details are left to our reader. \square

Theorem 3. *Let X be a closed subspace of $L^2(\mathbb{R})$. The following statements are equivalent:*

- (i) *X is a reducing subspace.*
- (ii) *$\hat{X} = L^2(\mathbb{R}) \cdot \chi_\Omega$, where $\Omega = 2\Omega$ is a Lebesgue measurable set.*
- (iii) *There exists a Bessel set E which is a wavelet set for X .*
- (iv) *There exists a Bessel set E which is an NTF wavelet set for X .*
- (v) *$X = X_E$ for a Bessel set E such that $E = E(\delta, 1) = E(\tau, 1)$.*

Proof. (i) \Rightarrow (ii) is from Proposition 4.3 in [2]. (ii) \Rightarrow (iii) is Theorem 4.4 in [2]. (iii) \Rightarrow (iv) is trivial and (v) \Rightarrow (i) is implied by Theorem 2 above. So we only need to show that (iv) \Rightarrow (v). Notice that in general we have $\langle H_E(\hat{f}), \hat{f} \rangle \geq \langle H_E^0(\hat{f}), \hat{f} \rangle$

for any $f \in L^2(\mathbb{R})$. Let $\hat{f} = \chi_E \in \hat{X}$; we have $\mu(E) = \|\hat{f}\|^2 = \langle H_E(\hat{f}), \hat{f} \rangle$. On the other hand, $H_E^0(f) = \sum_{m=1}^M m\chi_{E(\tau, m)}$ by Proposition 2. It follows that $\mu(E) \geq \sum_{m=1}^M \int_{\mathbb{R}} m\chi_{E(\tau, m)} ds = \sum_{m=1}^M m\mu(E(\tau, m))$, hence $\mu(E(\tau, m)) = 0$ for all $m > 1$. Similarly, one can show that $\mu(E(\delta, m)) = 0$ for all $m > 1$. $X = X_E$ since E is an NTF set for X as pointed out in the note after Definition 3. \square

§3. EXAMPLES AND DISCUSSIONS

We now conclude this paper with some examples and discussions.

1. Let X be a closed reducing subspace of D, T . Then for any $0 < \alpha < 2\pi$, we can find an NTF wavelet set E for X such that $\mu(E) = \alpha$ as shown below.

Let $\hat{X} = L^2(\mathbb{R}) \cdot \chi_{\Omega}$ where $\Omega = 2\Omega$. It is shown in [2] that there exists a measurable set S such that $S = S(\delta, 1)$, $\bigcup_{k \in \mathbb{Z}} 2^k S = \Omega$ and $S \stackrel{\sim}{\sim} [0, 2\pi]$. Let $n \in \mathbb{Z}$ be such that $\frac{\pi}{2^{n-1}} < \min\{\alpha, 2\pi - \alpha\}$. For any $t \in [\frac{2\pi-\alpha}{2}, \pi]$, define $A = \bigcup_{0 \neq l \in \mathbb{Z}} (S \cap ((-t, t) + 2l\pi))$ and $B = S \cap ((-t, -\frac{\pi}{2^n}) \cup (\frac{\pi}{2^n}, t))$. Let $C = S \setminus (A \cup B)$ and D be the subset of $(-\frac{\pi}{2^n}, -\frac{\pi}{2^{n+1}}) \cup (\frac{\pi}{2^{n+1}}, \frac{\pi}{2^n})$ that is 2-dilation equivalent to $A \cup B$. It can be verified that $S_t = C \cup D$ is an NTF wavelet set. Note that when $t = \frac{2\pi-\alpha}{2}$, the measure of S_t is greater than α , whereas when $t = \pi$, the measure of S_t is less than $\frac{\pi}{2^{n-1}}$ which is less than α . Since the measure of S_t is a continuous function of t , the conclusion follows.

2. Let E be a wavelet set (for $L^2(\mathbb{R})$ or a reducing subspace). Let F be a measurable subset of E . Then by Theorem 1, F is a normalized tight frame wavelet set for the subspace X_F with $\hat{X}_F = L^2(\mathbb{R}) \cdot \chi_{\Omega}$, where $\Omega = \bigcup_{n \in \mathbb{Z}} 2^n F$. In particular, any measurable subset in $[-2\pi, -\pi) \cup [\pi, 2\pi)$ with positive measure is such a set.

3. Any basic set E such that $E \neq E(\delta, 1)$ or $E \neq E(\tau, 1)$ is not an NTF wavelet set for X_E . This can be seen from Proposition 2.

4. Although Theorem 2 guarantees the existence of a maximal reducing subspace in X_E , it does not say when or whether X_E is itself a reducing subspace. The following example shows that in general X_E is not necessarily a reducing subspace. Let $E_1 = [2\pi, 2.75\pi)$, $E_2 = [4\pi, 4.75\pi)$, $E_3 = [4.75\pi, 5\pi)$, $E_4 = [10\pi, 10.75\pi)$, $E_5 = [10.75\pi, 11\pi)$ and $E = \bigcup_j E_j$. We have $E = E(\delta, 2)$, $E(\tau, 1) = \emptyset$, $E(\tau, 2) = E_3 \cup E_5$ and $E(\tau, 3) = E_1 \cup E_2 \cup E_4$. Define $\hat{f}(s) = 2\chi_{E_1} - \chi_{E_2} - \chi_{E_4} + \chi_{E_3} - \chi_{E_5}$. $H_E(\hat{f})(s) = \hat{f}(s)$ by Proposition 1 so $\hat{f}(s) \in \hat{X}_E$, but $H_E(|\hat{f}|)(s) \neq |\hat{f}(s)|$. Thus, X_E cannot be a reducing subspace. Since if it were, then $|\hat{f}(s)|$ would be in it as well.

5. Although we proved in Theorem 3 that any subspace admitting an NTF wavelet defined by a Bessel set is necessarily a reducing subspace, this is not true for general NTF wavelets. In fact, one can construct non-reducing subspaces of $L^2(\mathbb{R})$ that admit wavelets (hence NTF wavelets); see [2].

ACKNOWLEDGEMENT

The authors thank Professor David Larson for some inspiring discussions.

REFERENCES

- [1] R. J. Duffin and A. C. Shaffer, A Class of Nonharmonic Fourier Series, *Trans. Amer. Math. Soc.*, 72 (1952), 341-366. MR **13**:839a
- [2] X. Dai and S. Lu, Wavelets in Subspaces, *The Michigan Math. J.*, Vol. 43, (1996), 81-98. MR **97m**:42021
- [3] X. Dai, Y. Diao and Q. Gu, Frame Wavelet Sets in \mathbb{R} , *Proc. of AMS*, to appear. CMP 2001:11

- [4] I. Daubechies, The Wavelet Transform, Time-frequency Localization and Signal Analysis, IEEE Trans. Inform. Theory, 36 (1990), 961-1005. MR **91e**:42038
- [5] I. Daubechies, Ten Lectures on Wavelets, CBS-NSF Regional Conferences in Applied Mathematics, 61, SIAM, (1992). MR **93e**:42045
- [6] I. Daubechies, A. Grossman and Y. Meyer, Painless Nonorthogonal Expansions, J. Math. Phys., 27(5) (1986), 1271-1283. MR **87e**:81089
- [7] D. Han and D. Larson, Frames, Bases and Group Representation, Memoirs of the AMS, to appear. MR **2001a**:47013
- [8] E. Hernandez and G. Weiss, A first Course on Wavelets, CRC Press, Boca Raton, 1996. MR **97i**:42015
- [9] Y. Meyer, *Ondelettes et operateurs I*, Hermann editeurs des sciences et des arts, 1990; Eng. transl, *Wavelets and Operators*, Camb. Studies in Adv. Math., 37, 1992. MR **93i**:42002
- [10] B. Sz-Nagy, Expansion Theorems of Paley-Wiener Type, Duke Math. J., 14 (1947), 975-978.
- [11] R. M. Young, An Introduction to Nonharmonic Fourier Series, Academic Press, New York, 1980. MR **81m**:42027

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NORTH CAROLINA AT CHARLOTTE, CHARLOTTE, NORTH CAROLINA 28223-9998

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NORTH CAROLINA AT CHARLOTTE, CHARLOTTE, NORTH CAROLINA 28223-9998

DEPARTMENT OF MATHEMATICS, EAST CHINA NORMAL UNIVERSITY, SHANGHAI, PEOPLE'S REPUBLIC OF CHINA