

## LOGARITHMIC CONVEXITY OF EXTENDED MEAN VALUES

FENG QI

(Communicated by Carmen C. Chicone)

ABSTRACT. In this article, the logarithmic convexity of the extended mean values are proved and an inequality of mean values is presented. As by-products, two analytic inequalities are derived. Two open problems are proposed.

### 1. INTRODUCTION

The so-called extended mean values  $E(r, s; x, y)$  were first defined by Professor K. B. Stolarsky in [21] as follows:

$$(1) \quad E(r, s; x, y) = \left[ \frac{r}{s} \cdot \frac{y^s - x^s}{y^r - x^r} \right]^{1/(s-r)}, \quad rs(r-s)(x-y) \neq 0,$$

$$(2) \quad E(r, 0; x, y) = \left[ \frac{1}{r} \cdot \frac{y^r - x^r}{\ln y - \ln x} \right]^{1/r}, \quad r(x-y) \neq 0,$$

$$(3) \quad E(r, r; x, y) = e^{-1/r} \left( \frac{x^{x^r}}{y^{y^r}} \right)^{1/(x^r - y^r)}, \quad r(x-y) \neq 0,$$

$$(4) \quad \begin{aligned} E(0, 0; x, y) &= \sqrt{xy}, & x &\neq y, \\ E(r, s; x, x) &= x, & x &= y. \end{aligned}$$

Define a function  $g$  by

$$(5) \quad g(t) = g(t; x, y) = \begin{cases} \frac{(y^t - x^t)}{t}, & t \neq 0, \\ \ln y - \ln x, & t = 0. \end{cases}$$

It is easy to see that  $g$  can be expressed in integral form as

$$(6) \quad g(t; x, y) = \int_x^y u^{t-1} du, \quad t \in \mathbb{R},$$

and

$$(7) \quad g^{(n)}(t) = \int_x^y (\ln u)^n u^{t-1} du, \quad t \in \mathbb{R}.$$

---

Received by the editors May 31, 2000 and, in revised form, December 26, 2000.

2000 *Mathematics Subject Classification*. Primary 26A51; Secondary 26B25, 26D07.

*Key words and phrases*. Logarithmic convexity, extended mean values, inequality, exponential function, absolutely monotonic function.

The author was supported in part by NSF of Henan Province (no. 004051800), SF for Pure Research of the Education Department of Henan Province (no. 1999110004), SF for the Prominent Youth of Henan Province, Doctor Fund of Jiaozuo Institute of Technology, SF of Henan Innovation Talents at Universities, and NNSF (no. 10001016) of China.

Therefore, the extended mean values can be represented [8, 19] in terms of  $g$  by

$$(8) \quad E(r, s; x, y) = \begin{cases} \left(\frac{g(s; x, y)}{g(r; x, y)}\right)^{1/(s-r)}, & (r-s)(x-y) \neq 0, \\ \exp\left(\frac{\partial g(r; x, y)/\partial r}{g(r; x, y)}\right), & r = s, x - y \neq 0, \end{cases}$$

and

$$(9) \quad \ln E(r, s; x, y) = \begin{cases} \frac{1}{s-r} \int_r^s \frac{\partial g(t; x, y)/\partial t}{g(t; x, y)} dt, & (r-s)(x-y) \neq 0, \\ \frac{\partial g(r; x, y)/\partial r}{g(r; x, y)}, & r = s, x - y \neq 0. \end{cases}$$

Leach and Sholander [3] showed that  $E(r, s; x, y)$  are increasing with both  $r$  and  $s$ , or with both  $x$  and  $y$ . The monotonicities of  $E$  have also been researched by the author and others in [1] and [13]–[19] using different ideas and simpler methods.

Leach and Sholander [4] and Páles [7] solved the problem of comparison of  $E$ ; that is, they found necessary and sufficient conditions for the parameters  $r, s, u, v$  in order that

$$(10) \quad E(r, s; x, y) \leq E(u, v; x, y)$$

be satisfied for all positive  $x$  and  $y$ .

Most of two variable means are special cases of  $E$ , for example [2],

$$(11) \quad E(1, 2; x, y) = A(x, y), \quad E(1, 1; x, y) = I(x, y), \quad E(0, 1; x, y) = L(x, y).$$

They are called the arithmetic mean, the identric mean, and the logarithmic mean, respectively.

Recently, the concepts of mean values have been generalized by the author in [9]–[12].

The main purpose of this paper is to verify the logarithmic convexity of the extended mean values  $E(r, s; x, y)$ . As applications, an inequality among the arithmetic mean, the identric mean and the logarithmic mean is established; two open problems are proposed. As by-products, an inequality for the exponential function is obtained.

## 2. LOGARITHMIC CONVEXITY OF $E(r, s; x, y)$

In order to prove our main result, the following lemma is necessary.

**Lemma 1** ([19]). *If  $f(t)$  is an increasing integrable function on  $I$ , then the arithmetic mean of function  $f(t)$ ,*

$$(12) \quad \phi(r, s) = \begin{cases} \frac{1}{s-r} \int_r^s f(t) dt, & r \neq s, \\ f(r), & r = s, \end{cases}$$

*is also increasing with both  $r$  and  $s$  on  $I$ .*

*If  $f$  is a twice-differentiable convex function, then the function  $\phi(r, s)$  is also convex with both  $r$  and  $s$  on  $I$ .*

*Proof.* Direct calculation yields

$$(13) \quad \frac{\partial\phi(r, s)}{\partial s} = \frac{1}{(s - r)^2} \left[ (s - r)f(s) - \int_r^s f(t) dt \right],$$

$$(14) \quad \frac{\partial^2\phi(r, s)}{\partial s^2} = \frac{(s - r)^2 f'(s) - 2(s - r)f(s) + 2 \int_r^s f(t) dt}{(s - r)^3} \equiv \frac{\varphi(r, s)}{(s - r)^3},$$

$$(15) \quad \frac{\partial\varphi(r, s)}{\partial s} = (s - r)^2 f''(s).$$

In the case of  $f(t)$  being increasing, we have  $\partial\phi(r, s)/\partial s \geq 0$ , thus  $\phi(r, s)$  increases in both  $r$  and  $s$ , since  $\phi(r, s) = \phi(s, r)$ .

In the case of  $f''(t) \geq 0$ ,  $\varphi(r, s)$  increases with  $s$ . Since  $\varphi(r, r) = 0$ , we have  $\partial^2\phi(r, s)/\partial s^2 \geq 0$ . Therefore  $\phi(r, s)$  is convex with respect to either  $r$  or  $s$ , since  $\phi(r, s) = \phi(s, r)$ . This completes the proof.  $\square$

By formula (9) and the above Lemma, we can see that, in order to prove the logarithmic convexity of the extended mean values  $E(r, s; x, y)$ , it suffices to verify the convexity of function

$$g'(t)/g(t) \triangleq g'_t(t; x, y)/g(t; x, y) \triangleq (\partial g(t; x, y)/\partial t)/g(t; x, y)$$

with respect to  $t$ , where  $g(t) = g(t; x, y)$  is defined by (5) or (6).

Straightforward computation results in

$$(16) \quad \left( \frac{g'(t)}{g(t)} \right)' = \frac{g''(t)g(t) - [g'(t)]^2}{g^2(t)},$$

$$(17) \quad \left( \frac{g'(t)}{g(t)} \right)'' = \frac{g^2(t)g'''(t) - 3g(t)g'(t)g''(t) + 2[g'(t)]^3}{g^3(t)}.$$

For  $y > x = 1$ , expanding  $g(t; 1, y)$  into series at  $t_0 = 0$  with respect to  $t$  directly gives us

$$(18) \quad \begin{aligned} g(t; 1, y) &= \sum_{i=0}^{\infty} \frac{(\ln y)^{i+1}}{(i + 1)!} t^i, \\ g'_t(t; 1, y) &= \sum_{i=0}^{\infty} \frac{(i + 1)(\ln y)^{i+2}}{(i + 2)!} t^i, \\ g''_t(t; 1, y) &= \sum_{i=0}^{\infty} \frac{(i + 1)(i + 2)(\ln y)^{i+3}}{(i + 3)!} t^i, \\ g'''_t(t; 1, y) &= \sum_{i=0}^{\infty} \frac{(i + 1)(i + 2)(i + 3)(\ln y)^{i+4}}{(i + 4)!} t^i. \end{aligned}$$

From the four fundamental operations of arithmetic and suitable properties of series, we have

$$(19) \quad g^2(t; 1, y) = 2 \sum_{k=0}^{\infty} \frac{(2^{k+1} - 1)(\ln y)^{k+2}}{(k + 2)!} t^k,$$

$$(20) \quad g(t; 1, y)g'_t(t; 1, y) = \sum_{k=0}^{\infty} \frac{(k + 1)(2^{k+2} - 1)(\ln y)^{k+3}}{(k + 3)!} t^k,$$

$$(21) \quad g(t; 1, y)g_t''(t; 1, y) + [g_t'(t; 1, y)]^2 = \sum_{k=0}^{\infty} \frac{(k+1)(k+2)(2^{k+3}-1)(\ln y)^{k+4}}{(k+4)!} t^k.$$

By standard arguments for series, we can get two combinatorial identities:

$$(22) \quad \sum_{i=0}^k (i+1) \binom{k+3}{i+2} = (k+1)(2^{k+2}-1),$$

$$(23) \quad \sum_{i=0}^k (i+1)(k-i+1) \binom{k+4}{i+2} = 2(k+3)(1+k \cdot 2^{k+1}).$$

By further computation, the following expansions are obtained:

$$(24) \quad \begin{aligned} g^2(t; 1, y)g_t'''(t; 1, y) &\equiv \sum_{k=0}^{\infty} \alpha_k (\ln y)^{k+6} t^k \\ &= \sum_{k=0}^{\infty} \frac{2}{(k+6)!} \sum_{i=0}^k (i+1)(i+2)(i+3)(2^{k-i+1}-1) \binom{k+6}{i+4} (\ln y)^{k+6} t^k, \end{aligned}$$

$$(25) \quad \begin{aligned} g(t; 1, y)g_t'(t; 1, y)g_t''(t; 1, y) &\equiv \sum_{k=0}^{\infty} \beta_k (\ln y)^{k+6} t^k \\ &= \sum_{k=0}^{\infty} \frac{1}{(k+6)!} \sum_{i=0}^k (i+1)(i+2)(k-i+1)(2^{k-i+2}-1) \binom{k+6}{i+3} (\ln y)^{k+6} t^k, \end{aligned}$$

$$(26) \quad \begin{aligned} g_t'(t; 1, y)[g_t'(t; 1, y)]^2 + g(t; 1, y)g_t''(t; 1, y) &\equiv \sum_{k=0}^{\infty} \gamma_k (\ln y)^{k+6} t^k \\ &= \sum_{k=0}^{\infty} \frac{1}{(k+6)!} \sum_{i=0}^k (i+1)(i+2)(k-i+1)(2^{i+3}-1) \binom{k+6}{i+4} (\ln y)^{k+6} t^k. \end{aligned}$$

**Proposition 1.** For an arbitrary nonnegative integer  $k$ , we have

$$(27) \quad \begin{aligned} 2 \sum_{i=0}^k (i+1)(i+2) \{ (i+3)2^{k-i+1} + (k-i+1)2^{i+3} - (k+4) \} \binom{k+6}{i+4} \\ \leq 5 \sum_{i=0}^k (i+1)(i+2)(k-i+1)(2^{k-i+2}-1) \binom{k+6}{i+3}. \end{aligned}$$

*Proof.* The inequality (27) is equivalent to

$$(28) \quad \begin{aligned} \sum_{j=0}^{k+6} (j-3)(j-2) [(k-j+5)2^j + (j-1)2^{k-j+6} - 2(k+4)] \binom{k+6}{j} \\ + (k^2 + 12k + 38)(2^{k+6} + k + 3) \\ \leq 5 \sum_{j=0}^{k+6} (j-2)(j-1)(k-j+4)(2^{k-j+5}-1) \binom{k+6}{j}. \end{aligned}$$

Using expansions-into-series of  $(1 + 2x)^n$  and its derivatives, we get

$$\begin{aligned}
 \sum_{i=0}^n 2^i \binom{n}{i} &= 3^n, \\
 \sum_{i=0}^n i 2^i \binom{n}{i} &= 2n3^{n-1}, \\
 \sum_{i=0}^n i(i-1) 2^i \binom{n}{i} &= 4n(n-1)3^{n-2}, \\
 \sum_{i=0}^n i(i-1)(i-2) 2^i \binom{n}{i} &= 8n(n-1)(n-2)3^{n-3}.
 \end{aligned}
 \tag{29}$$

Similarly, using expansions-into-series of  $(1 + x/2)^n$  and its derivatives, we have

$$\begin{aligned}
 \sum_{j=0}^n \frac{1}{2^j} \binom{n}{j} &= \left(\frac{3}{2}\right)^n, \\
 \sum_{j=0}^n \frac{j}{2^j} \binom{n}{j} &= \frac{n}{2} \left(\frac{3}{2}\right)^{n-1}, \\
 \sum_{j=0}^n \frac{j(j-1)}{2^j} \binom{n}{j} &= \frac{n(n-1)}{4} \left(\frac{3}{2}\right)^{n-2}, \\
 \sum_{j=0}^n \frac{j(j-1)(j-2)}{2^j} \binom{n}{j} &= \frac{n(n-1)(n-2)}{8} \left(\frac{3}{2}\right)^{n-3}.
 \end{aligned}
 \tag{30}$$

The following formulae are also well-known:

$$\begin{aligned}
 \sum_{j=0}^n \binom{n}{j} &= 2^n, \\
 \sum_{j=0}^n j \binom{n}{j} &= n2^{n-1}, \\
 \sum_{j=0}^n j(j-1) \binom{n}{j} &= n(n-1)2^{n-2}, \\
 \sum_{j=0}^n j(j-1)(j-2) \binom{n}{j} &= n(n-1)(n-2)2^{n-3}.
 \end{aligned}
 \tag{31}$$

Substitution of (29), (30) and (31) into (28) and simplification give us

$$(k^3 + 15k^2 + 74k + 114) + (k^3 + 15k^2 + 74k + 168)2^{k+3} - 2 \times 3^{k+6} \leq 0.
 \tag{32}$$

From Taylor’s expansion

$$a^x = \sum_{i=0}^{\infty} \frac{(\ln a)^i}{i!} x^i,
 \tag{33}$$

the inequality (32) reduces to

$$\begin{aligned}
 & 1458 + (666 + 1344 \ln 2)k + 8[84(\ln 2)^2 + 74 \ln 2 + 15]k^2 + k^3 \\
 (34) \quad & + 8 \sum_{i=3}^{\infty} \left[ \frac{168(\ln 2)^3}{i(i-1)(i-2)} + \frac{74(\ln 2)^2}{(i-1)(i-2)} + \frac{15 \ln 2}{i-2} + 1 \right] \frac{(\ln 2)^{i-3}}{(i-3)!} \cdot k^i \\
 & \leq 1458 \sum_{i=0}^{\infty} \frac{(\ln 3)^i}{i!} \cdot k^i.
 \end{aligned}$$

To prove inequality (34), by equating the coefficients of  $k^i$  in (34), it is sufficient to verify that

$$(35) \quad 4[168(\ln 2)^3 + 74(\ln 2)^2 i + 15i(i-1) \ln 2 + i(i-1)(i-2)] \leq 729(\ln 2)^3 (\log_2 3)^i$$

holds for  $i \geq 4$ .

Let

$$(36) \quad \begin{aligned}
 \psi(x) = & 4\{x^3 + (15 \ln 2 - 3)x^2 + [2 - 15 \ln 2 + 74(\ln 2)^2]x + 168(\ln 2)^3\} \\
 & - 729(\ln 2)^3 (\log_2 3)^x
 \end{aligned}$$

for  $x \geq 4$ . By standard argument, we have

$$\begin{aligned}
 \psi'(x) &= 4\{3x^2 + 2(15 \ln 2 - 3)x + 2 - 15 \ln 2 + 74(\ln 2)^2\} \\
 &\quad - 729(\ln 2)^3 \ln(\log_2 3) (\log_2 3)^x, \\
 \psi''(x) &= 4\{6x + 2(15 \ln 2 - 3)\} - 729(\ln 2)^3 [\ln(\log_2 3)]^2 (\log_2 3)^x, \\
 \psi'''(x) &= 24 - 729(\ln 2)^3 [\ln(\log_2 3)]^3 (\log_2 3)^x, \\
 \psi^{(4)}(x) &= -729(\ln 2)^3 (\log_2 3)^x (\ln(\log_2 3))^4 < 0.
 \end{aligned}$$

Direct computation yields

$$\begin{aligned}
 \psi'''(4) &= 24 - 729(\ln 2)^3 [\ln(\log_2 3)]^3 (\log_2 3)^4 < -125, \\
 \psi''(4) &= 4(18 + 30 \ln 2) - 729(\ln 2)^3 [\ln(\log_2 3)]^2 (\log_2 3)^4 < -169, \\
 \psi'(4) &= 4\{26 + 105 \ln 2 + 74(\ln 2)^2\} - 729(\ln 2)^3 \ln(\log_2 3) (\log_2 3)^4 < -168, \\
 \psi(4) &= 4\{24 + 180 \ln 2 + 296(\ln 2)^2 + 168(\ln 2)^3\} - 729(\ln 2)^3 (\log_2 3)^4 < -144.
 \end{aligned}$$

Since the function  $\psi'''(x)$  is decreasing, then  $\psi'''(x) < 0$  for  $x \geq 4$ , and  $\psi''(x) < 0$ ,  $\psi'(x) < 0$ , and then  $\psi(x) < 0$  for  $x \geq 4$ . Inequality (35) follows. This completes the proof. □

*Remark 1.* In fact, inequality  $\psi(x) < 0$  holds for  $x \geq 0$ .

**Corollary 1.** For any nonnegative number  $x \geq 0$ , we have

$$(37) \quad \frac{1 + 3^{x+3}}{1 + 2^{x+3}} \geq \frac{x^3 + 15x^2 + 74x + 168}{54}.$$

**Proposition 2.** If  $y > x = 1$ , then, for  $t \geq 0$ ,

$$(38) \quad g^2(t; 1, y)g_t'''(t; 1, y) - 3g(t; 1, y)g_t'(t; 1, y)g_t''(t; 1, y) + 2[g_t'(t; 1, y)]^3 \leq 0.$$

*Proof.* It is clear that

$$\begin{aligned}
 & g^2(t)g'''(t) - 3g(t)g'(t)g''(t) + 2[g'(t)]^3 \\
 (39) \quad & = g^2(t)g'''(t) - 5(g(t)g'(t))g''(t) + 2g'(t)[g(t)g''(t) + (g'(t))^2] \\
 & = \sum_{k=0}^{\infty} (\alpha_k - 5\beta_k + 2\gamma_k)(\ln y)^{k+6}t^k.
 \end{aligned}$$

Furthermore, Proposition 1 implies

$$(40) \quad \alpha_k - 5\beta_k + 2\gamma_k \leq 0, \quad k \geq 0.$$

The proof of Proposition 2 is completed. □

**Theorem 1.** *For all fixed  $x, y > 0$  and  $s \in [0, +\infty)$  (or  $r \in [0, +\infty)$ , respectively), the extended mean values  $E(r, s; x, y)$  are logarithmically concave in  $r$  (or in  $s$ , respectively) on  $[0, +\infty)$ . For all fixed  $x, y > 0$  and  $s \in (-\infty, 0]$  (or  $r \in (-\infty, 0]$ , respectively), the extended mean values  $E(r, s; x, y)$  are logarithmically convex in  $r$  (or in  $s$ , respectively) on  $(-\infty, 0]$ .*

*Proof.* The combination of Proposition 2 with equality (17) proves that  $\frac{g'_t(t;1,y)}{g(t;1,y)}$  is concave on  $[0, +\infty)$  with  $t$  for  $y > x = 1$ . Therefore, from Lemma 1, it follows that the extended mean values  $E(r, s; 1, y)$  are logarithmically concave on  $[0, +\infty)$  with respect to either  $r$  or  $s$  for  $y > x = 1$ .

By standard arguments, we obtain

$$(41) \quad E(r, s; x, y) = xE(r, s; 1, \frac{y}{x}),$$

$$(42) \quad E(-r, -s; x, y) = \frac{xy}{E(r, s; x, y)}.$$

Hence,  $E(r, s; x, y)$  are logarithmically concave on  $[0, +\infty)$  with either  $r$  or  $s$ , respectively, and logarithmically convex on  $(-\infty, 0]$  in either  $r$  or  $s$ , respectively. The proof of Theorem 1 is completed. □

**Corollary 2.** *The logarithmic means  $L(r; x, y) = (L(x^r, y^r))^{1/r}$  and the extended logarithmic means  $S_r(x, y) = E(r, 1; x, y)$  are logarithmically concave on  $[0, +\infty)$  with respect to  $r$ .*

**Corollary 3.** *For any  $y > x > 0$ , if  $t \geq 0$ , then*

$$(43) \quad g^2(t; x, y)g_t'''(t; x, y) - 3g(t; x, y)g_t'(t; x, y)g_t''(t; x, y) + 2[g_t'(t; x, y)]^3 \leq 0.$$

*If  $t \leq 0$ , the inequality (43) reverses.*

As concrete applications of the logarithmic convexity of  $E(r, s; x, y)$ , an inequality of mean values is given as follows.

**Theorem 2.** *Let  $A(x, y)$ ,  $L(x, y)$  and  $I(x, y)$  denote the arithmetic mean, the logarithmic mean, and the identric mean of two variables  $x$  and  $y$ . Then, for  $x \neq y$ , we have*

$$(44) \quad I(x, y) > \frac{L(x, y) + A(x, y)}{2}.$$

*Proof.* P. Montel [5, p. 19] verified that a positive function  $f(x)$  is logarithmically convex if and only if  $x \mapsto e^{ax}f(x)$  is a convex function for all real values of  $a$ . Thus, applied this conclusion to the function  $e^{ar} E(r, s; x, y)$ , for any given  $a \in \mathbb{R}$  and  $x, y > 0, x \neq y$ , we get

$$(45) \quad I(x, y) \geq \frac{e^{-a}L(x, y) + e^aA(x, y)}{2} \geq \sqrt{A(x, y)L(x, y)}.$$

The proof is complete. □

### 3. MISCELLANEA

A function  $f(t)$  is said to be absolutely monotonic on  $(a, b)$  if it has derivatives of all orders and  $f^{(k)}(t) \geq 0, t \in (a, b), k \in \mathbb{N}$ .

A function  $f(t)$  is said to be completely monotonic on  $(a, b)$  if it has derivatives of all orders and  $(-1)^k f^{(k)}(t) \geq 0, t \in (a, b), k \in \mathbb{N}$ .

The famous Bernstein-Widder theorem [22] states that a function  $f(x), x \in (0, +\infty)$ , is absolutely monotone if and only if there exists a bounded and nondecreasing function  $\sigma(t)$  such that the integral

$$(46) \quad f(x) = \int_0^{+\infty} e^{xt} d\sigma(t)$$

converges for  $0 \leq x < +\infty$ , and a function  $f(x), x \in (0, +\infty)$ , is completely monotone if and only if there exists a bounded and nondecreasing function  $\eta(t)$  such that the integral

$$(47) \quad f(x) = \int_0^{+\infty} e^{-xt} d\eta(t)$$

converges for  $0 \leq x < +\infty$ .

**Proposition 3.** *Suppose  $F(t) = \int_a^b p(u)f^t(u) du, t \in \mathbb{R}, p(u) \not\equiv 0$ , is a nonnegative and continuous function, and  $f(u)$  a positive and continuous function on a given interval  $[a, b]$ . Then*

$$(48) \quad F^{(n)}(t) = \int_a^b p(u)f^t(u) [\ln f(u)]^n du.$$

*If  $f(u) \geq 1, F(t)$  is absolutely monotone on  $(-\infty, +\infty)$ ; if  $0 < f(u) < 1$ , then  $F(t)$  is completely monotone on  $(-\infty, +\infty)$ . Moreover,  $F(t)$  is absolutely convex on  $(-\infty, +\infty)$ .*

*Proof.* This is obvious. □

**Corollary 4** ([17, 18]). *The function  $g(t; x, y)$  is absolutely and regularly monotonic on  $(-\infty, +\infty)$  for  $y > x > 1$ , or on  $(0, +\infty)$  for  $y > x^{-1} > 1$ , completely and regularly monotonic on  $(-\infty, +\infty)$  for  $0 < x < y < 1$ , or on  $(-\infty, 0)$  for  $1 < y < x^{-1}$ . Furthermore,  $g(x)$  is absolutely convex on  $(-\infty, +\infty)$ .*

The generalized weighted mean values  $M_{p,f}(r, s; x, y)$ , with two parameters  $r$  and  $s$ , are defined in [9, 10] by

$$(49) \quad M_{p,f}(r, s; x, y) = \left( \frac{\int_x^y p(u) f^s(u) \, du}{\int_x^y p(u) f^r(u) \, du} \right)^{1/(s-r)}, \quad (r-s)(x-y) \neq 0,$$

$$(50) \quad M_{p,f}(r, r; x, y) = \exp \left( \frac{\int_x^y p(u) f^r(u) \ln f(u) \, du}{\int_x^y p(u) f^r(u) \, du} \right), \quad x-y \neq 0,$$

$$M_{p,f}(r, s; x, x) = f(x),$$

where  $x, y, r, s \in \mathbb{R}$ ,  $p(u) \not\equiv 0$  is a nonnegative and integrable function and  $f(u)$  a positive and integrable function on the interval between  $x$  and  $y$ .

It is clear that  $E(r, s; x, y)$  is a special case of  $M_{p,f}(r-1, s-1; x, y)$  applied to  $p(u) \equiv 1$ ,  $f(u) = u$ , and  $M^{[r]}(f; p; x, y) = M_{p,f}(r, 0; x, y)$ .

The basic properties and monotonicities of  $M_{p,f}(r, s; x, y)$  were studied in [9, 15, 16, 20].

At last, we propose the following two open problems.

**Open Problem 1.** If  $f(x)$  is an absolutely or completely monotonic function on the interval  $(-\infty, +\infty)$ , then the following inequality holds for  $0 \leq x < +\infty$  or reverses for  $-\infty < x \leq 0$ :

$$(51) \quad f^2(x)f'''(x) - 3f(x)f'(x)f''(x) + 2[f'(x)]^3 \leq 0.$$

**Open Problem 2.** Suppose  $p(u)$  is nonnegative and continuous, and  $f(u)$  positive and continuous on a given interval  $[a, b]$ . If  $f(u) \geq 1$  or  $0 < f(u) < 1$ , then the generalized weighted mean values  $M_{p,f}(r, s; x, y)$  are logarithmically concave on  $(0, +\infty)$  with respect to either  $r$  or  $s$ , respectively, or logarithmically convex on  $(-\infty, 0)$  with respect to either  $r$  or  $s$ , respectively.

#### ACKNOWLEDGEMENT

The author is thankful to the referee for his thoughtful suggestions, and to the Editor, Professor Carmen Chicone, for his kind help.

#### REFERENCES

- [1] Bai-Ni Guo, Shi-Qin Zhang, and Feng Qi, *Elementary proofs of monotonicity for extended mean values of some functions with two parameters*, Mathematics in Practice and Theory **29** (1999), no.2, 169–174. (Chinese) MR **2000g**:26007
- [2] Ji-Chang Kuang, *Applied Inequalities*, 2nd edition, Hunan Education Press, Changsha, China, 1993. (Chinese) MR **95j**:26001
- [3] E. Leach and M. Sholander, *Extended mean values*, Amer. Math. Monthly **85** (1978), 84–90. MR **58**:22428
- [4] E. Leach and M. Sholander, *Extended mean values II*, J. Math. Anal. Appl. **92** (1983), 207–223. MR **85b**:26007
- [5] D. S. Mitrinović, *Analytic Inequalities*, Springer-Verlag, Berlin, 1970. MR **43**:448
- [6] D. S. Mitrinović, J. E. Pečarić and A. M. Fink, *Classical and New Inequalities in Analysis*, Kluwer Academic Publishers, Dordrecht/Boston/London, 1993. MR **94c**:00004
- [7] Z. Páles, *Inequalities for differences of powers*, J. Math. Anal. Appl. **131** (1988), 271–281. MR **89f**:26023
- [8] J. E. Pečarić, Feng Qi, V. Šimić and Sen-Lin Xu, *Refinements and extensions of an inequality*, III, J. Math. Anal. Appl. **227** (1998), no. 2, 439–448. MR **99i**:26029
- [9] Feng Qi, *Generalized weighted mean values with two parameters*, Proceedings of the Royal Society of London Series A—Mathematical, Physical and Engineering Sciences **454** (1998), no. 1978, 2723–2732. MR **99k**:26027

- [10] Feng Qi, *On a two-parameter family of nonhomogeneous mean values*, Tamkang Journal of Mathematics **29** (1998), no. 2, 155–163. MR **99g**:26026
- [11] Feng Qi, *Studies on Problems in Topology and Geometry and on Generalized Weighted Abstracted Mean Values*, Thesis submitted for the degree of Doctor of Philosophy at University of Science and Technology of China, Hefei City, Anhui Province, China, Winter 1998. (Chinese)
- [12] Feng Qi, *Generalized abstracted mean values*, Journal of Inequalities in Pure and Applied Mathematics **1** (2000), no. 1, Article 4. [http://jipam.vu.edu.au/v1n1/013\\_99.html](http://jipam.vu.edu.au/v1n1/013_99.html). RGMIA Research Report Collection **2** (1999), no. 5, Article 4, 633–642. <http://rgmia.vu.edu.au/v2n5.html>. MR **2001d**:26048
- [13] Feng Qi and Qiu-Ming Luo, *Refinements and extensions of an inequality*, Mathematics and Informatics Quarterly **9** (1999), no. 1, 23–25.
- [14] Feng Qi and Qiu-Ming Luo, *A simple proof of monotonicity for extended mean values*, J. Math. Anal. Appl. **224** (1998), no. 2, 356–359. CMP 98:16
- [15] Feng Qi, Jia-Qiang Mei, Da-Feng Xia, and Sen-Lin Xu, *New proofs of weighted power mean inequalities and monotonicity for generalized weighted mean values*, Mathematical Inequalities and Applications **3** (2000), no. 3, 377–383. MR **2001d**:26040
- [16] Feng Qi, Jia-Qiang Mei, and Sen-Lin Xu, *Other proofs of monotonicity for generalized weighted mean values*, RGMIA Research Report Collection **2** (1999), no. 4, Article 6, 469–472. <http://rgmia.vu.edu.au/v2n4.html>.
- [17] Feng Qi and Sen-Lin Xu, *Refinements and extensions of an inequality*, II, J. Math. Anal. Appl. **211** (1997), 616–620. MR **99i**:26028
- [18] Feng Qi and Sen-Lin Xu, *The function  $(b^x - a^x)/x$ : Inequalities and properties*, Proc. Amer. Math. Soc. **126** (1998), no. 11, 3355–3359. MR **99a**:26001
- [19] Feng Qi, Sen-Lin Xu, and Lokenath Debnath, *A new proof of monotonicity for extended mean values*, Intern. J. Math. Math. Sci. **22** (1999), no. 2, 415–420. MR **2000c**:26019
- [20] Feng Qi and Shi-Qin Zhang, *Note on monotonicity of generalized weighted mean values*, Proceedings of the Royal Society of London Series A—Mathematical, Physical and Engineering Sciences **455** (1999), no. 1989, 3259–3260. CMP 2001:07
- [21] K. B. Stolarsky, *Generalizations of the logarithmic mean*, Math. Mag. **48** (1975), 87–92. MR **50**:10186
- [22] D. V. Widder, *The Laplace Transform*, Princeton University Press, Princeton, 1941. MR **3**:232d

DEPARTMENT OF MATHEMATICS, JIAOZUO INSTITUTE OF TECHNOLOGY, JIAOZUO CITY, HENAN 454000, PEOPLE'S REPUBLIC OF CHINA

*E-mail address*: [qifeng@jz.it.edu.cn](mailto:qifeng@jz.it.edu.cn)

*URL*: <http://rgmia.vu.edu.au/qi.html>