

## CONTINUOUS TRANSFORMATION OF BAIRE MEASURES INTO LEBESGUE MEASURE

HANS G. KELLERER

(Communicated by David Preiss)

ABSTRACT. A recent result by Wulbert on the existence of continuous functions with measure zero level sets is slightly extended and its proof is considerably simplified. As a by-product, a criterion is established for a Baire measure to allow a continuous transformation into Lebesgue measure.

### INTRODUCTION

As outlined in the survey [5] by Wulbert, the following result is of interest in  $L_1$ -approximation theory:

(\*) *Given a nonatomic  $\sigma$ -finite measure space  $(X, \mathfrak{A}, \mu)$ , for any finite dimensional subspace  $\mathcal{K}$  of  $\mathcal{L}_1(X, \mathfrak{A}, \mu)$  there is an extreme element  $g$  in the unit ball of the dual space, i.e. satisfying*

$$(1) \quad g(x) = \pm 1 \quad \text{for } \mu\text{-almost all } x \in X,$$

which annihilates  $\mathcal{K}$ , i.e. satisfies

$$(2) \quad \int_X fg d\mu = 0 \quad \text{for all } f \in \mathcal{K}.$$

One approach to this result uses Liapunov's convexity theorem, as has been carried out by Phelps and Dye [4, Theorem 2.5]. An alternative, yielding some additional insight, consists in applying the Borsuk–Ulam theorem on antipodal mappings, as has been done by Hobby and Rice [1] in the classical case  $X = [0, 1]$  and  $\mu = \lambda$  ( $:=$  Lebesgue measure). In fact, this approach has been anticipated in a stochastic context by the present author [2, Satz 3] for arbitrary measure spaces. The additional tool there is a transformation of the underlying measure  $\mu$  into Lebesgue measure, as is always available under the – obviously necessary – condition that  $\mu$  is nonatomic and  $\sigma$ -finite. Actually, all that is needed is an  $\mathfrak{A}$ -measurable function  $h$  with measure zero level sets, i.e.

$$(3) \quad \mu(\{x : h(x) = y\}) = 0 \quad \text{for all } y \in \mathbf{R}.$$

In a recent paper [6] Wulbert proves that in the topological situation the function  $h$  may even be chosen to be continuous, which yields for (\*) above a solution  $g$  being the sign of a continuous function. Since his construction is highly involved, a simple proof seemed to be desirable. On the way it turned out that by a slight generalization the question raised by the title of this paper finds a complete answer.

---

Received by the editors January 5, 2001.

2000 *Mathematics Subject Classification.* Primary 28C15; Secondary 46G10.

*Key words and phrases.* Baire measures, measure transformation, Lebesgue measure.

## PRELIMINARIES

In proving the main results, three simple lemmata will be used. First, the notion of moderatedness has to be extended from Borel to Baire measures:

**Definition 1.** For an arbitrary topological space  $X$  denote by  $\mathfrak{C}(X)$  the Baire  $\sigma$ -algebra, generated by the space  $\mathcal{C}(X)$  of continuous functions. Then a measure  $\mu$  on  $\mathfrak{C}(X)$  is called “moderated”, if  $\mathcal{C}(X)$  contains functions  $f_n, n \in \mathbf{N}$ , satisfying

$$X = \bigcup_{n \in \mathbf{N}} \{x : f_n(x) \neq 0\} \quad \text{and} \quad \mu(\{x : f_n(x) \neq 0\}) < \infty \quad \text{for all } n \in \mathbf{N}.$$

While any moderated Baire measure is  $\sigma$ -finite, the converse fails even for nonatomic measures. Indeed, choose an open base  $G_n, n \in \mathbf{N}$ , for  $X = [0, 1]$  and measurable functions  $f_n \geq 0, n \in \mathbf{N}$ , such that

$$\int_{G_n} f_n d\lambda = \infty \quad \text{and} \quad \sum_{n \in \mathbf{N}} \lambda(\{x : f_n(x) \neq 0\}) < \infty.$$

The last condition implies

$$\lambda(\{x : f_n(x) = 0 \text{ for almost all } n \in \mathbf{N}\}) = 1,$$

hence  $f := \sum_{n \in \mathbf{N}} f_n$  is  $\lambda$ -almost finite and thus  $d\mu = f d\lambda$  defines a  $\sigma$ -finite measure. Due to  $\mu(G) = \infty$  for all open sets  $G \neq \emptyset$ , however, it fails to be moderated (this example also solves the final problem in [6]).

The following criterion will be required in the sequel:

**Lemma 1.** *A Baire measure  $\mu|_{\mathfrak{C}(X)}$  is moderated if and only if  $\mathcal{C}(X)$  contains a  $\mu$ -integrable strictly positive function  $f$ .*

*Proof.* Given functions  $f_n, n \in \mathbf{N}$ , according to Definition 1, choose constants  $\gamma_n > 0$  such that

$$\sum_{n \in \mathbf{N}} \gamma_n < \infty \quad \text{and} \quad \sum_{n \in \mathbf{N}} \gamma_n \mu(\{x : f_n(x) \neq 0\}) < \infty.$$

Then the function

$$f := \sum_{n \in \mathbf{N}} (|f_n| \wedge \gamma_n)$$

meets all requirements. Conversely, if  $f$  is strictly positive and  $\mu$ -integrable, then the functions

$$f_n := \left(f - \frac{1}{n}\right)^+, n \in \mathbf{N},$$

satisfy the conditions of Definition 1. □

Next, some notation has to be introduced:

**Definition 2.** For a Hausdorff topological space  $Y$  denote by  $\mathfrak{B}(Y)$  the Borel  $\sigma$ -algebra and by  $M(Y)$  the space of all finite measures  $\nu$  on  $\mathfrak{B}(Y)$  endowed with the weak (narrow) topology. Then

$$\delta(\nu) := \max_{y \in Y} \nu(\{y\}) \quad \text{for } \nu \in M(Y).$$

The following fact will be used only for  $Y = \mathbf{R}$ , but is of independent interest:

**Lemma 2.** *If  $Y$  is a second countable metrizable space, then the functional  $\delta$  is upper semicontinuous.*

*Proof.* Fix  $\nu_0 \in M(Y)$  and  $\delta_0 > 0$  with  $\delta(\nu_0) < \delta_0$ . Since  $Y$  is second countable,  $\nu_0$ -atoms are singletons, which is easily seen to imply that  $Y$  may be partitioned into a finite number of sets  $B_m \in \mathfrak{B}(Y)$ ,  $1 \leq m \leq n$ , with  $\nu_0(B_m) < \delta_0$ . Since  $\nu_0$  is outer regular, this yields a cover of  $Y$  by open sets  $G_m$ ,  $1 \leq m \leq n$ , with  $\nu_0(G_m) < \delta_0$ . Now in a metrizable space each open set  $G$  can be approximated from inside by a closed set  $F$  with measure zero boundary  $\partial F$  (choose a metric  $d$  and consider the closed sets  $F_t := \{y : d(y, z) \geq t \text{ for } z \notin G\}$ , having disjoint boundaries for different values  $t > 0$ ). Thus there are closed sets  $F_m \subset G_m$  with

$$\nu_0(F_m) > \nu_0(G_m) - \frac{1}{n} \delta_0 \quad \text{and} \quad \nu_0(\partial F_m) = 0.$$

Adjoining to  $F_1, \dots, F_n$  the set

$$F_0 := Y \setminus \bigcup_{1 \leq m \leq n} (F_m \setminus \partial F_m)$$

yields a cover of  $Y$  by closed sets  $F_m$ ,  $0 \leq m \leq n$ , with  $\nu_0(F_m) < \delta_0$  for all  $m$ , because also

$$\begin{aligned} \nu_0(F_0) &= \nu_0\left(\bigcup_{1 \leq m \leq n} G_m\right) - \nu_0\left(\bigcup_{1 \leq m \leq n} F_m\right) \\ &\leq \sum_{1 \leq m \leq n} \nu_0(G_m \setminus F_m) < \delta_0. \end{aligned}$$

Thus the weakly open set

$$M_0 := \bigcap_{0 \leq m \leq n} \{\nu \in M(Y) : \nu(F_m) < \delta_0\}$$

contains  $\nu_0$  and satisfies  $\delta(\nu) < \delta_0$  for all  $\nu \in M_0$ , as had to be shown.  $\square$

The final auxiliary result is the essential content of [6, Lemma 4.1] (for a similar argument see the proof of [3, Lemma 1]):

**Lemma 3.** *If  $Y$  is a Hausdorff topological vector space, then for any  $\nu \in M(Y)$  there are only countably many one-dimensional affine subspaces  $S$  of  $Y$  satisfying the inequality  $\nu(S) > \delta(\nu)$ .*

*Proof.* Let  $\mathfrak{S}$  denote the class of all one-dimensional affine (closed, hence Borel measurable) subspaces  $S$  of  $Y$  and  $A$  be the set of points  $y \in Y$  with  $\nu(\{y\}) > 0$ . Let, moreover, the measure  $\nu_0$  be defined by  $d\nu_0 = 1_{X \setminus A} d\nu$ . Then  $S \in \mathfrak{S}$  satisfies  $\nu(S) \leq \delta(\nu)$ , whenever (a)  $S$  intersects  $A$  in at most one point and (b)  $S$  is a  $\nu_0$ -null set. But there are only countably many exceptions of (a), because  $A$  is countable, and this holds as well for (b), because two sets  $S \in \mathfrak{S}$  meet in at most one point and thus are  $\nu_0$ -almost disjoint.  $\square$

## MAIN RESULTS

No further auxiliary results on nonatomic measures (as in Section 3 of [6]) are needed to establish the following extension of Wulbert's main result:

**Proposition 1.** *Let  $\mu$  be a nonatomic  $\sigma$ -finite Baire measure on some topological space  $X$  and define the topology in  $\mathcal{C}(X)$  by the (finite or infinite) uniform norm. Then*

$$\mathcal{C}_0 := \{h \in \mathcal{C}(X) : \mu(\{x : h(x) = y\}) = 0 \text{ for all } y \in \mathbf{R}\}$$

*is a dense subset of  $\mathcal{C}(X)$ .*

*Proof.* 1. Since  $\mu$  may be replaced by any dominating measure, it is sufficient to consider the case of a probability measure. Since, moreover,  $\mu$  is nonatomic, there exists a Baire measurable function  $h : X \rightarrow [0, 1]$  with

$$\mu(\{x : h(x) = y\}) = 0 \quad \text{for } 0 \leq y \leq 1.$$

Let  $h_n \in \mathcal{C}(X), n \in \mathbf{N}$ , be an  $L_1$ -approximation of  $h$ , i.e. satisfying

$$\lim_{n \rightarrow \infty} \int_X |h_n - h| d\mu = 0,$$

where  $0 \leq h_n \leq 1$  may and will be assumed. Then  $h_n \rightarrow h$  in  $\mu$ -measure and thus the distributions  $\mu \circ h_n^{-1}$  converge weakly to  $\mu \circ h^{-1}$ . Since this distribution is nonatomic, Lemma 2 yields

$$\limsup_{n \rightarrow \infty} \delta(\mu \circ h_n^{-1}) \leq \delta(\mu \circ h^{-1}) = 0.$$

2. Now, with  $f \in \mathcal{C}(X)$  and  $n \in \mathbf{N}$  being fixed, define

$$g_t := f + t h_n \in \mathcal{C}(X) \quad \text{for } t \in \mathbf{R}.$$

Then an application of Lemma 3 to  $Y = \mathbf{R}^2$  and the joint distribution  $\nu$  of  $(f, h_n)$  shows the inequality  $\delta(\mu \circ g_t^{-1}) \leq \delta(\nu)$  to hold, except for countably many values of  $t$ . Combined with the trivial inequality  $\delta(\nu) \leq \delta(\mu \circ h_n^{-1})$  this provides a dense subset  $T$  of  $\mathbf{R}$  such that

$$\delta(\mu \circ g_t^{-1}) \leq \delta(\mu \circ h_n^{-1}) \quad \text{for } t \in T.$$

3. Next, consider the sets

$$\mathcal{G}_n := \left\{ g \in \mathcal{C}(X) : \delta(\mu \circ g^{-1}) < \delta(\mu \circ h_n^{-1}) + \frac{1}{n} \right\} \quad \text{for } n \in \mathbf{N}.$$

Since the mapping  $\mathcal{C}(X) \ni g \mapsto \mu \circ g^{-1} \in M(\mathbf{R})$  is continuous, they are open subsets of  $\mathcal{C}(X)$  by Lemma 2. Since, moreover, they are dense in  $\mathcal{C}(X)$  by part 2 of the proof and  $\mathcal{C}(X)$  is a Baire space, the set  $\bigcap_{n \in \mathbf{N}} \mathcal{G}_n$  is as well dense in  $\mathcal{C}(X)$ . Since, finally, this intersection equals  $\mathcal{C}_0$  by part 1 of the proof, the assertion is established.\* □

Now the concluding criterion is an easy consequence:

**Proposition 2.** *A Baire measure  $\mu$  on a topological space  $X$  allows a continuous transformation  $\varphi$  into Lebesgue measure on the interval  $[0, \mu(X)]$  resp.  $\mathbf{R}_+$  if and only if it is nonatomic and moderated.*

*Proof.* The condition is necessary, as is clear for “nonatomic” and follows for “moderated” from Lemma 1. To prove sufficiency choose a function  $f \in \mathcal{C}(X)$  according to Lemma 1 and a function  $h \in \mathcal{C}_0$  with  $\|h - (\frac{1}{f} + 1)\| < 1$  according to Proposition 1. Then  $h \geq 0$ , and it follows from

$$\mu(\{x : h(x) \leq y\}) \leq \mu\left(\left\{x : f(x) \geq \frac{1}{y}\right\}\right) < \infty \quad \text{for } y > 0,$$

that the “distribution function”

$$\psi(y) := \mu(\{x : h(x) \leq y\}) \quad \text{for } y \geq 0$$

---

\* The original proof of Proposition 1 used arguments from measure theory, resulting in

$$\delta(\mu \circ (f + \sum_{n \in \mathbf{N}} \frac{t_n}{2^n} h_n)^{-1}) = 0 \quad \text{for } \lambda^{\mathbf{N}}\text{-almost all } (t_n, n \in \mathbf{N}) \in [0, 1]^{\mathbf{N}}.$$

Following a suggestion by the referee, it was slightly simplified by using topological arguments.

is continuous and finite-valued. Using the “inverse function”

$$\psi^{-1}(z) := \max \{y : \psi(y) \leq z\} \quad \text{for } 0 \leq z < \mu(X),$$

it is now easily checked that

$$\mu(\{x : \psi \circ h(x) \leq y\}) = y \quad \text{for } 0 \leq y < \mu(X),$$

i.e. the continuous function  $\varphi := \psi \circ h$  is appropriate.  $\square$

Finally, it should be mentioned that for infinite  $\mu$  a continuous transformation into Lebesgue measure on the whole real line is as well possible. Indeed, to replace  $X = \mathbf{R}_+$  by  $Y = \mathbf{R}$ , it suffices to choose for the first step a continuous function  $h$  with the properties

$$\lambda(\{x : |h(x)| = y\}) = 0 \quad \text{and} \quad \lambda(\{x : |h(x)| \leq y\}) < \infty \quad \text{for all } y \geq 0,$$

satisfying in addition

$$\lambda(\{x : h(x) \leq 0\}) = \infty = \lambda(\{x : h(x) \geq 0\})$$

(take for instance  $h(x) = x^2 \sin x$ ). The second step then uses the “two-sided” distribution function  $\psi$ , being defined by  $\lambda(\{x : 0 \leq h(x) \leq y\})$  for  $y \geq 0$  and by  $-\lambda(\{x : y \leq h(x) \leq 0\})$  for  $y \leq 0$ , respectively.

#### ADDED IN PROOF

As the referee asked for a direct solution, here is a continuous transformation  $\varphi$  of Lebesgue measure on  $\mathbf{R}_+$  into Lebesgue measure on  $\mathbf{R}$ : define mappings  $\varphi_n : [0, 1] \mapsto \mathbf{R}_+$ ,  $n \in \mathbf{N}$ , by affine interpolation of the points

$$(0, 0), \left(\frac{1}{2(n+1)}, n\right), \left(\frac{1}{2}, n+1\right), \left(1 - \frac{1}{2(n+1)}, n\right), (1, 0)$$

and let  $\varphi$  be composed of the translates of  $\varphi_n$  resp.  $-\varphi_n$  to the intervals  $[2n - 2, 2n - 1]$  resp.  $[2n - 1, 2n]$ .

#### REFERENCES

1. Hobby, C.R. and J.R. Rice, *A moment problem in  $L_1$ -approximation*, Proc. Amer. Math. Soc. 16(1965)665-670. MR **31**:2550
2. Kellerer, H.G., *Zur Existenz analoger Bereiche*, Z. Wahrscheinlichkeitstheorie Verw. Geb. 1(1963)240-246. MR **28**:182
3. Kellerer, H.G., *A topological version of Liapunov's theorem*, Arch. Math. 72(1999) 206-213. MR **2000c**:28019
4. Phelps, R.R., *Uniqueness of Hahn-Banach extensions and unique best approximation*, Trans. Amer. Math. Soc. 95(1960)238-255. MR **22**:3964
5. Wulbert, D.E., *Liapunov's and related theorems*, Proc. Amer. Math. Soc. 108(1990) 955-960. MR **90m**:46074
6. Wulbert, D.E., *Annihilating a subspace of  $L_1$  with the sign of a continuous function*, Proc. Amer. Math. Soc. 128(2000)2431-2438. MR **2000m**:46036

MATHEMATISCHES INSTITUT DER UNIVERSITÄT MÜNCHEN, THERESIENSTRASSE 39, D-80333 MUNICH, GERMANY

*E-mail address:* [hans\\_g.kellerer@mathematik.uni-muenchen.de](mailto:hans_g.kellerer@mathematik.uni-muenchen.de)