

ON WAVELETS INTERPOLATED FROM A PAIR OF WAVELET SETS

ZIEMOWIT RZESZOTNIK AND DARRIN SPEEGLE

(Communicated by David R. Larson)

ABSTRACT. We show that any wavelet, with the support of its Fourier transform small enough, can be interpolated from a pair of wavelet sets. In particular, the support of the Fourier transform of such wavelets must contain a wavelet set, answering a special case of an open problem of Larson. The interpolation procedure, which was introduced by X. Dai and D. Larson, allows us also to prove the extension property.

1. INTRODUCTION

The L^2 theory of orthonormal wavelets is not fully developed yet. There are three ways of understanding the whole set of wavelets in $L^2(\mathbb{R})$. The first one relies on characterization equations introduced by Y. Meyer (see [Le], [BSW], [Gr], [HW]) or those conjectured by G. Weiss (see [B], [WW]). The second one is based on the idea of a generalized multiresolution analysis introduced by L. Baggett, H. Medina and K. Merrill in [BMM]. The final one, described by X. Dai and D. Larson in [DL], approaches wavelets from the operator theory point of view by treating them as wandering vectors. One of the advantages of this approach is the interpolation procedure, which allows a construction of “new” wavelets from “old” wavelets. The procedure is particularly simple in the case when the “old” wavelets are of the MSF type. As a result, these elementary MSF wavelets can be viewed as “building blocks” of a huge class of wavelets. Even the situation when we use only two wavelet sets to construct new wavelets is very interesting. [DL] contains a detailed study of this basic interpolation. For example, it is shown that Meyer’s original construction of wavelets follows naturally from this point of view. The construction of Meyer was based on the characterization equations mentioned above. In this paper we establish a further connection between these equations and wavelets interpolated from a pair of wavelet sets. To be more precise, we use the equations to prove that if the support of $\hat{\psi}$ is “small”, then the wavelet ψ can be interpolated from a pair of wavelet sets. To exhibit the usefulness of this basic interpolation, we apply it to prove that the local behavior of the Fourier transform $\hat{\psi}$ of a wavelet ψ is as arbitrary as possible at any point on the real line, except the origin. This is closely related to [BGRW], where it was shown that the origin is a point with a special meaning for $\hat{\psi}$.

Received by the editors September 19, 2000 and, in revised form, March 22, 2001.

2000 *Mathematics Subject Classification*. Primary 42C40.

Key words and phrases. Orthonormal wavelets, MSF wavelets, interpolated wavelets.

2. INTERPOLATION PAIRS OF WAVELET SETS

Following [DL], we shall define an interpolation pair of wavelet sets. Let E and F be wavelet sets. Since both these sets partition \mathbb{R} under translations by $2k\pi$, $k \in \mathbb{Z}$, we can consider a natural mapping $\sigma : E \rightarrow F$ such that $\sigma(\xi) = \xi + 2k\pi$, where $k \in \mathbb{Z}$. The *interpolation map* σ_E^F is the unique extension of σ to \mathbb{R} such that

$$(2.1) \quad \sigma_E^F(\xi) = 2^j \sigma_E^F(2^{-j}\xi) \quad \text{for } \xi \in 2^j E, \quad j \in \mathbb{Z}.$$

That the extension exists and is unique follows from the fact that both E and F partition \mathbb{R} under dilations by 2^j , $j \in \mathbb{Z}$. The ordered pair (E, F) is called an *interpolation pair* if $(\sigma_E^F)^2$ is equal to the identity. If that is the case, then every function ψ given by

$$\hat{\psi} = h_1 \mathbf{1}_E + h_2 \mathbf{1}_F,$$

where h_1 and h_2 are 2-dilation-periodic, is an orthonormal wavelet¹ provided that the matrix

$$(2.2) \quad \begin{bmatrix} h_1 & h_2 \\ h_2 \circ \sigma_E^F & h_1 \circ \sigma_E^F \end{bmatrix}$$

is unitary a.e. on E . We refer to such wavelets as wavelets interpolated from a pair (E, F) . In this paper, an interpolation pair will be called *non-trivial* if $E \neq F$. We note here that the trivial case corresponds to MSF wavelets.

The question we are concerned about is: “How big is the class of wavelets interpolated from a pair of wavelet sets?” Although this question is pretty vague, we shall try to give a precise answer to it.

In [La] it is shown that if (E, F) is an interpolation pair and ψ is a wavelet such that the support of $\hat{\psi}$ is contained in $E \cup F$, then ψ can be interpolated from the pair (E, F) . Therefore, at least in the case of interpolation pairs, it is enough to concentrate on the support of $\hat{\psi}$ to find out if a given wavelet ψ comes from an interpolation; however, left open is the question of how to know whether the support of $\hat{\psi}$ is contained in the union of an interpolation pair of wavelet sets. This motivates the following approach. Let $\psi \in L^2(\mathbb{R})$ be a wavelet. Denote the support of $\hat{\psi}$ by K and define two maps

$$T_K(\xi) = \sum_{k \in \mathbb{Z}} \mathbf{1}_K(\xi + 2k\pi)$$

and

$$D_K(\xi) = \sum_{j \in \mathbb{Z}} \mathbf{1}_K(2^j \xi),$$

where $\xi \in \mathbb{R}$. The maps are defined a.e., and the supremum norms $\|T_K\|_\infty$, $\|D_K\|_\infty$ determine the “size” of the set K . To see how this notion operates let us recall the Meyer–Weiss equations characterizing wavelets (see [HKLS]):

$$(2.3) \quad \sum_{k \in \mathbb{Z}} |\hat{\psi}(\xi + 2k\pi)|^2 = 1,$$

¹ We use the Fourier transform defined by

$$\hat{\psi}(\xi) = \int_{\mathbb{R}} \psi(x) e^{-ix\xi} dx.$$

$$(2.4) \quad \sum_{k \in \mathbb{Z}} \hat{\psi}(\xi + 2k\pi) \overline{\hat{\psi}(2^j(\xi + 2k\pi))} = 0 \quad \text{for } j \geq 1,$$

$$(2.5) \quad \sum_{j \in \mathbb{Z}} |\hat{\psi}(2^j \xi)|^2 = 1,$$

$$(2.6) \quad t_q(\xi) := \sum_{j=0}^{\infty} \hat{\psi}(2^j \xi) \overline{\hat{\psi}(2^j(\xi + 2q\pi))} = 0, \quad \text{for } q \in 2\mathbb{Z} + 1.$$

A fundamental result in the L^2 theory of wavelets is that a function $\psi \in L^2$ is a wavelet if and only if $\|\psi\|_2 = 1$ and equations (2.5) and (2.6) hold. Moreover, in this case, (2.3) and (2.4) also hold. These facts were proven by various authors in different generalities; see [HW] for details.

From equations (2.3) and (2.5) it follows that the case $\|T_K\|_\infty = \|D_K\|_\infty = 1$ gives us the most elementary wavelets, that is, MSF wavelets (even if only one of the norms is equal to 1, then ψ must be an MSF wavelet). We will prove that the second simplest case (that is, $\|T_K\|_\infty = \|D_K\|_\infty = 2$) corresponds exactly to wavelets interpolated from a non-trivial pair of wavelet sets. Of course, in light of [La], we can achieve this by focusing on the support of the Fourier transform of a wavelet.

Theorem 2.1. *Let $\psi \in L^2(\mathbb{R})$ be an orthonormal wavelet, and let K denote the support of $\hat{\psi}$. There exists a non-trivial interpolation pair (E, F) such that $K = E \cup F$ if and only if $\|T_K\|_\infty = \|D_K\|_\infty = 2$.*

Proof. Only one implication of the above statement requires a proof (the other one follows immediately from the fact that E and F are wavelet sets and $E \neq F$). The real task is to decompose K into the union of two wavelet sets, provided that K is the support of the Fourier transform of a wavelet ψ and T_K and D_K are bounded by 2 (proving that these wavelet sets will form an interpolation pair will not be difficult).

The proof consists of several easily-justified claims.

Claim A. $T_K(\xi) = D_K(\xi)$ for a.e. $\xi \in K$.

Proof of Claim A. To prove Claim A we observe that T_K and D_K can only attain values 1 and 2 on \mathbb{R} (value 0 is restricted because of (2.3) and (2.5)). From equation (2.3) it follows that $\{\xi \in \mathbb{R} : |\hat{\psi}(\xi)| = 1\} = \{\xi \in K : T_K(\xi) = 1\}$. Similarly, equation (2.5) asserts that $\{\xi \in \mathbb{R} : |\hat{\psi}(\xi)| = 1\} = \{\xi \in K : D_K(\xi) = 1\}$. Thus, Claim A is true. □

Let $R := \{\xi \in K : T_K(\xi) = D_K(\xi) = 1\}$ (we know already that $R = \{\xi \in \mathbb{R} : |\hat{\psi}(\xi)| = 1\}$). From Claim A it follows that if $\xi \in K \setminus R$, then there is a unique integer $k \neq 0$ such that $\xi + 2k\pi \in K \setminus R$, and also a unique integer $j \neq 0$ such that $2^j \xi \in K \setminus R$. Thus, we can define a map $\tau_K : K \setminus R \rightarrow K \setminus R$ by setting

$$\tau_K(\xi) = \xi + 2k\pi,$$

where $k \in \mathbb{Z} \setminus \{0\}$ (k depends on ξ) and a map $d_K : K \setminus R \rightarrow K \setminus R$ by setting

$$d_K(\xi) = 2^j \xi,$$

where $j \in \mathbb{Z} \setminus \{0\}$ (j depends on ξ).

Claim B. τ_K and d_K commute.

Proof of Claim B. We prove Claim B by applying equation (2.6). Let $\xi \in K \setminus R$, $\tau_K(\xi) = \xi + 2k\pi$ and $d_K(\xi) = 2^j\xi$. Clearly, $k = 2^nq$ for some $n \in \mathbb{N} \cup \{0\}$ and $q \in 2\mathbb{Z} + 1$. Since, by (2.6), $t_q(2^{-n}\xi) = 0$, we have

$$(2.7) \quad \sum_{l=0}^{\infty} \hat{\psi}(2^{l-n}\xi) \overline{\hat{\psi}(2^{l-n}(\xi + 2k\pi))} = 0.$$

It is easy to see that the above sum contains a non-zero term. Indeed, the term corresponding to $l = n$ is simply $\hat{\psi}(\xi) \overline{\hat{\psi}(\xi + 2k\pi)}$, which cannot be zero, since both ξ and $\xi + 2k\pi$ are in K . Thus, to make the whole sum equal to 0 we need to have at least one more non-zero term. On the other hand, since $\xi \in K$ and $D_K \leq 2$, we conclude that we can have at most one more non-zero term, the one corresponding to $l = j + n$. Therefore, (2.7) reduces to

$$\hat{\psi}(\xi) \overline{\hat{\psi}(\xi + 2k\pi)} + \hat{\psi}(2^j\xi) \overline{\hat{\psi}(2^j(\xi + 2k\pi))} = 0.$$

The above allows us to draw two conclusions: $2^j(\xi + 2k\pi)$ belongs to $K \setminus R$ and $2^jk = 2^lq$ is a non-zero integer. Thus, we have

$$d_K \circ \tau_K(\xi) = d_K(\xi + 2k\pi) = 2^j(\xi + 2k\pi) = 2^j\xi + 2^lq\pi = \tau_K(2^j\xi) = \tau_K \circ d_K(\xi),$$

which concludes the proof of the claim. \square

The fact that the maps τ_K and d_K commute can be stated as: “the dilation brother of my translation brother is the translation brother of my dilation brother” and represented by the following diagram:

$$(2.8) \quad \begin{array}{ccc} \xi & \xrightarrow{\tau_K} & \xi + 2k\pi \\ d_K \downarrow & & d_K \downarrow \\ 2^j\xi & \xrightarrow{\tau_K} & 2^j(\xi + 2k\pi), \end{array}$$

where $2^jk \in \mathbb{Z} \setminus \{0\}$. The above diagram, or rather its simplicity, is the main reason why we can decompose K into a union of two wavelet sets. Of course, the set R must be a part of these wavelet sets; thus we only need to investigate the remaining set $K \setminus R$.

For every $j, k \in \mathbb{Z} \setminus \{0\}$ we define $K_{j,k} := \{\xi \in K : \xi + 2k\pi \in K, 2^j\xi \in K\}$; that is,

$$(2.9) \quad K_{j,k} = K \cap (K - 2k\pi) \cap 2^{-j}K.$$

Since T_K and D_K are bounded by 2, it follows that the sets $K_{j,k}$ are mutually disjoint and

$$(2.10) \quad \bigcup_{j,k \in \mathbb{Z} \setminus \{0\}} K_{j,k} = K \setminus R.$$

Claim C. We have the following equalities:

$$K_{j,k} + 2k\pi = K_{j,-k},$$

$$2^j K_{j,k} = K_{-j,2^j k},$$

$$2^j(K_{j,k} + 2k\pi) = K_{-j,-2^j k}.$$

Proof of Claim C. Let $\xi \in K_{j,k}$. By diagram (2.8) we have $2^j(\xi + 2k\pi) \in K$. Therefore,

$$(2.11) \quad K_{j,k} \subset 2^{-j}K - 2k\pi.$$

From (2.9) and (2.11) it follows that

$$(2.12) \quad K_{j,k} = K \cap (K - 2k\pi) \cap 2^{-j}K \cap (2^{-j}K - 2k\pi).$$

Thus, since (2.12) holds for $-k$ as well,

$$K_{j,k} + 2k\pi = (K + 2k\pi) \cap K \cap (2^{-j}K + 2k\pi) \cap 2^{-j}K = K_{j,-k},$$

which proves the first equality of the claim. Applying (2.12) in the same manner yields the two remaining equalities. \square

Claim C allows us to obtain an analogous version of diagram (2.8):

$$(2.13) \quad \begin{array}{ccc} K_{j,k} & \xrightarrow[\substack{\tau\kappa \\ +2k\pi}]{} & K_{j,-k} \\ d\kappa \downarrow \cdot 2^j & & d\kappa \downarrow \cdot 2^j \\ K_{-j,2^j k} & \xrightarrow[\substack{\tau\kappa \\ +2^j 2k\pi}]{} & K_{-j,-2^j k}, \end{array}$$

where $2^j k \in \mathbb{Z} \setminus \{0\}$. Let $A_{j,k}$ denote the union of all the sets in the above diagram, that is,

$$A_{j,k} := K_{j,k} \cup K_{j,-k} \cup K_{-j,2^j k} \cup K_{-j,-2^j k},$$

where, to avoid redundancy, we consider only $j, k \in \mathbb{N}$. (\mathbb{N} does not contain 0.)

Claim D. The sets $A_{j,k}$ are mutually disjoint for all $j, k \in \mathbb{N}$, and

$$\bigcup_{j,k \in \mathbb{N}} A_{j,k} = K \setminus R.$$

Proof of Claim D. The disjointness follows from the fact that the sets $K_{j,k}$ are mutually disjoint. By (2.10) we have

$$(2.14) \quad K \setminus R = \bigcup_{j,k \in \mathbb{N}} K_{j,k} \cup \bigcup_{j,k \in \mathbb{N}} K_{j,-k} \cup \bigcup_{j,k \in \mathbb{N}} K_{-j,k} \cup \bigcup_{j,k \in \mathbb{N}} K_{-j,-k}.$$

However, if we consider the set $K_{-j,k}$, where $j \in \mathbb{N}$ and $k \in \mathbb{Z} \setminus \{0\}$, then according to diagram (2.13) we must have $2^{-j}k \in \mathbb{Z} \setminus \{0\}$. Therefore, it follows that either $K_{-j,k}$ is empty or $k = 2^j k'$ for some non-zero integer k' . This observation allows us to reduce the last two unions in (2.14), which leads to

$$K \setminus R = \bigcup_{j,k \in \mathbb{N}} K_{j,k} \cup \bigcup_{j,k \in \mathbb{N}} K_{j,-k} \cup \bigcup_{j,k' \in \mathbb{N}} K_{-j,2^j k'} \cup \bigcup_{j,k' \in \mathbb{N}} K_{-j,-2^j k'} = \bigcup_{j,k \in \mathbb{N}} A_{j,k}$$

and proves our claim. \square

At this point we are ready to construct the wavelet sets. The intersection with $K \setminus R$ of one of them is obtained by unioning all the terms on the diagonal of diagram (2.13) for $j, k \in \mathbb{N}$, and the intersection of the other one by unioning all the terms on the antidiagonal. Thus, we define

$$E' := \bigcup_{j,k \in \mathbb{N}} (K_{j,k} \cup K_{-j,-2^j k}), \quad F' := \bigcup_{j,k \in \mathbb{N}} (K_{j,-k} \cup K_{-j,2^j k}),$$

and

$$E := E' \cup R, \quad F := F' \cup R.$$

Claim D simply assures that

$$(2.14) \quad K = E' \cup F' \cup R$$

and that the union is disjoint.

Claim E. E and F are wavelet sets.

Proof of Claim E. From diagram (2.13) it follows that both sets are translation and dilation congruent to each other; therefore, it is enough to prove that E is a wavelet set.

From diagram (2.13) it follows that

$$(2.15) \quad K \cap \bigcup_{n \in \mathbb{Z} \setminus \{0\}} (E + 2n\pi) = F' \quad \text{and} \quad K \cap \bigcup_{n \in \mathbb{Z} \setminus \{0\}} 2^n E = F'.$$

Thus, since $E \subset K$, it follows that $\emptyset = E \cap F' = E \cap K \cap \bigcup_{n \in \mathbb{Z} \setminus \{0\}} (E + 2n\pi) = E \cap \bigcup_{n \in \mathbb{Z} \setminus \{0\}} (E + 2n\pi)$, and the 2π -translations of E are disjoint. Similarly, the 2-dilations of E are disjoint. Moreover, by (2.14) and (2.15) we obtain

$$K \cap \bigcup_{n \in \mathbb{Z}} (E + 2n\pi) = K \quad \text{and} \quad K \cap \bigcup_{n \in \mathbb{Z}} 2^n E = K.$$

Since T_K and D_K are bounded below by 1, this proves that 2π -translations, as well as 2-dilations, of E cover the real line. \square

Claim F. (E, F) is an interpolation pair.

Proof of Claim F. We need to prove that the interpolation map σ_E^F composed with itself gives us the identity. From (2.1) it follows that it is enough to show that $(\sigma_E^F)^2(\xi) = \xi$ for a.e. $\xi \in E$. Since σ_E^F restricted to R is the identity, we can focus on the case when $\xi \in E'$. Clearly, $\sigma_E^F(\xi) = \tau_K(\xi)$ for $\xi \in E'$. Moreover, the same is true if $\xi \in F'$. Indeed, from diagram (2.13) it follows that for such ξ 's we have $\sigma_E^F(\xi) = d_K \circ \tau_K \circ d_K(\xi) = \tau_K(\xi)$. Thus, σ_E^F restricted to $K \setminus R$ is equal to τ_K , which shows that $(\sigma_E^F)^2 = id$. \square

Since we proved that (E, F) is an interpolation pair and, by (2.14), $K = E \cup F$, this concludes the proof of the theorem. \square

As an immediate consequence of Theorem 2.1 and the remarks preceding it, we obtain the following characterization.

Corollary 2.2. *If $\psi \in L^2(\mathbb{R})$ is an orthonormal wavelet and K denotes the support of $\hat{\psi}$, then ψ is interpolated from a pair of wavelet sets (E, F) if and only if $\|T_K\|_\infty$ and $\|D_K\|_\infty$ are bounded above by 2. Moreover, if this is the case, then either $\|T_K\|_\infty = \|D_K\|_\infty = 2$, in which case $K = E' \cup F'$, where (E', F') is a non-trivial interpolation pair, or $\|T_K\|_\infty = \|D_K\|_\infty = 1$, in which case ψ is an MSF wavelet.*

3. EXTENSION PROPERTY

The behavior at the origin of the Fourier transform of wavelets was studied in [BGRW]. The starting point of this investigation was the observation that, if ψ is a wavelet and we set $\hat{\psi}(0) = 0$, then the origin becomes a Lebesgue point of the function $\hat{\psi}$. It turns out that the origin is the only point on the real line at which the behavior of $\hat{\psi}$ is limited. We shall prove that for any other point $\xi_0 \in \mathbb{R}$ there exists a neighborhood U such that every measurable function b on U with $|b| \leq 1$ can be extended to the real line in such a way that the extension is the Fourier transform of a wavelet (this result was already announced in [BGRW] as Theorem (1.6); the case $|b| = 1$ was proven in [IP]). This will be achieved by finding an open interval $U \ni \xi_0$, small enough to be a subset of a wavelet set E , which will form an interpolation pair with another wavelet set F . When this is done, we shall use interpolation based on (E, F) to find a wavelet ψ such that $\hat{\psi}|_U = b$. This last step of the proof will be familiar to the reader who is accustomed to Meyer’s construction of wavelets, which was mentioned before, or its generalizations, presented both in [HWW] and [DL].

We start by exhibiting a special class of interpolation pairs which we shall use for our construction; similar classes were considered in [Gu].

Proposition 3.1. *Let T and U be measurable subsets of \mathbb{R} , and let both $n \in \mathbb{N}$ and $k \in \mathbb{Z}$ be given. If $E := U \cup (2^{-n}U - 2k\pi) \cup T$ is a wavelet set and the union is disjoint, then $F := 2^{-n}U \cup (U - 2^n 2k\pi) \cup T$ is a wavelet set and the union is disjoint. Moreover, (E, F) is an interpolation pair.*

Proof. If the sets U , $2^{-n}U - 2k\pi$ and T are pairwise disjoint and their union is a wavelet set, then the sets $2^{-n}U$, $U - 2^n 2k\pi$ and T must be mutually disjoint as well. Therefore, F is translation and dilation congruent to E , which shows that F is a wavelet set. We shall prove that (E, F) is an interpolation pair. As we mentioned before, it is enough to show that the interpolation map σ_E^F satisfies $(\sigma_E^F)^2(\xi) = \xi$ for $\xi \in E$. Computing σ_E^F on E yields

$$\sigma_E^F(\xi) = \begin{cases} \xi & \text{for } \xi \in T, \\ \xi - 2^n 2k\pi & \text{for } \xi \in U, \\ \xi + 2k\pi & \text{for } \xi \in 2^{-n}U - 2k\pi. \end{cases}$$

Therefore, on F , σ_E^F has the form

$$\sigma_E^F(x) = \begin{cases} \xi & \text{for } \xi \in T, \\ 2^{-n}\sigma_E^F(2^n\xi) = \xi - 2k\pi & \text{for } \xi \in 2^{-n}U, \\ 2^n\sigma_E^F(2^{-n}\xi) = \xi + 2^n 2k\pi & \text{for } \xi \in U - 2^n 2k\pi. \end{cases}$$

So, it is easily checked that for $\xi \in E$, $(\sigma_E^F)^2(\xi) = \xi$. □

Although k given in the above proposition can be any integer, we will only need to consider $k = 1$ to proceed with our construction. As a technical lemma we shall prove that if $\xi_0 > 0$, then there is a wavelet set E , given as in Proposition 3.1, such that U is an open interval containing ξ_0 . We also would like to point out that, according to [BGRW], if ψ is an MRA wavelet and m is an arbitrary integer, then $4m\pi$ is a Lebesgue point of $\hat{\psi}$, provided $\hat{\psi}(4m\pi) = 0$. This indicates that the case when $\xi_0 = 4m\pi$ may be harder to consider. Indeed, as we will see, the most general case $\xi_0 \neq 2m\pi$ can be treated in a simple way based on the approach offered in

[DLS]. The special case $\xi_0 = 2m\pi$ requires more advanced techniques described in [IP]. Both of these methods rely on introducing the maps $\tau : \mathbb{R} \rightarrow [-\pi, \pi)$ defined by $\tau(\xi) = \xi + 2k\pi$ (where k is the unique integer such that $\xi + 2k\pi \in [-\pi, \pi)$) and $d : \mathbb{R} \setminus \{0\} \rightarrow [-2\pi, -\pi) \cup [\pi, 2\pi)$ defined by $d(\xi) = 2^j\xi$ (where j is the unique integer such that $2^j\xi \in [-2\pi, -\pi) \cup [\pi, 2\pi)$). Ionascu and Pearcy [IP] proved that a measurable set V is a subset of a wavelet set E if and only if there exist $A \supset V$ and $B \supset V$ such that $\tau|_A, \tau|_B, d|_A$ and $d|_B$ are injections, $\tau(A) = [-\pi, \pi)$ and $d(B) = [-2\pi, -\pi) \cup [\pi, 2\pi)$. Moreover, in this case, the wavelet set E may be chosen as a subset of $A \cup B$. A sufficient condition for a set V to be contained in a wavelet set was previously noted in [DLS], namely, $\tau|_V$ and $d|_V$ are injective, $d(V) \neq [-2\pi, -\pi) \cup [\pi, 2\pi)$ and there exists a $\delta > 0$ such that $\tau(V) \cap [-\delta, \delta] = \emptyset$. We shall use these two results to prove the technical lemma.

Lemma 3.2. *For every $\xi_0 > 0$ there exist an open interval $U \ni \xi_0$, a measurable set $T \subset \mathbb{R}$ and some natural n such that $E := U \cup (2^{-n}U - 2\pi) \cup T$ is a wavelet set and the union is disjoint.*

Proof. Case 1: $\xi_0 \neq 2m\pi$ for any $m \in \mathbb{N}$. Let U be an open neighborhood of ξ_0 such that $2U \cap U = \emptyset$ and $2k\pi \notin U$ for any $k \in \mathbb{Z}$. Write $U = (2k\pi + \epsilon_1, 2(k+1)\pi - \epsilon_2)$, where $\epsilon_1, \epsilon_2 > 0$ and $k \in \mathbb{Z}$. Pick $n \in \mathbb{N}$ such that $2^{-n}U \subset [0, \epsilon_1] \cap [0, \pi/2)$ and denote the disjoint union $U \cup (2^{-n}U - 2\pi)$ by V . Notice that $\tau|_V$ is 1-1, $d|_V$ is 1-1, $\tau(V) \cap [-\delta, \delta] = \emptyset$ for $\delta = \min(2^{-n}(2k\pi + \epsilon_1), \epsilon_2)$, and $d(V) \neq [-2\pi, -\pi) \cup [\pi, 2\pi)$. Hence, by [DLS], there is a $T \subset \mathbb{R}$ such that $V \cup T$ is a wavelet set, as desired.

Case 2: $\xi_0 = 2m\pi$ for some $m \in \mathbb{N}$. For $m = 1$ we can choose $U = (4\pi/3, 8\pi/3)$, $T = \emptyset$ and $n = 1$. For $m \neq 1$, choose $n \in \mathbb{N}$ such that $2^{-n}\xi \in (2\pi/3, 4\pi/3)$, and choose U such that $2^{-n}U \subset (2\pi/3, 4\pi/3)$ and $U \subset (2m\pi - \pi/3, 2m\pi + \pi/3)$. Notice that $\tau(2^{-n}U) \subset (2\pi/3, \pi) \cup [-\pi, -2\pi/3)$ and $\tau(U) \subset (-\pi/3, \pi/3)$, so, setting $V = U \cup (2^{-n}U - 2\pi)$, we see that $\tau|_V$ and $d|_V$ are 1-1.

We wish to apply the theorem from [IP] which characterizes sets which are subsets of wavelet sets. In order to do so, we must find sets T_1 and T_2 (both disjoint from V) such that the restrictions of τ and d to $T_1 \cup V$ and $T_2 \cup V$ are 1-1, $\tau(T_1 \cup V) = [-\pi, \pi)$ and $d(T_2 \cup V) = [-2\pi, -\pi) \cup [\pi, 2\pi)$.

We simply note that $T_1 = [2(m-1)\pi + \pi/3, 2m\pi + \pi/3) \cap U^c \cap (2^{-n}U + 2(m-1)\pi)^c$ and $T_2 = 1/2([2\pi/3, 4\pi/3) \cap (2^{-n}U)^c) \cup 1/2([-4\pi/3, -2\pi/3) \cap (2^{-n}U - 2\pi)^c)$ satisfy the conditions above. Therefore, by Ionascu and Pearcy's characterization, there is a set $T \subset T_1 \cup T_2$ such that the disjoint union $U \cup (2^{-n}U - 2\pi) \cup T$ is a wavelet set. \square

From Proposition 3.1 it follows that the set $E = U \cup (2^{-n}U - 2\pi) \cup T$ from Lemma 3.2 and the set $F = 2^{-n}U \cup (U - 2^n 2\pi) \cup T$ form an interpolation pair. The standard interpolation procedure allows us to prove the main result of this section, which generalizes the special case $|b| = 1$ proven in [IP].

Theorem 3.3. *If $\xi_0 \neq 0$, then there exists a neighborhood U of ξ_0 with the property that if b is any measurable function on U with $|b| \leq 1$, then we can find a wavelet ψ such that $\hat{\psi}(\xi) = b(\xi)$ for all $\xi \in U$.*

Proof. First we observe that if ψ is a wavelet, then $\hat{\psi}(\cdot)$ is the Fourier transform of a wavelet (this follows from the Meyer-Weiss equations). Therefore, we can restrict our attention to the case $\xi_0 > 0$.

Let U , T and n be as in Lemma 3.2. We will interpolate from the pair (E, F) , where $E = U \cup (2^{-n}U - 2\pi) \cup T$ and $F = 2^{-n}U \cup (U - 2^n 2\pi) \cup T$. In order to do

so we define b_1, b_2, b_3 and b_4 by

$$\begin{aligned} b_1(\xi) &= \sqrt{1 - |b(2^n \xi)|^2} \text{ for } \xi \in 2^{-n}U \text{ and } 0 \text{ otherwise,} \\ b_2(\xi) &= \sqrt{1 - b_1^2(\xi + 2\pi)} \text{ for } \xi \in 2^{-n}U - 2\pi \text{ and } 0 \text{ otherwise,} \\ b_3(\xi) &= \sqrt{1 - b_2^2(2^{-n}\xi)} \text{ for } \xi \in U - 2^n 2\pi \text{ and } 0 \text{ otherwise,} \\ b_4(\xi) &= 1 \text{ for } \xi \in T \text{ and } 0 \text{ otherwise.} \end{aligned}$$

A simple calculation shows that if we set $h_1(\xi) = e^{i\xi/2}(|b(\xi)| + b_2(\xi) + b_4(\xi))$ on E and $h_2(\xi) = e^{i\xi/2}(b_1(\xi) + b_3(\xi))$ on F and extend 2-dilation-periodically, then the matrix (2.2) is unitary on E and, therefore, ψ given by $\hat{\psi} = h_1 \mathbf{1}_E + h_2 \mathbf{1}_F$ is a wavelet. To finish the proof we observe that there exists a unimodular function ν defined on U such that $b(\xi) = \nu(\xi)e^{i\xi/2}|b(\xi)|$. Moreover, since U is an open interval whose measure does not exceed 2π , we can extend ν to a 2π -periodic unimodular function $\tilde{\nu}$. Then, by the Meyer-Weiss equations, $\tilde{\nu}(\xi)\hat{\psi}(\xi)$ is the Fourier transform of a wavelet and $\tilde{\nu}(\xi)\hat{\psi}(\xi) = b(\xi)$ for all $\xi \in U$, as desired. \square

We also point out that applying the above theorem to a point $\xi_0 = 4m\pi$, where m is a non-zero integer, and a bell-shaped smooth function b yields an example of a non-MRA wavelet which behaves nicely on the interval U . A similar result is proven in [BCM]; the authors became aware of it after submitting this paper.

4. CONCLUSION

We answered the question: “How big is the class of wavelets interpolated from a pair of wavelet sets?” in two ways. The proof of the extension property indicated that, at least locally, every wavelet can be approximated by those wavelets in the frequency domain. The characterization theorem, presented as Corollary 2.2, asserted that a minimal necessary condition on the support of the Fourier transform of an interpolated wavelet is also sufficient to assure that a wavelet comes from interpolation. In this sense the class is as big as possible. The main difficulty in proving this characterization was in finding a wavelet set contained in the support of the Fourier transform of a given wavelet. The problem of doing so in general was already formulated by D. Larson in [La], and its solution does not follow immediately from the present L^2 theory of wavelets.

ACKNOWLEDGMENTS

We would like to thank Professor Guido Weiss for bringing our attention to the extension property.

REFERENCES

[B] Bownik, M., *On Characterizations of multiwavelets in $L^2(\mathbb{R}^n)$* , Proc. Amer. Math. Soc. **129** (2001), 3263–3274. CMP 2001:16

[BCM] Baggett, L., Courter, J. and Merrill, K., *The construction of wavelets from generalized conjugate mirror filters in $L^2(\mathbb{R}^n)$* , preprint.

[BGRW] Brandolini L., Garrigós G., Rzeszutnik Z. and Weiss G., *The behaviour at the origin of a class of band-limited wavelets*, Contemporary Mathematics **247** (1999), 75–91. MR **2001b**:42045

[BMM] Baggett, L.W., Medina, H.A. and Merrill, K.D., *Generalized multiresolution analyses, and a construction procedure for all wavelet sets in \mathbb{R}^n* , J. Fourier. Anal. Appl. **5** (no.6) (1999), 563–573. MR **2001f**:42055

- [BSW] Bonami, A., Soria, F. and Weiss, G., *Band-limited wavelets*, Jour. of Geom. Anal. **3** no.6 (1993), 543–578. MR **94k**:42046
- [DL] Dai, X. and Larson, D.R., *Wandering vectors for unitary systems and orthogonal wavelets*, Mem. Amer. Math. Soc. **134** (no. 640) (1998). MR **98m**:47067
- [DLS] Dai, X., Larson, D.R. and Speegle, D., *Wavelet sets in \mathbb{R}^n , II*, Contemp. Math. **216** (1998), 15–40. MR **99d**:42054
- [Gr] Gripenberg, G., *A necessary and sufficient condition for the existence of a father wavelet*, Studia Math. **114** (1995), 207–226. MR **96d**:42049
- [Gu] Gu, Q., *On interpolation families of wavelet sets*, Proc. Amer. Math. Soc. **128** (2000), 2973–2979. MR **2000m**:42026
- [HKLS] Ha, Y., Kang, H., Lee, J. and Seo, J., *Unimodular wavelets for L^2 and the Hardy space H^2* , Michigan Math. J. **41** (1994), 345–361. MR **95g**:42050
- [HW] Hernández, E. and Weiss, G., *A first course on wavelets*, Studies in Advanced Mathematics, CRC Press, Boca Raton, FL, 1996. MR **97i**:42015
- [HWW] Hernández, E., Wang, X. and Weiss, G., *Smoothing minimally supported frequency wavelets. I*, J. Fourier Anal. Appl. **2** no. 4 (1996), 329–340. MR **97h**:42015
- [IP] Ionascu, E.J. and Percy, C.M., *On subwavelet sets*, Proc. Amer. Math. Soc. **126** (1998), 3549–3552. MR **99b**:42040
- [La] Larson, D.R., *Von Neumann algebras and wavelets*, Operator algebras and applications (Samos, 1996), 267–312, NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci., 495, Kluwer Acad. Publ., Dordrecht, 1997. MR **98g**:46091
- [Le] Lemarié-Rieusset, P.G., *Ondolettes à localisation exponentielle*, J. Math. Pure et Appl **67** (1988), 227–236. MR **89m**:42024
- [WW] Weiss G. and Wilson E.N., *The Mathematical Theory of Wavelets*, preprint.

INSTITUTE OF MATHEMATICS, UNIVERSITY OF WROCLAW, PL GRUNWALDZKI 2/4, 50-384 WROCLAW, POLAND

E-mail address: zioma@math.uni.wroc.pl

DEPARTMENT OF MATHEMATICS & COMPUTER SCIENCE, SAINT LOUIS UNIVERSITY, ST. LOUIS, MISSOURI 63103

E-mail address: speegled@slu.edu