

REMARKS ON GINZBURG’S BIVARIANT CHERN CLASSES

SHOJI YOKURA

(Communicated by Paul Goerss)

ABSTRACT. The convolution product is an important tool in the geometric representation theory. Ginzburg constructed the bivariant Chern class operation from a certain convolution algebra of Lagrangian cycles to the convolution algebra of Borel-Moore homology. In this paper we give some remarks on the Ginzburg bivariant Chern classes.

§1. INTRODUCTION

The basic references are V. Ginzburg’s survey article “Geometric methods in the representation theory of Hecke algebras and quantum groups” [G2], Chriss-Ginzburg’s book [CG] and H. Nakajima’s survey article “Quiver varieties and quantum affine algebras” (in Japanese) [N1] (cf. its original paper [N2]).

In [G1] Ginzburg introduced the notion of “bivariant” Chern class from the abelian group of certain Lagrangian cycles satisfying some special conditions to the Borel-Moore homology group and showed that it is “convolutive”, i.e., it commutes with the convolution product. The Ginzburg bivariant Chern class is, in short, expressed as the Chern-Schwartz-MacPherson class multiplied by the pullback of the Segre (cohomology) class of the nonsingular target variety. In this paper we make some remarks on the Ginzburg bivariant Chern classes and also we give a naïve “constructible function version” of the above Ginzburg’s result concerning convolution.

§2. RELATIVE CHERN-MATHER CLASS AND GINZBURG BIVARIANT CHERN CLASS

The construction or definition of the Ginzburg bivariant Chern class c^{biv} given in [G1] is not direct, but in his survey article [G2] he gives an explicit description of it. It assigns to a Lagrangian cycle, i.e., $\Lambda_Y, Y \subset X_1 \times X_2$, the *relative Chern-Mather class of the fibers of the projection* $p_Y : Y \rightarrow X_2$. The projection p_Y is the restriction of the projection $p_2 : X_1 \times X_2 \rightarrow X_2$ to the subvariety Y . Let $\nu : \widehat{Y} \rightarrow Y$ be the Nash blow-up and \widehat{TY} the tautological Nash tangent bundle over \widehat{Y} . Then

Received by the editors May 25, 2001 and, in revised form, July 6, 2001.

1991 *Mathematics Subject Classification*. Primary 14C17, 14F99, 55N35.

Key words and phrases. Bivariant theory, Chern-Schwartz-MacPherson class, constructible function, convolution.

The author was partially supported by Grant-in-Aid for Scientific Research (C) (No.12640081), the Japanese Ministry of Education, Science, Sports and Culture.

the above relative Chern-Mather class of the fibers of the projection $p_Y : Y \rightarrow X_2$ is

$$c^{\text{biv}}(\Lambda_Y) := i_{Y*} \nu_* \left(c(\widehat{TY} - \nu^* p_Y^* TX_2) \cap [\widehat{Y}] \right)$$

where $i_Y : Y \rightarrow X_1 \times X_2$ is the inclusion. A more concrete expression of this is

$$\begin{aligned} c^{\text{biv}}(\Lambda_Y) &= i_{Y*} \left(\frac{1}{p_Y^* c(TX_2)} \cap c^M(Y) \right) \\ &= p_2^* s(TX_2) \cap i_{Y*} c^M(Y). \end{aligned}$$

Here $s(TX_2)$ denotes the Segre class of the tangent bundle of the manifold X_2 and $c^M(Y)$ is the Chern-Mather class of Y .

Therefore, the Ginzburg bivariant Chern class can be simply extended to a S -variety X with S being nonsingular, i.e., any morphism $\pi : X \rightarrow S$ from a possibly singular variety X to a nonsingular variety S and defined for any constructible functions on the source variety. Thus we can define

$$c_*^\pi = \pi^* s(TS) \cap c_* : F(X) \rightarrow H_*(X, \mathbb{Q})$$

where $c_* : F(X) \rightarrow H_*(X; \mathbb{Q})$ is the usual Chern-Schwartz-MacPherson class transformation with rational coefficients [M]. In fact, furthermore, we can modify the base variety a bit more. For a local complete intersection variety S in a smooth variety M , we can take the virtual tangent bundle

$$TS := TM|_S - N_S M$$

where $N_S M$ is the normal bundle of S in M (see [F]).

Theorem (2.1). *For the category of S -varieties with S being a smooth variety or a local complete intersection in a smooth variety, $c_*^\pi : F(X) \rightarrow H_*(X; \mathbb{Q})$ is the unique natural transformation from the constructible functions to the homology theory satisfying the condition that for a smooth variety X*

$$c_*^\pi(\mathbb{1}_X) = c(T_\pi) \cap [X],$$

where $T_\pi := TX - \pi^* TS$ is the relative virtual tangent bundle. Namely, for any commutative diagram

$$\begin{array}{ccc} X_1 & \xrightarrow{f} & X_2 \\ \pi_1 \downarrow & & \downarrow \pi_2 \\ S & \xrightarrow{\text{id}_S} & S \end{array}$$

we have the following commutative diagram:

$$\begin{array}{ccc} F(X_1) & \xrightarrow{f_*} & F(X_2) \\ c_*^{\pi_1} \downarrow & & \downarrow c_*^{\pi_2} \\ H_*(X_1) & \xrightarrow{f_*} & H_*(X_2). \end{array}$$

□

This follows from MacPherson's theorem [M] and the projection formula, and the proof of the uniqueness is the same as in [M].

Now it is easy to see that, for any vector bundle \mathcal{E} given on the base variety S , we can replace the "twisting" cohomology $\pi^*s(TS)$ by the pullback of any characteristic class $c\ell(\mathcal{E})$ of the bundle \mathcal{E} , i.e., we can consider

$$c_*^{c\ell(\mathcal{E})} := \pi^*c\ell(\mathcal{E}) \cap c_* : F(X) \rightarrow H_*(X; \mathbb{Q}).$$

Note that in this situation the base variety S can be any variety, singular or non-singular. Then this is the unique natural transformation satisfying the smooth condition that for a smooth variety X we have

$$c_*^{c\ell(\mathcal{E})}(\mathbb{1}_X) = (\pi^*c\ell(\mathcal{E}) \cup c(TX)) \cap [X].$$

The requirement of S being smooth or a local complete intersection in a smooth variety is essential in the above theorem or the set up of Ginzburg. The most naive and interesting question is whether or not we can generalize the definition of the Ginzburg bivariant Chern class to the case when the base variety S is arbitrarily singular. Consider the following fiber squares:

$$\begin{array}{ccccc} \widehat{W} \times_S \widehat{S} & \xrightarrow{\widehat{\nu}_W} & W \times_S \widehat{S} & \xrightarrow{\widehat{f}_W} & \widehat{S} \\ \downarrow \widetilde{\nu}_S & & \downarrow \widetilde{\nu}_S & & \downarrow \nu_S \\ \widehat{W} & \xrightarrow{\nu_W} & W & \xrightarrow{f_W} & S, \end{array}$$

where $\nu_W : \widehat{W} \rightarrow W$ and $\nu_S : \widehat{S} \rightarrow S$ are the Nash blow-ups and the morphisms $\widetilde{\nu}_S, \widetilde{\nu}_W, \widetilde{\nu}_S$ and \widehat{f}_W are the projections from the corresponding fiber products. Then it is quite natural and reasonable to define the following homology class:

$$c_*^\pi(\mathbb{1}_W) := i_{W*} \nu_{W*} \widetilde{\nu}_{S*} \left(c(\widetilde{\nu}_S^* T\widehat{W} - \widehat{\nu}_W^* \widehat{f}_W^* TS) \cap [\widehat{W} \times_S \widehat{S}] \right),$$

which is also equal to

$$i_{W*} \widehat{\nu}_{S*} \widehat{\nu}_{W*} \left(c(\widehat{\nu}_S^* T\widehat{W} - \widehat{\nu}_W^* \widehat{f}_W^* TS) \cap [\widehat{W} \times_S \widehat{S}] \right).$$

Then our obvious question is whether Theorem (2.1) holds or not with this definition of $c_*^\pi(\mathbb{1}_W)$ for the base variety S being arbitrarily singular. It turns out that the answer is negative, which implies that the nonsingularity of the base variety S is essential in the set-up of Ginzburg, i.e., the existence of the (virtual) tangent bundle TS is essential.

To see this negative result, let us consider the following special case. Let $\nu : \widehat{S} \rightarrow S$ be the Nash blow-up and consider the diagram

$$\begin{array}{ccc} \widehat{S} & \xrightarrow{\nu} & S \\ \nu \downarrow & & \downarrow \text{id}_S \\ S & \xrightarrow{\text{id}_S} & S. \end{array}$$

Then we ask if the following diagram commutes:

$$\begin{CD} F(\widehat{S}) @>\nu_*>> F(S) \\ @Vc_*^\nu VV @VVc_*^{\text{id}_S}V \\ H_*(\widehat{S}) @>\nu_*>> H_*(S). \end{CD}$$

In particular, we ask if the following equality holds:

$$\nu_*c_*^\nu(\mathbb{1}_{\widehat{S}}) = c_*^{\text{id}_S}(\nu_*\mathbb{1}_{\widehat{S}}).$$

In fact, we can show the following:

Theorem (2.2). *Let S be a singular variety with one isolated singularity x_0 such that the Nash blow-up \widehat{S} of S is nonsingular and furthermore such that the Euler-Poincaré characteristic $\chi(\nu^{-1}(x_0))$ of the fiber of the singularity x_0 is not equal to 1. Then we have*

$$\nu_*c_*^\nu(\mathbb{1}_{\widehat{S}}) \neq c_*^{\text{id}_S}(\nu_*\mathbb{1}_{\widehat{S}}).$$

Proof. To define $c_*^\nu(\mathbb{1}_{\widehat{S}})$, we need the following fiber square:

$$(2.2.1) \quad \begin{CD} \widehat{S} \times_S \widehat{S} @>\widehat{\nu}>> \widehat{S} \\ @V\widehat{\nu}VV @VV\nu V \\ \widehat{S} @>\nu>> S. \end{CD}$$

By the definition we have

$$\nu_*c_*^\nu(\mathbb{1}_{\widehat{S}}) = \nu_*\left(c(\widehat{\nu}^*T\widehat{S} - \widehat{\nu}^*T\widehat{S}) \cap [\widehat{S} \times_S \widehat{S}]\right).$$

On the other hand, we have that

$$\nu_*\mathbb{1}_{\widehat{S}} = \mathbb{1}_S + \left(\chi(\nu^{-1}(x_0)) - 1\right) \cdot \mathbb{1}_{x_0}.$$

Hence by the definition we have

$$c_*^{\text{id}_S}(\nu_*\mathbb{1}_{\widehat{S}}) = c_*^{\text{id}_S}(\mathbb{1}_S) + \left(\chi(\nu^{-1}(x_0)) - 1\right)c_*^{\text{id}_S}(\mathbb{1}_{x_0}).$$

Now, to define $c_*^{\text{id}_S}(\mathbb{1}_S)$ we need the following fiber squares:

$$\begin{CD} \widehat{S} \times_S \widehat{S} @>\widehat{\nu}>> \widehat{S} @>\text{id}_{\widehat{S}}>> \widehat{S} \\ @V\widehat{\nu}VV @VV\nu V @VV\nu V \\ \widehat{S} @>\nu>> S @>\text{id}_S>> S, \end{CD}$$

which is nothing but the above diagram (2.2.1). Therefore we get the following:

$$c_*^{\text{id}_S}(\mathbb{1}_S) = \nu_*c_*^\nu(\mathbb{1}_{\widehat{S}}).$$

It is easy to see that $c_*^{\text{id}_S}(\mathbb{1}_{x_0}) = 1$. Therefore, the hypothesis that $\chi(\nu^{-1}(x_0)) \neq 1$ implies that

$$\nu_*c_*^\nu(\mathbb{1}_{\widehat{S}}) \neq c_*^{\text{id}_S}(\nu_*\mathbb{1}_{\widehat{S}}).$$

□

A typical example of such a variety S in Theorem (2.2) is, for example, more than two lines intersecting at one point.

§3. CONVOLUTION

The notion of *convolution* (product) is an important technique ubiquitous in the geometric representation theory. Here we recall the convolution on the Borel-Moore homology theory.

In this paper the homology theory $H_*(X)$ is the Borel-Moore homology group of a locally compact Hausdorff space X , i.e., the ordinary (singular) cohomology group of the pair (\bar{X}, ∞) where $\bar{X} = X \cup \infty$ is the one-point compactification of X .

For any closed subsets X and X' in a smooth manifold M , we have the cup product

$$\cup : H^p(M, M \setminus X) \otimes H^q(M, M \setminus X') \rightarrow H^{p+q}(M, M \setminus (X \cap X')),$$

which implies, by the Alexander duality isomorphism $H^\bullet(M, M \setminus A) \cong H_{\dim M - \bullet}(A)$, the following intersection product:

$$\cdot : H_i(X) \otimes H_j(X') \rightarrow H_{i+j-\dim M}(X \cap X').$$

Let M_1, M_2, M_3 be smooth oriented manifolds. Let $p_{ij} : M_1 \times M_2 \times M_3 \rightarrow M_i \times M_j$ be the canonical projections. Let $Z \subset M_1 \times M_2$ and $Z' \subset M_2 \times M_3$ be closed subsets and assume that the restricted map

$$p_{13} : p_{12}^{-1}(Z) \cap p_{23}^{-1}(Z') \rightarrow M_1 \times M_3$$

is proper. Then its image is denoted by the $Z \circ Z'$, i.e., the composite of the two correspondences Z and Z' (see Fulton's book [F]). With this set-up, the convolution

$$\star : H_i(Z) \otimes H_j(Z') \rightarrow H_{i+j-\dim M}(Z \circ Z')$$

is defined by

$$(3.1) \quad \alpha \star \beta := p_{13*}(p_{12}^* \alpha \cdot p_{23}^* \beta).$$

In particular, when $M_1 = M_2 = M_3 = M$, for any closed subvariety $Z \subset M \times M$, we have $Z \circ Z = Z$ and therefore $H_*(Z)$ is a convolution algebra.

As one can see in the above construction of convolution, as long as the operations of product, pullback and pushforward are available on certain algebraic objects defined on (topological) spaces, one can always define a convolution product. For example, the convolution of constructible functions is the obvious one defined in the same way as (3.1), using the usual product, pullback and pushforward of constructible functions.

For varieties X, Y , we set

$$\widetilde{\mathbb{F}}(X \times Y) := \mathbb{1}_X \times F(Y) = \{\mathbb{1}_X \times \alpha \mid \alpha \in F(Y)\}.$$

Let $\pi : X \times Y \rightarrow X$ be the projection to the first factor. Then we can see that for a nonsingular variety X we have

$$c_*^\pi(\mathbb{1}_X \times \alpha) = [X] \times c_*(\alpha).$$

This observation follows from the multiplicativity of Chern-Schwartz-MacPherson class, $c_*(\gamma \times \delta) = c_*(\gamma) \times c_*(\delta)$ ([K] and also see [KY]). Indeed,

$$\begin{aligned} c_*^\pi(\mathbb{1}_X \times \alpha) &= \pi^*s(TX) \cap c_*(\mathbb{1}_X \times \alpha) \\ &= (s(TX) \times 1) \cap (c_*(\mathbb{1}_X) \times c_*(\alpha)) \\ &= (s(TX) \times 1) \cap ((c(TX) \cap [X]) \times c_*(\alpha)) \\ &= (s(TX) \cap (c(TX) \cap [X])) \times (1 \cap c_*(\alpha)) \\ &= [X] \times c_*(\alpha). \end{aligned}$$

Or, we can simply take this as the definition of the homomorphism $c_*^\pi : \tilde{\mathbb{F}}(X \times Y) \rightarrow H_*(X \times Y)$ with $\pi : X \times Y \rightarrow X$ being the projection. The following theorem is a naïve “constructible function version” of Ginzburg’s theorem [G2, Theorem 6.7]:

Theorem (3.2). *Let X be a nonsingular variety and Y be an arbitrary variety. Then the homomorphism $c_*^{\text{biv}} : \tilde{\mathbb{F}}(X \times Y) \rightarrow H_*(X \times Y)$ defined by $c_*^{\text{biv}}(\mathbb{1}_X \times \alpha) = [X] \times c_*(\alpha)$ is convolutive; i.e., the following diagram commutes:*

$$\begin{array}{ccc} \tilde{\mathbb{F}}(X_1 \times X_2) \otimes \tilde{\mathbb{F}}(X_2 \times X_3) & \xrightarrow{\star} & \tilde{\mathbb{F}}(X_1 \times X_3) \\ c_*^{\text{biv}} \otimes c_*^{\text{biv}} \downarrow & & \downarrow c_*^{\text{biv}} \\ H_*(X_1 \times X_2) \otimes H_*(X_2 \times X_3) & \xrightarrow{\star} & H_*(X_1 \times X_3). \end{array}$$

Proof. By the definition of the convolution of constructible functions it follows that for constructible functions $\mathbb{1}_{X_1} \times \alpha \in \tilde{\mathbb{F}}(X_1 \times X_2)$, $\mathbb{1}_{X_2} \times \beta \in \tilde{\mathbb{F}}(X_2 \times X_3)$, we have

$$(\mathbb{1}_{X_1} \times \alpha) \star (\mathbb{1}_{X_2} \times \beta) = \left(\int_{X_2} \alpha \right) (\mathbb{1}_{X_1} \times \beta),$$

where $\int_{X_2} \alpha$ denotes the Euler-Poincaré characteristic of the constructible function α , $\chi(\alpha)$, in other words, it is nothing but the degree of the 0-dimensional component of the Chern-Schwartz-MacPherson class $c_*(\alpha)$ of the constructible function α . Therefore we get

$$c_*^{\text{biv}}((\mathbb{1}_{X_1} \times \alpha) \star (\mathbb{1}_{X_2} \times \beta)) = \left(\int_{X_2} \alpha \right) ([X_1] \times c_*(\beta)).$$

Let $\pi_{13}^{123} : X_1 \times X_2 \times X_3 \rightarrow X_1 \times X_3$ be the projection and let $\mathcal{D}_X : H^*(X) \rightarrow H_*(X)$ be the Poincaré duality isomorphism. Then, by the definition of convolution of Borel-Moore homology we get

$$\begin{aligned} &c_*^{\text{biv}}(\mathbb{1}_{X_1} \times \alpha) \star c_*^{\text{biv}}(\mathbb{1}_{X_2} \times \beta) \\ &= ([X_1] \times c_*(\alpha)) \star ([X_2] \times c_*(\beta)) \\ &= (\pi_{13}^{123})_* \left(((1 \times \mathcal{D}_{X_2}^{-1}(c_*(\alpha)) \times 1) \cup (1 \times 1 \times \mathcal{D}_{X_3}^{-1}(c_*(\beta)))) \cap [X_1 \times X_2 \times X_3] \right) \\ &= (\pi_{13}^{123})_* \left(\left((1 \times \mathcal{D}_{X_2}^{-1}(c_*(\alpha)) \times 1) \cup \mathcal{D}_{X_3}^{-1}(c_*(\beta)) \right) \cap ([X_1] \times [X_2] \times [X_3]) \right) \\ &= (\pi_{13}^{123})_* ([X_1] \times c_*(\alpha) \times c_*(\beta)) \\ &= \left(\int_{X_2} \alpha \right) ([X_1] \times c_*(\beta)). \end{aligned}$$

Thus the above diagram is commutative. □

Motivated by Theorem (3.2), in [Y2] we proved the following theorem:

Theorem (3.3). *For nonsingular varieties X_1, X_2, X_3 , the homomorphism $c_*^{\text{biv}} : \mathbb{F}(X \times Y \xrightarrow{\pi} X) \rightarrow H_*(X \times Y)$ defined by $c_*^{\text{biv}} := c_*^\pi$ is convolutive; i.e., the following diagram commutes:*

$$\begin{array}{ccc} \mathbb{F}(X_1 \times X_2 \rightarrow X_1) \otimes \mathbb{F}(X_2 \times X_3 \rightarrow X_2) & \xrightarrow{\star} & \mathbb{F}(X_1 \times X_3 \rightarrow X_1) \\ c_*^{\text{biv}} \otimes c_*^{\text{biv}} \downarrow & & \downarrow c_*^{\text{biv}} \\ H_*(X_1 \times X_2) \otimes H_*(X_2 \times X_3) & \xrightarrow{\star} & H_*(X_1 \times X_3). \end{array}$$

Here $\mathbb{F}(X \times Y \rightarrow X)$ is the Fulton-MacPherson bivariant group of constructible functions (see [B] and [FM]).

The key ingredient of the proof of the above theorem is Brasselet's bivariant Chern class theorem [B, III, Théorème] (see [Y2] for the details). For some other properties of the Ginzburg bivariant Chern classes, see [Y1], [Y2], [Y3].

REFERENCES

- [B] J.-P. Brasselet, *Existence des classes de Chern en théorie bivariante*, Astérisque **101-102** (1981), 7–22. MR **85j**:32019
- [CG] N. Chriss and V. Ginzburg, *Representation theory and complex geometry*, Birkhäuser, 1997. MR **98i**:22021
- [F] W. Fulton, *Intersection Theory*, Springer-Verlag, 1984. MR **85k**:14004
- [FM] W. Fulton and R. MacPherson, *Categorical frameworks for the study of singular spaces*, Memoirs of Amer. Math. Soc. **31** (1981). MR **83a**:55015
- [G1] V. Ginzburg, \mathfrak{G} -Modules, *Springer's Representations and Bivariant Chern Classes*, Adv. in Maths. **61** (1986), 1–48. MR **87k**:17014
- [G2] ———, *Geometric methods in the representation theory of Hecke algebras and quantum groups*, in “Representation theories and algebraic geometry (Montreal, PQ, 1997)” (ed. by A. Broer and A. Daigneault), Kluwer Acad. Publ., Dordrecht, 1998, pp. 127–183. MR **99j**:17020
- [K] M. Kwieciński, *Formule du produit pour les classes caractéristiques de Chern-Schwartz-MacPherson et homologie d'intersection*, C. R. Acad. Sci. Paris **314** (1992), 625–628. MR **93b**:55008
- [KY] M. Kwieciński and S. Yokura, *Product formula of the twisted MacPherson class*, Proc. Japan Acad **68** (1992), 167–171. MR **94d**:32052
- [M] R. MacPherson, *Chern classes for singular algebraic varieties*, Ann. of Math. **100** (1974), 423–432. MR **50**:13587
- [N1] H. Nakajima, *Quiver varieties and quantum affine algebras (in Japanese)*, Suugaku **52** (2000), 337–359. CMP 2001:06
- [N2] ———, *Quiver varieties and finite dimensional representations of quantum affine algebras*, J. Amer. Math. Soc. **14** (2001), 145–238. CMP 2001:07
- [Y1] S. Yokura, *On the uniqueness problem of the bivariant Chern classes*, preprint (2001).
- [Y2] ———, *On Ginzburg's bivariant Chern classes*, preprint (2001).
- [Y3] ———, *On Ginzburg's bivariant Chern classes, II*, preprint (2001).

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, FACULTY OF SCIENCE, UNIVERSITY OF KAGOSHIMA, 21-35 KORIMOTO 1-CHOME, KAGOSHIMA 890-0065, JAPAN
E-mail address: yokura@sci.kagoshima-u.ac.jp