

REMARKS ON SMALL SETS OF REALS

TOMEK BARTOSZYNSKI

(Communicated by Alan Dow)

ABSTRACT. We show that the Dual Borel Conjecture implies that $\mathfrak{d} > \mathfrak{N}_1$ and find some topological characterizations of perfectly meager and universally meager sets.

1. INTRODUCTION

For $f, g \in \omega^\omega$ let $f \leq^* g$ mean that $f(n) \leq g(n)$ for all but finitely many n . Let

$$\mathfrak{b} = \min\{|F| : F \subseteq \omega^\omega \ \& \ \forall h \in \omega^\omega \ \exists f \in F \ f \not\leq^* h\},$$

$$\mathfrak{d} = \min\{|F| : F \subseteq \omega^\omega \ \& \ \forall h \in \omega^\omega \ \exists f \in F \ h \leq^* f\}.$$

Let \mathcal{N} be the ideal of measure zero subsets of 2^ω with respect to the standard product measure μ , and let \mathcal{M} be the ideal of meager subsets of 2^ω . Let $+$ be the addition mod 2 on 2^ω . For $A, B \subseteq 2^\omega$ let $A + B = \{a + b : a \in A, b \in B\}$.

Definition 1. Let $X \subseteq 2^\omega$. We say that:

- (1) X has strong measure zero if $X + F \neq 2^\omega$ for all $F \in \mathcal{M}$,
- (2) X is strongly meager if $X + F \neq 2^\omega$ for all $F \in \mathcal{N}$,
- (3) X is meager additive if $X + F \in \mathcal{M}$ for all $F \in \mathcal{M}$,
- (4) X is null additive if $X + F \in \mathcal{N}$ for all $F \in \mathcal{N}$.

Let the Borel Conjecture be the statement that there are no uncountable strong measure zero sets and the Dual Borel Conjecture that there are no uncountable strongly meager sets. Both the Borel Conjecture and the Dual Borel Conjecture are (separately) known to be consistent with the ZFC [7], [4], and moreover the Dual Borel Conjecture is consistent with $\mathfrak{b} = \mathfrak{N}_1$, while Rothberger showed that if $\mathfrak{b} = \mathfrak{N}_1$, then there is an uncountable strong measure zero set. We will strengthen this result by showing that:

Theorem 2. (1) *If $\mathfrak{b} = \mathfrak{N}_1$, then there exists an uncountable meager additive set.*
(2) *If $\mathfrak{d} = \mathfrak{N}_1$, then there exists an uncountable null additive set. In particular, the Dual Borel Conjecture implies that $\mathfrak{d} > \mathfrak{N}_1$.*

Received by the editors July 24, 2001 and, in revised form, September 19, 2001.

2000 *Mathematics Subject Classification.* Primary 03E17.

Key words and phrases. Perfectly meager, universally meager, cardinal invariants.

The author was partially supported by NSF grant DMS 9971282 and the Alexander von Humboldt Foundation.

Proof. The construction presented here is a modification of a construction invented by Todorćević. Part (1) was also proved in [5] using different methods.

Let $F = \{f_\alpha : \alpha < \omega_1\}$ be a family of functions in ω^ω such that

- (1) f_α is strictly increasing for $\alpha < \omega_1$,
- (2) $\forall \alpha < \beta f_\alpha \leq^* f_\beta$.

For a perfect tree $p \subseteq 2^{<\omega}$ let $[p]$ denote the set of its branches. Every perfect subset of 2^ω is a set of branches of a perfect tree.

We will build an ω_1 -tree \mathbf{T} of perfect trees of $2^{<\omega}$. Let \mathbf{T}_α denote the α -th level of \mathbf{T} . We require that

- (1) $\forall \beta > \alpha \forall n \in \omega \forall p \in \mathbf{T}_\alpha \exists q \in \mathbf{T}_\beta (q \subseteq p \ \& \ q \cap 2^n = p \cap 2^n)$,
- (2) $\forall p \in \mathbf{T}_{\alpha+1} \forall^\infty n |p \cap 2^{f_\alpha(n)}| \leq 2^n$,
- (3) $\forall \alpha \mathbf{T}_\alpha$ is countable.

SUCCESSOR STEP. Suppose that \mathbf{T}_α is given. For each $p \in \mathbf{T}_\alpha$ choose $\{q_n : n \in \omega\}$ such that

- (1) $\forall n q_n \subseteq p$,
- (2) $q_n \cap 2^n = p \cap 2^n$,
- (3) $[q_n] \cap [q_m] = \emptyset$ for $n \neq m$,
- (4) $\forall n \forall^\infty k |q_n \cap 2^{f_\alpha(k)}| \leq 2^k$.

Set $\{q_n : n \in \omega\}$ to be the successors of p on level $\mathbf{T}_{\alpha+1}$.

LIMIT STEP. Suppose that γ is a limit ordinal and $\{\mathbf{T}_\alpha : \alpha < \gamma\}$ are already constructed.

For each $p \in \bigcup_{\alpha < \gamma} \mathbf{T}_\alpha$ and $n \in \omega$ we will construct an element $q = q(p, n)$ belonging to the level \mathbf{T}_γ as follows.

Suppose $p = p_0 \in \mathbf{T}_{\alpha_0}$, $n = n_0$ and construct sequences $\langle \alpha_k : k \in \omega \rangle$, $\langle n_k : k \in \omega \rangle$ and $\langle p_k : k \in \omega \rangle$ such that

- (1) $p_k \in \mathbf{T}_{\alpha_k}$,
- (2) $\sup_k \alpha_k = \gamma$, $\lim_k n_k = \infty$,
- (3) $p_{k+1} \subseteq p_k$ for all k ,
- (4) $p_{k+1} \cap 2^{n_k} = p_k \cap 2^{n_k}$,
- (5) $q = \bigcap_k p_k$ is a perfect tree.

The last condition is guaranteed by the careful choice of the sequence $\langle n_k : k \in \omega \rangle$.

Let X be the set obtained by selecting one element out of every tree $p \in \mathbf{T}$.

The following lemma gives the first part of the theorem.

Lemma 3. *If F is an unbounded family in ω^ω , then X is meager additive.*

Proof. Suppose that $H \subseteq 2^\omega$ is a meager set. It is well known (see [2], Theorem 2.2.4) that there exists $x_H \in 2^\omega$ and a strictly increasing function $f_H \in \omega^\omega$ such that

$$H \subseteq \{x \in 2^\omega : \forall^\infty n \exists j \in [f_H(n), f_H(n+1)) x(j) \neq x_H(j)\}.$$

Since we work with translations without loss of generality we can assume that $x_H(k) = 0$ for all k . As F is unbounded there is $\alpha_0 < \omega_1$ such that

$$\exists^\infty n \exists k f_{\alpha_0}(n) < f_H(k) < f_H(k+1) < \dots < f_H(k+2^n) < f_{\alpha_0}(n+1).$$

Fix sequences $\langle u_n, k_n : n \in \omega \rangle$ such that

$$\forall n f_{\alpha_0}(u_n) < f_H(k_n) < f_H(k_n+1) < \dots < f_H(k_n+2^{u_n}) < f_{\alpha_0}(u_n+1).$$

Fix $p \in \mathbf{T}_{\alpha_0+1}$ and let $z_p \in 2^\omega$ be defined as follows: given $n \in \omega$, let $\{s_1, \dots, s_{2^{u_n}}\}$ be an enumeration of $p \cap 2^{f_{\alpha_0}(u_n+1)}$. Define $z_p \upharpoonright [f_H(k_n + i), f_H(k_n + i + 1)) = s_i \upharpoonright [f_H(k_n + i), f_H(k_n + i + 1))$ for $i \leq 2^{u_n}$. This defines z_p on an infinite subset of ω ; extend it arbitrarily to a total function. Let

$$G = \{x \in 2^\omega : \forall^\infty n \ x \upharpoonright [f_{\alpha_0}(u_n), f_{\alpha_0}(u_n + 1)) \neq z_p \upharpoonright [f_{\alpha_0}(u_n), f_{\alpha_0}(u_n + 1))\}.$$

We claim that $[p] + H \subseteq G$. Note that if $x \in [p] + H$, then there exists $y \in [p]$ such that

$$\forall^\infty n \ x \upharpoonright [f_H(n), f_H(n + 1)) \neq y \upharpoonright [f_H(n), f_H(n + 1)).$$

Fix one such y and note that for sufficiently large n there is an i such that

$$\begin{aligned} y \upharpoonright [f_H(k_n + i), f_H(k_n + i + 1)) &= s_i \upharpoonright [f_H(k_n + i), f_H(k_n + i + 1)) \\ &= z_p \upharpoonright [f_H(k_n + i), f_H(k_n + i + 1)), \end{aligned}$$

which implies that $x \upharpoonright [f_H(k_n + i), f_H(k_n + i + 1)) \neq z_p \upharpoonright [f_H(k_n + i), f_H(k_n + i + 1))$. It follows that $x \upharpoonright [f_{\alpha_0}(u_n), f_{\alpha_0}(u_n + 1)) \neq z_p \upharpoonright [f_{\alpha_0}(u_n), f_{\alpha_0}(u_n + 1))$, which means that $x \in G$.

Let X_{α_0} be the collection of points selected from levels $\bigcup_{\alpha < \alpha_0} \mathbf{T}_\alpha$. We have

$$X + G \subseteq (X_{\alpha_0} + G) \cup \bigcup_{p \in \mathbf{T}_{\alpha_0}} [p] + G \in \mathcal{M},$$

which finishes the proof. □

To prove the second part of the theorem, we will use the following lemma:

Lemma 4. *If F is a dominating family, then X is null additive.*

Proof. The following is well known: □

Lemma 5. *Suppose that $H \subseteq 2^\omega$ is a null set. There exists a sequence of clopen sets $\{C_n : n \in \omega\}$ such that that for all n ,*

- (1) $\mu(C_n) < 4^{-n}$,
- (2) $\forall x \in H \exists^\infty n \ x \in C_n$.

Proof. Since H has measure zero, there are open sets $\langle U_n : n \in \omega \rangle$ covering H such that $\mu(U_n) < 4^{-n-2}$, for $n \in \omega$. Write each set U_n as a disjoint union of open basic intervals, $U_n = \bigcup_{m=1}^\infty [s_m^n]$ for $n \in \omega$, and order these sequences lexicographically in a single sequence $\{t_n : n \in \omega\}$. Put $k_0 = 0$ and for $n > 0$ let $k_n = \min\{k : \sum_{j \geq k} \mu([t_j]) < 4^{-n-1}\}$. Let

$$C_n = \bigcup_{k \in [k_n, k_{n+1})} [t_k].$$

Clearly $\mu(C_n) < 4^{-n}$ and, since all basic sets have been accounted for,

$$\forall x \in H \exists^\infty n \ x \in C_n.$$

□

Let $H \subseteq 2^\omega$ be a measure zero set and $\{C_n : n \in \omega\}$ the sequence given by the lemma. Since each clopen set is a union of finitely many basic sets, we can find a function $f_H \in \omega^\omega$ such that for every n , $C_n \subseteq 2^{f_H(n)}$. Let α_0 and n_0 be such that

$$\forall n > n_0 \ f_H(n) < f_{\alpha_0}(n).$$

By modifying finitely many C_n 's, we can assume that $n_0 = 0$. Suppose that $p \in \mathbf{T}_{\alpha_0+1}$. Note that

$$[p] + C_n \subseteq \left[p \cap 2^{f_H(n)} \right] + C_n = D_n.$$

Since $|p \cap 2^{f_H(n)}| \leq |p \cap 2^{f_{\alpha_0}(n)}|$, it follows that

$$\mu([p] + C_n) = \mu(D_n) \leq 2^n \cdot 4^{-n} \leq 2^{-n}.$$

Therefore, $[p] + H \subseteq G = \{x \in 2^\omega : \exists^\infty n \ x \in D_n\}$, and $\mu(G) = 0$. The rest of the proof is identical to the proof of the first part. \square

2. PERFECTLY MEAGER AND UNIVERSALLY MEAGER SETS

A set $X \subseteq 2^\omega$ is perfectly meager ($X \in \mathbf{PM}$) if $P \cap X$ is meager in P for every perfect set P .

A set $X \subseteq 2^\omega$ is universally meager ($X \in \mathbf{UM}$, [10]) if for every Borel isomorphism $F : 2^\omega \rightarrow 2^\omega$, $F''(X) \in \mathcal{M}$.

A set $X \subseteq 2^\omega$ is universally null ($X \in \mathbf{UN}$) if for every Borel isomorphism $F : 2^\omega \rightarrow 2^\omega$, $F''(X) \in \mathcal{N}$.

In [10] it is shown that the universally meager sets are a category analog of universally null sets. Yet, for quite a while perfectly meager sets were viewed in this role. Theorems 6 and 7 explain this phenomenon, which has to do with the fact that the families \mathbf{PM} and \mathbf{UM} are not *very* different. It is easy to see that $\mathbf{UM} \subseteq \mathbf{PM}$. The Continuum Hypothesis or Martin's Axiom imply that $\mathbf{UM} \neq \mathbf{PM}$, but it is also consistent that $\mathbf{UM} = \mathbf{PM}$, [1].

The first theorem gives a simple proof of a characterization of perfectly meager sets found in [3].

Theorem 6. *The following are equivalent:*

- (1) $X \in \mathbf{PM}$,
- (2) for every countable dense-in-itself set A there exists a set $B \subseteq A$ such that $\text{cl}(A) = \text{cl}(B)$ and B is a G_δ -set relative to $X \cup A$,
- (3) for every countable set A there exists a set $B \subseteq A$ such that $\text{cl}(A) = \text{cl}(B)$ and B is a G_δ -set relative to $X \cup A$,
- (4) for every perfect set P there exists an F_σ -set F such that $X \subseteq F$ and F is meager in P .

Proof. (2) \rightarrow (1). Let P be a perfect set and $Q \subseteq P$ a countable dense set. By (2) without loss of generality we can assume that Q is a G_δ relative to $X \cup Q$. In other words $Q = (X \cup Q) \cap \bigcap G_n$, where G_n are open sets. Clearly G_n 's are dense in P . It follows that $X \cap P \subseteq Q \cup P \setminus \bigcap G_n$.

(1) \rightarrow (4). Since $X \in \mathbf{PM}$, there exists an F_σ -set F_1 such that $X \subseteq F_1$, and there exists an F_σ -set F_2 such that $F_2 \cap P$ is meager in P and $X \cap P \subseteq F_2$. Now the set $F = (F_1 \setminus P) \cup (F_2 \cap P)$ is the set we are looking for.

(4) \rightarrow (3). Suppose that A is countable. By the Cantor-Bendixson Theorem, [6], $\text{cl}(A) = P \dot{\cup} C$, where P is perfect, C is countable and open relative to P (that is, C is contained in an open set disjoint with P). If $P = \emptyset$, then C is countable and closed and $A = C \setminus (C \setminus A)$ is a G_δ -set in 2^ω . Thus assume that $P \neq \emptyset$ and let $\langle F'_n : n \in \omega \rangle$ be closed nowhere dense sets such that $X \subseteq \bigcup_n F'_n$ and $F'_n \cap P$ is closed nowhere dense in P for each n . Let $\{F_n : n \in \omega\}$ be closed sets such that $\bigcup_n F_n = (\bigcup_{n \in \omega} F'_n) \setminus C$. Let $A' = A \setminus C$ and consider sets $A_n = A' \setminus F_n$. Since the

family $\langle A_n : n \in \omega \rangle$ has the finite intersection property, we can find a set $B' \subseteq A'$ such that

- (1) $B' \setminus A_n$ is finite for all n ,
- (2) B' is dense in P .

Let $F_n^* = F_n \setminus B'$. Note that each set F_n^* is F_σ since it differs from F_n by a finite set. Put $B = (X \cup A) \cap \bigcap_n (2^\omega \setminus F_n^*)$. It follows that $B' \cup (A \cap C) \subseteq B \subseteq A$, which finishes the proof.

(3) \rightarrow (2) is obvious. □

Theorem 7. *The following are equivalent:*

- (1) $X \in \mathbf{UM}$.
- (2) For every sequence of countable dense-in-itself sets $\{A_n : n \in \omega\}$ there exists a sequence $B_n \subseteq A_n$ such that $\text{cl}(A_n) = \text{cl}(B_n)$ and $\bigcup_n B_n$ is a G_δ -set relative to $X \cup \bigcup_n A_n$.
- (3) For every sequence of perfect sets $\{P_n : n \in \omega\}$, there exists an F_σ -set F such that $X \subseteq F$ and F is meager in P_n for every n .

Proof. (2) \rightarrow (1). The following argument is a small modification of a proof from [9]. Suppose that $X \notin \mathbf{UM}$. Let $G : X \rightarrow Y'$ be a Borel isomorphism onto a non-meager set. By Kuratowski's Theorem, G^{-1} is continuous on a dense G_δ -set. It follows that there exists a continuous one-to-one function $F : Y \rightarrow X$, where $Y \notin \mathcal{M}$. Let $\{U_n : n \in \omega\}$ be enumeration of clopen subsets of 2^ω such that $U_n \cap Y$ is uncountable. For each n , choose a countable dense-in-itself set $A_n \subseteq F''(Y \cap U_n)$. We will show that every G_δ -set which is disjoint from $X \setminus \bigcup_n A_n$ is also disjoint from one of the A_n 's. Let $F = \bigcup_n F_n$ be an F_σ -set containing $X \setminus \bigcup_n A_n$. For every n let H_n be a closed set such that $H_n \cap Y = G^{-1}(F_n)$. If for every n , $Y_n = \text{interior}(H_n) \cap Y$ is countable, then $Y \subseteq \bigcup_n Y_n \cup \bigcup_n (H_n \setminus \text{interior}(H_n)) \in \mathcal{M}$, which is impossible. Thus there exists $m, n \in \omega$ such that $U_m \subseteq H_n$ and therefore $A_m \subseteq F''(U_m \cap Y) \subseteq F_n$.

(1) \rightarrow (3). Let \mathbb{C} denote the Cohen algebra. The following is a (small) fragment of Theorem 2.1 from [10].

Theorem 8. *For a subset X of a perfect Polish space \mathbf{X} , the following are equivalent:*

- (1) $X \in \mathbf{UM}$.
- (2) For every σ -ideal \mathcal{J} in $\text{Borel}(\mathbf{X})$ such that $\text{Borel}(\mathbf{X})/\mathcal{J} \cong \mathbb{C}$ there is a Borel set $B \in \mathcal{J}$ such that $X \subseteq B$.

Proof. The implication (1) \rightarrow (2) that is of interest to us is a consequence of Sikorski's Theorem ([6], 15.10): if \mathcal{J} is a σ -ideal in $\text{Borel}(\mathbf{X})$ such that $\text{Borel}(\mathbf{X})/\mathcal{J} \cong \mathbb{C}$, then there is a Borel automorphism $F : \mathbf{X} \rightarrow \mathbf{X}$ such that

$$\forall X \in \text{Borel}(\mathbf{X}) \ X \in \mathcal{M} \iff F''(X) \in \mathcal{J}. \quad \square$$

Suppose that $\{P_n : n \in \omega\}$ are given. Consider the ideal

$$\mathcal{J} = \{A : A \text{ is Borel and } \forall n \ A \cap P_n \text{ is meager in } P_n\}.$$

It is easy to see that $\text{Borel}(\mathbf{X})/\mathcal{J} \cong \mathbb{C}$ (as $\text{Borel}(\mathbf{X})/\mathcal{J}$ is atomless and has a countable dense subset). Therefore there exists a set $F \in \mathcal{J}$ such that $X \subseteq F$.

(3) \rightarrow (2). Let $P_n = \text{cl}(A_n)$. Let $F = \bigcup_n F_n$ be an F_σ -set containing X such that F is meager in each P_n . As in the proof of Theorem 6, build by induction a

set $B' \subseteq \bigcup_n A_n$ such that for $n \in \omega$,

- (1) $\text{cl}(B' \cap A_n) = \text{cl}(A_n)$,
- (2) $B' \setminus F_n$ is finite.

As before $F_n^* = F_n \setminus B'$ is an F_σ -set and let $B = (X \cup \bigcup_n A_n) \cap \bigcap_n (2^\omega \setminus F_n^*)$. The sets $B_n = B \cap A_n$ for $n \in \omega$ are as required. \square

The assumption in Theorem 7(2) that the sets A_n are dense-in-itself is necessary since we have the following:

Theorem 9. *There exists a set $X \in \mathbf{UM}$ and a family $\{A_n : n \in \omega\}$ of countable subsets of X such that if $B_n \subseteq A_n$ is such that $\text{cl}(B_n) = \text{cl}(A_n)$ for each n , then $\bigcup_n B_n$ is not a G_δ relative to X .*

Proof. Recall that a set of reals is called a λ -set if all of its countable subsets are relative G_δ -sets. Theorem 7 implies readily that all λ -sets and unions of countable sets with λ -sets are universally meager. Rothberger showed (5.6 of [8]) that there exists a set $Y \subseteq 2^\omega$ and a countable set $C \subseteq 2^\omega$ such that

- (1) Y is a λ -set,
- (2) $X = Y \cup C$ is not a λ -set, that is C is not a G_δ -set relative to Y .

Let Y and C be as above. Write $C = \bigcup_n A_n$, where each A_n is infinite and discrete, that is $x \notin \text{cl}(A_n \setminus \{x\})$ for $x \in A_n$. It follows that if $B_n \subseteq A_n$ and $\text{cl}(B_n) = \text{cl}(A_n)$, then $A_n = B_n$, which finishes the proof. \square

REFERENCES

- [1] Tomek Bartoszynski. On perfectly meager sets. available from front.math.ucdavis.edu.
- [2] Tomek Bartoszyński and Haim Judah. *Set Theory: on the structure of the real line*. A.K. Peters, 1995. MR **96k**:03002
- [3] Harold R. Bennett, Masami Hosobuchi, and David J. Lutzer. A note on perfect generalized ordered spaces. *Rocky Mountain J. Math.*, 29(4):1195–1207, 1999. MR **2000k**:54027
- [4] Timothy J. Carlson. Strong measure zero and strongly meager sets. *Proceedings of the American Mathematical Society*, 118(2):577–586, 1993. MR **94b**:03086
- [5] Fred Galvin and Arnold W. Miller. γ -sets and other singular sets of real numbers. *Topology and its Applications*, 17(2):145–155, 1984. MR **85f**:54011
- [6] Alexander Kechris. *Classical Descriptive Set Theory*, volume 156 of *Graduate Texts in Mathematics*. Springer Verlag, 1995. MR **96e**:03057
- [7] Richard Laver. On the consistency of Borel's conjecture. *Acta-Math.*, 137(3-4):151–169, 1976. MR **54**:10019
- [8] Arnold W. Miller. Special subsets of the real line. In K. Kunen and J. E. Vaughan, editors, *Handbook of Set Theoretic Topology*, pages 201–235. North-Holland, Amsterdam, 1984. MR **86i**:54037
- [9] Andrzej Nowik and Tomasz Weiss. Not every Q -set is perfectly meager in the transitive sense. *Proceedings of the American Mathematical Society*, 128(10):3017–3024, 2000. MR **2000m**:03116
- [10] Piotr Zakrzewski. Universally meager sets. *Proceedings of the American Mathematical Society*, 129(6):1793–1798, 2001. MR **2001m**:03097

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, BOISE STATE UNIVERSITY, BOISE, IDAHO 83725

E-mail address: tomek@math.boisestate.edu

URL: <http://math.boisestate.edu/~tomek>