

AUTOMORPHISMS OF THE ENDOMORPHISM SEMIGROUP OF A FREE MONOID OR A FREE SEMIGROUP

G. MASHEVITZKY AND BORIS M. SCHEIN

(Communicated by Stephen D. Smith)

ABSTRACT. We determine all isomorphisms between the endomorphism semigroups of free monoids or free semigroups and prove that automorphisms of the endomorphism semigroup of a free monoid or a free semigroup are inner or “mirror inner”. In particular, we answer a question of B. I. Plotkin.

INTRODUCTION

One of the venerable algebraic problems, the first instance of which was considered by É. Galois, is (using the words of S. Ulam [7]) “determination of a mathematical structure from a given set of endomorphisms”. Let $\mathbf{End}(A)$ and $\mathbf{Aut}(A)$ denote the endomorphism monoid and the automorphism group of an algebraic system A , respectively. What can be said about systems A and B if $\mathbf{End}(A)$ is isomorphic to $\mathbf{End}(B)$? This problem has been considered by numerous authors.

We consider this problem for $\mathbf{End}(X^*)$ and $\mathbf{End}(X^+)$, where X^* and X^+ denote, respectively, the free monoid and the free semigroup generated by a set X . This particular problem about $\mathbf{End}(A)$, for A a free algebra in a certain variety, was raised by B. I. Plotkin [2] in his lectures on universal algebraic geometry. An analogous problem for $\mathbf{End}(F)$ with F a free group was solved by Formanek [1]. Other examples are given in a remark at the end of this paper.

Recall that the elements of X^* are words over X , including the empty word 1. The elements of X^+ are nonempty words. If $u = x_{i_1} \dots x_{i_k} \in X^*$, then \bar{u} denotes the “opposite” word $x_{i_k} \dots x_{i_1}$. In particular, $\bar{1} = 1$. Every bijection $f : X \rightarrow Y$ induces an isomorphism $\iota_f : X^* \rightarrow Y^*$ and an anti-isomorphism $\bar{\iota}_f : X^* \rightarrow Y^*$ defined as follows: $\iota_f(u) = f(x_{i_1}) \dots f(x_{i_k})$ and $\bar{\iota}_f(u) = \overline{\iota_f(u)} = f(x_{i_k}) \dots f(x_{i_1})$. Analogous facts are true for X^+ and Y^+ .

Let $\iota : S \rightarrow T$ be an isomorphism or an anti-isomorphism of a semigroup S onto a semigroup T . Define the mapping $\iota \square \iota : \mathbf{End}(S) \rightarrow \mathbf{End}(T)$ by $\iota \square \iota(\varphi) = \iota \circ \varphi \circ \iota^{-1}$ for all $\varphi \in \mathbf{End}(S)$. Thus, if $\iota(s_1) = t_1$, $\iota(s_2) = t_2$, and $\varphi(s_1) = s_2$ for some $s_1, s_2 \in S$ and $t_1, t_2 \in T$, then $\iota \square \iota(\varphi)(t_1) = t_2$. It is easy to see that $\iota \square \iota$ is an isomorphism of $\mathbf{End}(S)$ onto $\mathbf{End}(T)$. We call it the isomorphism *induced* by ι .

Let $|X|$ denote the cardinality of X and \mathbb{P} the set of prime numbers. A *permutation* of a finite or an infinite set is a bijection of that set onto itself.

Received by the editors December 5, 2001.

1991 *Mathematics Subject Classification.* Primary 20M20, 20M05, 08A35.

Key words and phrases. Free semigroup, free monoid, endomorphism, automorphism.

©2003 American Mathematical Society

Theorem 1. Let $\mathbf{End}(X^*)$ and $\mathbf{End}(Y^*)$ be isomorphic.

If $|X| = 1$, then $|Y| = 1$ and the isomorphisms of $\mathbf{End}(X^*)$ onto $\mathbf{End}(Y^*)$ are in a natural one-to-one correspondence with permutations of \mathbb{P} (explained in the proof).

If $|X| \geq 1$, then every isomorphism $\alpha : \mathbf{End}(X^*) \rightarrow \mathbf{End}(Y^*)$ is induced either by the isomorphism ι_f or by the anti-isomorphism $\bar{\iota}_f$ of X^* onto Y^* for a uniquely determined bijection $f : X \rightarrow Y$. In other words, either $\alpha = \iota_f \square \iota_f$ or $\alpha = \bar{\iota}_f \square \bar{\iota}_f$.

The same results hold for every isomorphism $\alpha : \mathbf{End}(X^+) \rightarrow \mathbf{End}(Y^+)$.

Let $\mathbf{Aut}(X)$ and C_2 denote the symmetric group on X and a 2-element group, respectively. Also, μ is the so-called mirror automorphism (see Definition 1.4(iii)).

Theorem 2. The groups $\mathbf{Aut}(\mathbf{End}(X^+))$ and $\mathbf{Aut}(\mathbf{End}(X^*))$ are isomorphic.

If $|X| > 1$, every automorphism of $\mathbf{End}(X^*)$ and of $\mathbf{End}(X^+)$ is either inner or a product of an inner automorphism and the mirror automorphism μ . In this case $\mathbf{Aut}(\mathbf{End}(X^*))$ is isomorphic to the direct product $\mathbf{Aut}(X) \times C_2$.

If $|X| = 1$, $\mathbf{Aut}(\mathbf{End}(X^*))$ is isomorphic to the symmetric group on a countably infinite set.

In Theorem 2 an automorphism is *inner* if it is of the form ι_f , where f is a permutation of X .

1. NOTATIONS AND PRELIMINARIES

We give the proofs in the case of the free monoid X^* . The proofs in the case of the free semigroup X^+ are almost *verbatim* the same, so we give only a few remarks in the case of free semigroups. Each endomorphism φ of X^* and of X^+ is uniquely determined by a mapping $X \rightarrow X^*$ or, respectively, $X \rightarrow X^+$. To define φ , it suffices to define $\varphi(x)$ for all $x \in X$. The mapping $\varphi \mapsto \varphi^*$ such that $\varphi^*(x) = \varphi(x)$ and $\varphi^*(1) = 1$ defines an injective homomorphism of $\mathbf{End}(X^+)$ into $\mathbf{End}(X^*)$. For simplicity we identify $\mathbf{End}(X^+)$ with a subsemigroup of $\mathbf{End}(X^*)$.

Clearly, φ is an automorphism precisely when its restriction to X is a permutation of X . Thus the automorphism groups $\mathbf{Aut}(X^*)$ and $\mathbf{Aut}(X^+)$ of X^* and X^+ are isomorphic to the symmetric group $\mathbf{Aut}(X)$ of all permutations of X .

Definition 1.1. (i) Let $u = x_{i_1} \dots x_{i_k} \in X^*$. Denote the length k of u by $|u|$. The empty word 1 has length 0.

(ii) Let $\mathbf{c}(u)$ be the set of all letters of u .

(iii) An endomorphism $\varphi \in \mathbf{End}(X^*)$ is *linear* if $\varphi(x) \in X \cup \{1\}$ for every $x \in X$. In the case of $\mathbf{End}(X^+)$, φ is linear when $\varphi(x) \in X$ for all $x \in X$.

(iv) If $u \in X^*$ is a fixed word, let c_u be the endomorphism of X^+ such that $c_u(x) = u$ for all $x \in X$. We call c_u a *constant* endomorphism. Observe that the range of c_u does not consist of a single word, unless $u = 1$. Clearly, c_1 is the zero element of $\mathbf{End}(X^*)$. We denote it by 0 , that is, $c_1 = 0$. If $v \in X^*$, then $c_u(v) = u^{|v|}$.

(v) An endomorphism $\varphi \in \mathbf{End}(X^*)$ is called *full* if $\varphi\psi = 0 \Rightarrow \psi = 0$ for all $\psi \in \mathbf{End}(X^*)$.

Lemma 1.2. (i) φ is a constant endomorphism of $\mathbf{End}(X^*)$ if and only if $\varphi\alpha = \varphi$ for all $\alpha \in \mathbf{Aut}(X^*)$. The same holds for $\mathbf{End}(X^+)$;

(ii) $\varphi c_u = c_{\varphi(u)}$ for all $\varphi \in \mathbf{End}(X^*)$;

- (iii) $\varphi \in \mathbf{End}(X^*)$ is a constant idempotent if and only if either $\varphi = c_x$ for some $x \in X$ or $\varphi = 0$;
- (iv) $\varphi \in \mathbf{End}(X^*)$ is full if and only if $\varphi(x) \neq 1$ for all $x \in X$.

Proof. (i) If $x \in X$, then $\alpha(x) = y \in X$. Varying α , we obtain $c_u\alpha(x) = c_u(y) = u$ for all $y \in X$. Thus $c_u\alpha = c_u$. Conversely, let $\varphi\alpha = \varphi$ for all automorphisms α . If $\alpha(x) = y$ for some $x, y \in X$, then $\varphi(x) = \varphi\alpha(x) = \varphi(y)$. Varying α , we obtain $\varphi(x) = \varphi(y)$ for all $x, y \in X$. It follows that $\varphi = c_u$, where $u = \varphi(x)$ for any $x \in X$.

(ii) $\varphi c_u(x) = \varphi(u)$ for every $x \in X$. Thus $\varphi c_u = c_{\varphi(u)}$.

(iii) If φ is a constant idempotent, that is, $\varphi = c_u$ for some $u \in X^*$ and $c_u c_u = c_u$, then, by (ii), $c_u = c_u c_u = c_{c_u(u)} = c_{u^{|u|}}$. Thus $u^{|u|} = u$, that is, $|u| \leq 1$. The converse is obvious.

(iv) If $\varphi(x) = 1$ for some $x \in X$, then $c_x \neq 0$ but $\varphi c_x = 0$. Thus φ is not full.

Conversely, let $\varphi(x) \neq 1$ for all $x \in X$. Thus $\varphi(u) \neq 1$ for all $u \in X^+$. If $\psi \neq 0$, then $\psi(x) = u \neq 1$ for some $x \in X$, and hence $\psi c_x = c_{\psi(x)} = c_u \neq 0$. Thus $\varphi\psi c_x = \varphi c_u = c_{\varphi(u)} \neq 0$, so that $\varphi\psi \neq 0$. It follows that φ is full. \square

Lemma 1.3. *The mapping $\mu(\varphi) = \bar{\varphi}$ is an automorphism of $\mathbf{End}(X^*)$.*

Proof. Obviously, $f : u \rightarrow \bar{u}$ is an antiautomorphism of X^* and $f^{-1} = f$. Thus $f^{-1}\varphi f = \bar{\varphi}$, and Lemma 1.3 follows from this equality. \square

Definition 1.4. (i) Let $\alpha : \mathbf{End}(X^*) \rightarrow \mathbf{End}(Y^*)$ be an isomorphism. By parts (i) and (iii) of Lemma 1.2, for every $x \in X$ there exists $y \in Y$ such that $\alpha(c_x) = c_y$. Define a bijection $f : X \rightarrow Y$ by $f(x) = y$. We say that f is *induced* by α . We make an analogous definition for $\mathbf{End}(X^+) \rightarrow \mathbf{End}(Y^+)$.

(ii) An automorphism α of $\mathbf{End}(X^*)$ or of $\mathbf{End}(X^+)$ is *stable* if it induces the identity permutation of X , that is, $\alpha(c_x) = c_x$ for all $x \in X$.

(iii) The mapping μ of Lemma 1.3 is the *mirror automorphism* of $\mathbf{End}(X^*)$.

2. THE CASE $|X| = 1$

If $|X| > 1$, let x and y be distinct elements of X . Then $c_{xy}(c_x(x)) = c_{xy}(x) = xy \neq xx = c_x(xy) = c_x c_{xy}(x)$, and hence $c_{xy}c_x \neq c_x c_{xy}$. Thus $\mathbf{End}(X^*)$ and $\mathbf{End}(X^+)$ are not commutative.

If $X = \{x\}$, a singleton, then $X^+ = \{x, x^2, x^3, \dots\}$ consists of all powers of x and is isomorphic to the additive semigroup $(\mathbb{N}, +)$ of positive integers. Every element of $\mathbf{End}(X^+)$ corresponds to $\varphi_k = \begin{pmatrix} x \\ x^k \end{pmatrix}$ for some $k \in \mathbb{N}$, and $\mathbf{End}(X^+)$ is isomorphic to the multiplicative semigroup (\mathbb{N}, \cdot) of positive integers. (\mathbb{N}, \cdot) is a free commutative monoid with the countably infinite set \mathbb{P} of free generators that are prime numbers. Therefore, $\mathbf{End}(X^+)$ is a free commutative semigroup and $\{\varphi_k\}_{k \in \mathbb{P}}$ is its set of free generators. If $\alpha : \mathbf{End}(X^+) \rightarrow \mathbf{End}(Y^+)$ is an isomorphism, then $\mathbf{End}(Y^+)$ is commutative, and hence $|Y| = 1$. Thus α is uniquely determined by a bijection between the free generators of $\mathbf{End}(X^+)$ and $\mathbf{End}(Y^+)$. These bijections (and hence the isomorphisms) are in a one-to-one correspondence with permutations of \mathbb{P} . The elements of $\mathbf{Aut}(\mathbf{End}(X^+))$ correspond to permutations of generators, and thus $\mathbf{Aut}(\mathbf{End}(X^+))$ is an infinite group isomorphic to the symmetric group $\mathbf{Aut}(\mathbb{P})$ of all permutations of \mathbb{P} , and also isomorphic to $\mathbf{Aut}(\mathbb{P}^+)$ because \mathbb{P} is countably infinite.

Similarly, X^* is isomorphic to the additive semigroup $(\mathbb{N}_0, +)$ of nonnegative integers, $\mathbf{End}(X^*)$ is isomorphic to the multiplicative semigroup (\mathbb{N}_0, \cdot) of nonnegative integers, and hence $\mathbf{Aut}(\mathbf{End}(X^*))$ is isomorphic to $\mathbf{Aut}(\mathbf{End}(X^+))$.

3. STABLE AUTOMORPHISMS OF $\mathbf{End}(X^*)$ AND $\mathbf{End}(X^+)$

Lemma 3.1. (i) If α is a stable automorphism of $\mathbf{End}(X^*)$, $\varphi \in \mathbf{End}(X^*)$, and $x \in X$, then $\alpha(c_{\varphi(x)}) = c_{\alpha(\varphi)(x)}$.

(ii) If φ is linear, then $\alpha(\varphi) = \varphi$.

Proof. (i) By Lemma 1.2(ii), $\alpha(c_{\varphi(x)}) = \alpha(\varphi c_x) = \alpha(\varphi)\alpha(c_x) = \alpha(\varphi)c_x = c_{\alpha(\varphi)(x)}$.

(ii) If φ is linear, then $\varphi(x) \in X \cup \{1\}$, and hence $c_{\alpha(\varphi)(x)} = \alpha(c_{\varphi(x)}) = c_{\varphi(x)}$. Therefore, $\alpha(\varphi)(x) = \varphi(x)$ for every $x \in X$, and so $\alpha(\varphi) = \varphi$. \square

Lemma 3.2. If $\alpha(c_u) = c_v$, where α is a stable automorphism of $\mathbf{End}(X^*)$ and $u, v \in X^*$, then $\mathbf{c}(u) = \mathbf{c}(v)$.

Proof. If $z \in \mathbf{c}(u) \setminus \mathbf{c}(v)$, choose $x \in X$, $\varphi \in \mathbf{End}(X^*)$, and $g \in \mathbf{End}(X^+)$ such that $x \neq z$, $\varphi(x) = u$, $g(z) = x$, and $g(y) = y$ for all $y \neq z$, $y \in X$. Then g is linear, $g(v) = v$, $\alpha(g) = g$ by Lemma 3.1(ii), and $\alpha(c_u) = c_v = c_{g(v)} = gc_v = \alpha(g)\alpha(c_u) = \alpha(gc_u) = \alpha(c_{g(u)})$. By injectivity of α , $c_u = c_{g(u)}$, so that $u = g(u)$, which is not true. Thus $\mathbf{c}(u) \setminus \mathbf{c}(v) = \emptyset$. Similarly, $\mathbf{c}(v) \setminus \mathbf{c}(u) = \emptyset$. Therefore, $\mathbf{c}(u) = \mathbf{c}(v)$. \square

Lemma 3.3. If α is a stable automorphism of $\mathbf{End}(X^*)$, then $|\varphi(x)| = |\alpha(\varphi)(x)|$ for all $\varphi \in \mathbf{End}(X^*)$ and $x \in X$.

Proof. Suppose that $|\varphi_1(x)| = |\varphi_2(x)| = m$, $|\alpha(\varphi_1)(x)| = k$, and $|\alpha(\varphi_2)(x)| = l$. Then $c_x \varphi_1 c_x = c_x \varphi_2 c_x = c_{x^m}$. Also, $\alpha(c_x) = c_x$. Therefore, $\alpha(c_{x^m}) = \alpha(c_x \varphi_1 c_x) = c_x \alpha(\varphi_1) c_x = c_{x^k}$ and $\alpha(c_{x^m}) = \alpha(c_x \varphi_2 c_x) = c_x \alpha(\varphi_2) c_x = c_{x^l}$. Thus $k = l$.

If Y is a finite n -element subset of X and m a nonnegative integer, define $\mathbf{End}_Y^m(x) = \{\varphi \in \mathbf{End}(X^*) \mid |\varphi(x)| = m \text{ and } \mathbf{c}(\varphi(x)) = Y\}$. By the previous paragraph and Lemma 3.2, for every m there exists k such that $\alpha(\mathbf{End}_Y^m(x)) \subseteq \mathbf{End}_Y^k(x)$. There are n^k words of length k and n^m words of length m over Y . Since α is injective, then $k \geq m$. Since α is surjective, $k = m$. Thus $\alpha(\mathbf{End}_Y^m(x)) = \mathbf{End}_Y^m(x)$, and hence $|\varphi(x)| = |\alpha(\varphi)(x)|$ for all $\varphi \in \mathbf{End}(X^*)$ and $x \in X$. \square

Corollary 3.4. Choose two distinct elements $x_1, x_2 \in X$. If α is a stable automorphism of $\mathbf{End}(X^*)$, then $\alpha(c_{x_1 x_2})$ is either $c_{x_1 x_2}$ or $c_{x_2 x_1}$.

Proof. By Lemma 1.2(i), $\alpha(c_{x_1 x_2}) = c_u$ for some $u \in X^*$. By Lemma 3.2, $\mathbf{c}(u) = \{x_1, x_2\}$. By Lemma 3.3, $|u| = 2$. Thus u is either $x_1 x_2$ or $x_2 x_1$. \square

Lemma 3.5. If α is a stable automorphism of $\mathbf{End}(X^*)$ and $\alpha(c_{x_1 x_2}) = c_{x_1 x_2}$ for some $x_1, x_2 \in X$, $x_1 \neq x_2$, then α is the identity automorphism of $\mathbf{End}(X^*)$.

Proof. First we prove that $\alpha(c_u) = c_u$ for all $u \in X^*$ by induction on $|u|$. The base of induction is obvious for $|u| = 0$ and follows from the stability of α for $|u| = 1$. Let the claim hold for $|u| < k$, and let $|u| = k$, where $u = u_1 y$. If $\varphi(x_1) = u_1$, $\varphi(x_2) = y$, and $\varphi(x) = x$ for other $x \in X$, then $\varphi c_{x_1 x_2} = c_u$. By the induction hypothesis, $\alpha(c_{u_1}) = c_{u_1}$. Also, $\alpha(c_x) = c_x$ for every $x \in X$. By Lemma 3.1(i), $\alpha(\varphi)(x) = \varphi(x)$ for all $x \in X$. Thus $\alpha(\varphi) = \varphi$, and hence $\alpha(c_u) = \alpha(\varphi c_{x_1 x_2}) = \varphi c_{x_1 x_2} = c_u$.

By Lemma 3.1(i), $c_{\varphi(x)} = \alpha(c_{\varphi(x)}) = c_{\alpha(\varphi)(x)}$, and so $\varphi(x) = \alpha(\varphi(x))$ for all $x \in X$. Thus $\varphi = \alpha(\varphi)$ for all $\varphi \in \mathbf{End}(X^*)$. \square

Lemma 3.6. *If α is a stable automorphism of $\mathbf{End}(X^*)$ and $\alpha(c_{x_1x_2}) = c_{x_2x_1}$ for some $x_1, x_2 \in X$, $x_1 \neq x_2$, then $\alpha = \mu$.*

Proof. First we prove that $\alpha(c_u) = c_{\bar{u}}$ for all $u \in X^*$ by induction on $|u|$. The base of induction is obvious for $|u| = 0$ and follows from the stability of α for $|u| = 1$. Let the claim hold for $|u| < k$, and let $|u| = k$, where $u = u_1y$. If $\varphi(x_1) = u_1, \varphi(x_2) = y$, and $\varphi(x) = x$ for other $x \in X$, then $\varphi c_{x_1x_2} = c_u$. By the induction hypothesis, $\alpha(c_{u_1}) = c_{\bar{u}_1}$. Also, $\alpha(c_x) = c_x$ for every $x \in X$. By Lemma 3.1(i), $\alpha(\varphi)(x) = \overline{\varphi(x)}$ for all $x \in X$. Thus $\alpha(\varphi) = \overline{\varphi}$, and hence $\alpha(c_u) = \alpha(\varphi c_{x_1x_2}) = \overline{\varphi} c_{x_2x_1} = c_{y\bar{u}_1} = c_{\bar{u}}$.

By Lemma 3.1(i), $\alpha(\varphi) = \overline{\varphi}$ for all $\varphi \in \mathbf{End}(X^*)$. Thus $\alpha = \mu$. □

We skip the proof of the following proposition (it is obtained as an obvious simplification of our proof of Theorem 2).

Proposition 3.7. *Let $F(X)$ be the free commutative monoid with a set X , $|X| > 1$, of free generators. Every automorphism of $\mathbf{End}(F(X))$ is inner.*

Remark 3.8. Similarly, automorphisms of a free commutative semigroup are inner.

4. GROUP OF AUTOMORPHISMS OF $\mathbf{End}(X^*)$

Proof of Theorem 1. For $|X| = 1$ Theorem 1 was proved in Section 2. Suppose that $|X| > 1$, $\alpha : \mathbf{End}(X^*) \rightarrow \mathbf{End}(Y^*)$ is an isomorphism, and $f : X \rightarrow Y$ is a bijection induced by α , that is, $\alpha(c_x) = c_{f(x)}$ for every $x \in X$. Then f extends to an isomorphism $\iota_f : X^* \rightarrow Y^*$ and ι_f induces an isomorphism $\alpha_f = \iota_f \square \iota_f : \mathbf{End}(X^*) \rightarrow \mathbf{End}(Y^*)$. Consider the automorphism $\beta = \alpha_f^{-1} \alpha$ of $\mathbf{End}(X^*)$. Then $\beta(c_x) = \alpha_f^{-1}(\alpha(c_x)) = \alpha_f^{-1}(c_{f(x)}) = c_{f^{-1}(f(x))} = c_x$ for every $x \in X$ and β is stable. By Corollary 3.4 and Lemmas 3.5 and 3.6, β is an identity automorphism Δ or a mirror automorphism μ . In the former case, $\alpha_f^{-1} \alpha = \Delta$, and hence $\alpha = \alpha_f$, the automorphism induced by ι_f . In the latter case, $\alpha_f^{-1} \alpha = \mu$, whence $\alpha = \alpha_f \mu$, the automorphism induced by $\bar{\iota}_f$. □

Proof of Theorem 2. If $X = Y$, Theorem 1 becomes the first part of Theorem 2. Thus we prove here the second part of Theorem 2 for $|X| > 1$.

Obviously, $\{\Delta, \mu\}$ is a two-element group of automorphisms of $\mathbf{End}(X^*)$. It is isomorphic to C_2 . Define a mapping $i : \mathbf{Aut}(\mathbf{End}(X^*)) \rightarrow \mathbf{Aut}(X) \times C_2$ as follows: if $\alpha \in \mathbf{Aut}(\mathbf{End}(X^*))$, then $i(\alpha) = (f, \Delta)$ for $\alpha = \alpha_f$, and $i(\alpha) = (f, \mu)$ for $\alpha = \alpha_f \mu$. Clearly, i is a bijection. Also, i is a homomorphism because $\alpha_g \circ \alpha_f = \alpha_{g \circ f}$ and $\mu \circ \alpha = \alpha \circ \mu$ for all $f, g \in \mathbf{Aut}(X)$ and $\alpha \in \mathbf{Aut}(\mathbf{End}(X^*))$. Thus i is an isomorphism of $\mathbf{Aut}(\mathbf{End}(X^*))$ onto $\mathbf{Aut}(X) \times C_2$. □

Remark 4.1. Let \mathbf{L} be the variety of all left zero semigroups (that is, the semigroups satisfying the identity $x_1x_2 = x_1$). It is easily seen that every semigroup in this variety is free. If X is a left zero semigroup, then every transformation $X \rightarrow X$ is an endomorphism of X , and hence $\mathbf{End}(X)$ is merely the full transformation semigroup \mathcal{T}_X of the set X . Then $\mathbf{Aut}(\mathbf{End}(X)) = \mathbf{Aut}(\mathcal{T}_X)$. Schreier [5] proved that all automorphisms of \mathcal{T}_X are inner and $\mathbf{Aut}(\mathcal{T}_X)$ is isomorphic to $\mathbf{Aut}(X)$, the symmetric group on X .

Let \mathbf{Z} be the variety of all zero semigroups (that is, the semigroups satisfying the identity $x_1x_2 = x_3x_4$). They are semigroups $X^0 = X \cup \{0\}$, $0 \notin X$, with zero 0 such that $xy = 0$ for all $x, y \in X^0$. Every semigroup in this variety is free. Every partial transformation f of X corresponds bijectively to an endomorphism \bar{f} of X^0 :

for every $s \in X^0$, $\bar{f}(s) = f(s)$ if $f(s)$ is defined, and $\bar{f}(s) = 0$ otherwise. Clearly, \mathfrak{F}_X , the semigroup of all partial transformations of X , is naturally isomorphic to $\mathbf{End}(X^0)$ under the correspondence $f \mapsto \bar{f}$, and hence $\mathbf{Aut}(\mathbf{End}(X))$ is isomorphic to $\mathbf{Aut}(\mathfrak{F}_X)$. As proved by Shutov [6], all automorphisms of \mathfrak{F}_X are inner and $\mathbf{Aut}(\mathfrak{F}_X)$ is isomorphic to $\mathbf{Aut}(X)$, the symmetric group on X .

REFERENCES

1. E. Formanek, *A question of B. Plotkin about the semigroup of endomorphisms of a free group*, Proc. American Math. Society **130**(2001), 935-937. MR **2002h**:20032
2. B. I. Plotkin, *Seven Lectures on the Universal Algebraic Geometry*, Preprint, Institute of Mathematics, Hebrew University, 2000.
3. Boris M. Schein and Beimnet Teclezghi, *Endomorphisms of finite full transformation semigroups*, Proc. American Math. Soc. **126**(1998), 2579-2587. MR **99b**:20093
4. Б. М. Шайн, *Атомные полугруппы и инволютированные полугруппы*, Известия Высш. Учебн. Завед., Математика, **1965**, №3, 172-184. In Russian: B. M. Schein, *Atomic semigroups and involuted semigroups*, Izvestiya Vyssh. Uchebn. Zaved. (Matematika), **1965**, no. 3, 172-184.
5. J. Schreier, *Über Abbildungen einer abstrakten Menge auf ihre Teilmengen*, Fundamenta Mathematica **28**(1936), 261-264.
6. Э. Г. Шутов, *Гомоморфизмы полугруппы всех частичных преобразований*, Известия Высш. Учебн. Завед., Математика, **1961**, №3, 178-184. In Russian: È. G. Shutov, *Homomorphisms of the semigroup of all partial transformations*, Izvestiya Vyssh. Uchebn. Zaved. (Matematika), **1961**, no. 3, 178-184.
7. S. M. Ulam, *A Collection of Mathematical Problems*, Interscience Publishers, New York, 1960. MR **22**:10884

DEPARTMENT OF MATHEMATICS, BEN GURION UNIVERSITY OF THE NEGEV, BEER SHEVA, ISRAEL
E-mail address: gmash@math.bgu.ac.il

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF ARKANSAS, FAYETTEVILLE,
 ARKANSAS 72701
E-mail address: bschein@uark.edu