

ON ASYMMETRY OF THE FUTURE AND THE PAST FOR LIMIT SELF-JOININGS

OLEG N. AGEEV

(Communicated by Michael Handel)

ABSTRACT. Let Δ_T be an off-diagonal joining of a transformation T . We construct a non-typical transformation having asymmetry between limit sets of Δ_{T^n} for positive and negative powers of T . It follows from a correspondence between subpolymorphisms and positive operators, and from the structure of limit polynomial operators. We apply this technique to find all polynomial operators of degree 1 in the weak closure (in the space of positive operators on L_2) of powers of Chacon's automorphism and its generalizations.

INTRODUCTION

In [1], D. Rudolph introduced the notion of joinings. This notion turned out to be very fruitful (see [2]–[6]). It is known that every automorphism or endomorphism can be characterized both as a measure on a graph of this map, i.e., a joining or, more generally, a polymorphism [7], and as an operator on L_2 . This provides a way to study the structure of joinings by operator methods, and automorphisms by joinings.

It is not difficult to show that limit sets for positive and negative powers of a transformation T are equal in the space of all automorphisms if T is rigid. If T is not rigid, then these sets are empty. Moreover, for rigid or mixing transformations (see Section 1) these limit sets are also equal in the space of all linear operators on L_2 .

Nevertheless we prove (Theorem 2.4) that there exist rank-one transformations T with different limit sets of off-diagonal joinings for positive and negative powers of T or, in terms of operators, that the sets of limit operators are different.

We apply a new approach, via limit polynomials. This approach recently gave a solution to an old problem of Rokhlin (see [8], [9]) (see also references [10]–[13] about this problem) and answered some other well-known questions. See for example [8], which contains an answer to a question of Katok.

In Section 1 we introduce subpolymorphisms and a natural homeomorphism between the space of such measures and a subspace of positive operators on L_2 .

Received by the editors April 19, 2001.

2000 *Mathematics Subject Classification*. Primary 37Axx, 28D05, 28D15, 20M14, 47B65; Secondary 47A05, 47A15, 47Dxx, 60Gxx.

Key words and phrases. Joinings, Chacon's automorphism, weak operator convergence.

The author was supported in part by the Max Planck Institute of Mathematics, Bonn, and RFBR Grants #100-15-96107, #99-01-01104.

Note that, for any transformation T , the space of its self-joinings lies in the space of subpolymorphisms.

Theorem 2.1 describes all possible limit linear polynomials in $T_{(k)}$ for powers of transformations $T_{(k)}$, where $T_{(k)}$ are constructed in Section 1. In particular, this implies that only one linear polynomial $1/2E + 1/2T$ is a limit of powers for the classical Chacon transformation T (see [3], [14], [15]).

In Section 3 we show that the future and the past for limit self-joinings cannot be completely different for any automorphism. Moreover, their intersection contains an abelian semigroup that is trivial (i.e., $\{\mu \times \mu\}$) only for mixing transformations.

1. BASIC DEFINITIONS AND NOTATIONS

Let T be a transformation defined on a non-atomic standard Borel probability space (X, \mathcal{F}, μ) . A transformation and a unitary operator on $L_2(\mu) : Tf(x) = f(Tx)$ are often called automorphisms and denoted by the same symbol T . It is clear that T is contained in the following space of positive operators on $L_2(\mu)$: $\mathcal{L}^+ = \{U : (Uf \in L_2 \text{ if } f \in L_2) \ \& \ (Ug \geq 0 \text{ if } g \geq 0)\}$ equipped with the weak operator convergence. Everywhere below the identity automorphism will be denoted by E . The group of all automorphisms of (X, \mathcal{F}, μ) , say $\mathbf{Aut}(\mu)$, becomes a completely metrizable topological group when endowed with the weak convergence of transformations ($T_n \rightarrow T$ iff for any measurable A we have $\mu(T_n(A) \Delta T(A)) \rightarrow 0$ as $n \rightarrow \infty$). Note that this topology is a restriction of the weak operator topology in \mathcal{L}^+ to the non-closed $\mathbf{Aut}(\mu)$. Denote by $C(T)$ the commutant of T , i.e., the set $\{S \in \mathbf{Aut}(\mu) : ST = TS\}$.

Let ν be a finite Borel measure on $X \times X$ with marginal measures, say $\pi_1\nu$ and $\pi_2\nu$, such that

$$(1.1) \quad \pi_i\nu(A) \leq \mu(A) \text{ for any } \mu\text{-measurable set } A.$$

Obviously, marginal measures are μ -absolutely continuous and $\|\nu\| = \nu(X \times X) \leq 1$. The set of all such measures, say $M(\mu)$, is a convex compact metrizable space with respect to the topology determined by

$$\mu_n \rightarrow \mu_0 \Leftrightarrow \mu_n(A \times B) \rightarrow \mu_0(A \times B)$$

for any μ -measurable sets A and B . Now we shall give the following definition.

Definition 1.1. Each measure from $M(\mu)$ is called a subpolymorphism.

Fix $T \in \mathbf{Aut}(\mu)$. The set $J_c(T, T)$ of c -self-joinings, i.e., $T \times T$ -invariant elements ν of $M(\mu)$, with $\|\nu\| = c$, is a closed subspace of $M(\mu)$ for any $c \in [0, 1]$. If T is ergodic, then each ν from $J_c(T, T)$ has $c\mu$ as marginal measures. This gives that $J(T, T) = J_1(T, T)$ is exactly the set of well-known self-joinings. Let $S \in C(T)$. As usual, by an off-diagonal joining Δ_S we mean a measure from $J(T, T)$ completely defined by $\Delta_S(A \times B) = \mu(S^{-1}A \cap B)$, where A, B are any μ -measurable sets. Denote by $LJ_+(T)$ and $LJ_-(T)$ the limit sets of Δ_{T^n} in $M(\mu)$ for all positive n and negative n respectively. It is clear that $LJ_{\pm}(T) \subseteq J(T, T)$. If T is mixing, then $LJ_{\pm}(T)$ consist of only one point $\mu \times \mu$. It is well known that T is weakly mixing iff $LJ_{\pm}(T)$ contain at least $\mu \times \mu$. We say T is rigid if $T^{n_k} \rightarrow E$ for some sequence $n_k \rightarrow \infty$.

Proposition 1.2. $LJ_-(T) = LJ_+(T)$ for rigid T .

Proposition 1.3. $LJ_+(T^*) = LJ_-(T) = \sigma LJ_+(T)$, where σ is the flip map (i.e., $\sigma(x, y) = (y, x)$), and S^* denotes the operator adjoint to S .

Corollary 1.4. $LJ_-(T) = LJ_+(T)$ iff $LJ_+(T)$ is invariant with respect to σ .

We leave simple proofs of the above statements to the reader. It seems to be well known that the set of rigid transformations is a dense G_δ -set in $\mathbf{Aut}(\mu)$ (see [16] regarding close results in different subclasses of the set of non-singular transformations). It turns out that $LJ_-(T) = LJ_+(T)$ for a typical transformation T .

1.1. Construction of $T_{(k)}$. We consider the following “generalized” Chacon’s automorphisms. For each $k \geq 3$, let $T_{(k)}$ be a rank-one transformation, where each column C_{n+1} is obtained by cutting C_n into k subcolumns, say $C_n(i)$, of equal width, placing a spacer only on the subcolumn $C_n(k - 1)$, and then stacking the subcolumn $C_n(i + 1)$ on top of $C_n(i)$ for $1 \leq i < k$. It is clear that $T_{(3)}$ is exactly Chacon’s automorphism. For the column C_n , let h_n be its height and let d_n be the measure of its one level, where $n \geq 1$.

1.2. Correspondence between positive operators and $M(\mu)$. Consider

$$\begin{aligned} \mathcal{L}_\mu^+ &= \{U \in \mathcal{L}^+ : \int_A U(\mathbf{1})d\mu \leq \mu(A) \\ &\& \int_A U^*(\mathbf{1})d\mu \leq \mu(A) \text{ for any } \mu\text{-measurable set } A\} \end{aligned}$$

with a restriction of the weak operator topology to this set. Obviously,

$$\begin{aligned} \mathcal{L}_\mu^+ &= \{U \in \mathcal{L}^+ : U(\mathbf{1}) \leq \mathbf{1} \\ &\& U^*(\mathbf{1}) \leq \mathbf{1} \text{ for a.e. } x \text{ with respect to } \mu\}. \end{aligned}$$

Proposition 1.5. *The natural correspondence given by*

$$(1.2) \quad \langle U_\nu f, g \rangle = \int f \otimes \bar{g}d\nu,$$

for any $f, g \in L_2(\mu)$, defines a linear homeomorphism, say ϕ , between the topological spaces $M(\mu)$ and \mathcal{L}_μ^+ .

Note that in (1.2) f and $U_\nu f$ are from different spaces $L_2(\mu)$, but we naturally identify these spaces.

Proof. Indeed, the right part of (1.2) is a linear bounded functional for every $f \in L_2(\mu)$ because

$$(1.3) \quad \left| \int f \otimes \bar{g}d\nu \right| \leq \|f\|_\nu \|g\|_\nu = \|f\|_{\pi_1\nu} \|g\|_{\pi_2\nu} \leq \|f\|_\mu \|g\|_\mu,$$

where $\|h\|_l$ means the norm of $h \in L_2(l)$. Thus $U_\nu f \in L_2(\mu)$ for each $f \in L_2(\mu)$. The remaining properties of the map ϕ are obvious. \square

Some properties of operators U_ν were considered in [17] due to Vershik for polymorphisms ν , i.e., for elements of $M(\mu)$ with exact equality in (1.1) for each A .

Remark 1.6. Clearly, $\phi(\Delta_T) = T$, where $T \in \mathbf{Aut}(\mu)$. Also, U_ν commutes with T if and only if $\nu \in J_{\|\nu\|}(T, T)$.

Note that

$$\int f \otimes \bar{g} d\nu = \int_X \bar{g}(y) d\pi_2 \nu \int_X f(x) d\nu_y(x) = \int_X \bar{g}(y) \rho_2(y) d\mu(y) \int_X f(x) d\nu_y(x),$$

where $\rho_2(y)$ is a density of $\pi_2 \nu$ with respect to μ , and $\nu_y(x)$ is a canonical system of conditional measures corresponding to ν . Hence, changing g to $U_\nu f$ in (1.2) and (1.3), we get

Corollary 1.7. \mathcal{L}_μ^+ is a compact convex metrizable space. For any subpolymorphism $\nu \in M(\mu)$, $\|U_\nu\| \leq 1$, and

$$U_\nu(f) = \rho_2(y) \int_X f(x) d\nu_y(x).$$

Remark 1.8. The space \mathcal{L}_μ^+ is a semigroup, and closed with respect to taking parts of operators, i.e., if $0 \leq V \leq U$ and $U \in \mathcal{L}_\mu^+$, then $V \in \mathcal{L}_\mu^+$.

Remark 1.9. It is readily seen that $\phi(LJ_+(T))$ and $\phi(LJ_-(T))$ are T -invariant closed semigroups.

2. LIMIT POLYNOMIALS

The following theorem completely determines the simplest limit polynomial.

Theorem 2.1. Let $\mathcal{P}_1[x]$ be the set of polynomials of degree at most 1. Then

$$\phi(LJ_+(T_{(k)})) \cap \mathcal{P}_1[T_{(k)}] = \left\{ \frac{1}{k-1} E + \frac{k-2}{k-1} T_{(k)} \right\}.$$

Proof. In order to prove Theorem 2.1 we need some definitions and a technical lemma. For clarity we restrict our attention to the case $k = 4$. The proof for $k \neq 4$ is analogous.

Let $m_i \rightarrow +\infty$ as $i \rightarrow +\infty$. Fix i and choose n such that $h_n \leq m_i < h_{n+1}$. Consider the $(n + 1)^{\text{st}}$ column. Number its levels by $1, 2, \dots, 4h_n + 1$ from the base consequently. There exists i_0 such that the $(3h_n - m_i)^{\text{th}} \pmod{h_{n+1}}$ level belongs to $C_n(i_0)$ (we put $i_0 = 4$ in the remaining case $m_i = 4h_n$). Let p_i be a number of higher levels in $C_n(i_0)$. Denote by $B_i(j)$ the set of the top p_i levels of $C_n(j)$ and $A_i(j) = C_n(j) \setminus B_i(j)$ ($A_i(j)$ or $B_i(j)$ can be empty). Define $O_i = \{x \in C_{n+1} : T_{(4)}^r x \in C_{n+1} \text{ for } r = 1, 2, \dots, h_{n+1}\}$. It is clear that on each level a measure of x from O_i is $2/3d_{n+1}$.

Define operators

$$Q_i = \chi_{A_i(i_0)} T_{(4)}^{m_i}, \quad R_i = \chi_{B_i(i_0)} T_{(4)}^{m_i}, \\ Q'_i = \chi_{O_i} Q_i, \quad R'_i = \chi_{O_i} R_i.$$

Note that Q_i, Q'_i, R_i, R'_i are in \mathcal{L}_μ^+ because they are less than $T_{(4)}^{m_i}$.

Lemma 2.2.

$$(2.1) \quad T_{(4)}^{m_i} - P_i \rightarrow 0 \text{ as } i \rightarrow +\infty,$$

where

- (1) $P_i = (4/3T_{(4)}^{-1} + 8/3E)Q_i + (1/3T_{(4)}^{-1} + 2E + 5/3T_{(4)})R_i$, if $h_n \leq m_i < 2h_n$.
- (2) $P_i = (1/3T_{(4)}^{-2} + 2T_{(4)}^{-1} + 5/3E)Q_i + (2/3T_{(4)}^{-1} + 8/3E + 2/3T_{(4)})R_i$, if $2h_n \leq m_i < 3h_n$.

$$(3) \quad P_i = (2/3T_{(4)}^{-2} + 8/3T_{(4)}^{-1} + 2/3E)3/2Q'_i + (4/3T_{(4)}^{-1} + 8/3E)3/2R'_i, \text{ if } 3h_n \leq m_i < h_{n+1}.$$

Proof. It is enough to show (2.1) on pairs of functions from some dense set in $L_2(\mu)$. Therefore we can assume that f and g are constant, say $f_n(j)$ and $g_n(j)$, on each j^{th} level of C_n for sufficiently large n .

Obviously, if $\mu(D_i) \rightarrow 0$, then $\chi_{D_i}T^{m_i} \rightarrow 0$. Thus

$$(2.2) \quad T_{(4)}^{m_i} - \sum_j \chi_{A_i(j)}T_{(4)}^{m_i} - \sum_j \chi_{B_i(j)}T_{(4)}^{m_i} \rightarrow 0.$$

Next we will calculate the connection between components of each sum in (2.2), using that $T_{(4)}^{m_i}f$ has a “regular” structure on sets $A_i(j)$ and $B_i(j)$ and g is independent of j . Consider the $(n + 1)^{\text{st}}$ column.

1. If $m_i < 2h_n$, then $i_0 = 2$. Clearly,

$$(2.3) \quad \langle \chi_{A_i(1)}T_{(4)}^{m_i}f, g \rangle = d_{n+1} \sum_{l=1}^{h_n-p_i} f_n(l + p_i)\bar{g}_n(l) = \langle Q_i f, g \rangle,$$

$$\langle \chi_{B_i(1)}T_{(4)}^{m_i}f, g \rangle = d_{n+1} \sum_{l=h_n-p_i+1}^{h_n} f_n(l + p_i - h_n)\bar{g}_n(l) = \langle T_{(4)}R_i f, g \rangle.$$

Here and next values of f and g can be written incorrectly on bases and tops of $A_i(j)$ and $B_i(j)$, but this fact is not essential for the convergence of such operators. It is clear that

$$\chi_{A_i(1)}T_{(4)}^{m_i} - Q_i \rightarrow 0,$$

$$\chi_{B_i(1)}T_{(4)}^{m_i} - T_{(4)}R_i \rightarrow 0.$$

Analogously,

$$\chi_{A_i(3)}T_{(4)}^{m_i} - T_{(4)}^{-1}Q_i \rightarrow 0.$$

By construction of $T_{(4)}$, the function $f(T_{(4)}^{m_i}x)$ has at most two values on each level from $B_i(3), A_i(4), B_i(4)$ up to two base levels of $B_i(3)$. The first one is exactly at x from O_i , and the second one is at the remaining part of this level. Indeed, fix some x from such a level. It is clear that $N(T_{(4)}^{m_i}x) = N(x) + m_i \pmod{h_{n+1}}$ for $x \in O_i$, where $N(y)$ means a number of the level having y . The set $B_i(3)$ starts from the $(h_{n+1} - m_i)^{\text{th}}$ level. Thus $\{x, T_{(4)}x, \dots, T_{(4)}^{m_i}x\} \setminus C_{n+1} \neq \emptyset$, if x is not in O_i . Therefore $N(T_{(4)}^{m_i}x) = N(x) + m_i - 1 \pmod{h_{n+1}}$.

Thus

$$\begin{aligned} \langle \chi_{B_i(3)}T_{(4)}^{m_i}f, g \rangle &= \langle \chi_{B_i(3) \cap O_i}T_{(4)}^{m_i}f, g \rangle + \langle \chi_{B_i(3) \setminus O_i}T_{(4)}^{m_i}f, g \rangle \\ &= \frac{2}{3}d_{n+1} \sum_{l=h_n-p_i+2}^{h_n} f_n(l + p_i - h_n - 1)\bar{g}_n(l) \\ &\quad + \frac{1}{3}d_{n+1} \sum_{l=h_n-p_i+3}^{h_n} f_n(l + p_i - h_n - 2)\bar{g}_n(l) \\ &= \frac{2}{3}\langle R_i f, g \rangle + \frac{1}{3}\langle T_{(4)}^{-1}R_i f, g \rangle. \end{aligned}$$

Hence

$$\chi_{B_i(3)}T_{(4)}^{m_i} - \left(\frac{2}{3}E + \frac{1}{3}T_{(4)}^{-1}\right)R_i \rightarrow 0.$$

In the same way, we get

$$\chi_{A_i(4)}T_{(4)}^{m_i} - \left(\frac{2}{3}E + \frac{1}{3}T_{(4)}^{-1}\right)Q_i \rightarrow 0,$$

$$\chi_{B_i(4)}T_{(4)}^{m_i} - \left(\frac{2}{3}T_{(4)} + \frac{1}{3}E\right)R_i \rightarrow 0.$$

This completes the calculation in the case 1.

2. Here $i_0 = 1$. As above, we have

$$\begin{aligned} \chi_{A_i(2)}T_{(4)}^{m_i} - T_{(4)}^{-1}Q_i &\rightarrow 0, \quad \chi_{B_i(2)}T_{(4)}^{m_i} - \left(\frac{2}{3}E + \frac{1}{3}T_{(4)}^{-1}\right)R_i \rightarrow 0, \\ \chi_{A_i(3)}T_{(4)}^{m_i} - \left(\frac{2}{3}T_{(4)}^{-1} + \frac{1}{3}T_{(4)}^{-2}\right)Q_i &\rightarrow 0, \quad \chi_{B_i(3)}T_{(4)}^{m_i} - \left(\frac{2}{3}E + \frac{1}{3}T_{(4)}^{-1}\right)R_i \rightarrow 0, \\ \chi_{A_i(4)}T_{(4)}^{m_i} - \left(\frac{2}{3}E + \frac{1}{3}T_{(4)}^{-1}\right)Q_i &\rightarrow 0, \quad \chi_{B_i(4)}T_{(4)}^{m_i} - \left(\frac{2}{3}T_{(4)} + \frac{1}{3}E\right)R_i \rightarrow 0. \end{aligned}$$

3. In this case, $i_0 = 4$, and $f(T_{(4)}^{m_i}x)$ has two values on each level, except for from $A_i(1)$. Using

$$\langle Q'_i f, g \rangle = \frac{2}{3}d_{n+1} \sum_{l=1}^{h_n-p_i} f_n(l+p_i)\bar{g}_n(l),$$

calculate

$$\begin{aligned} \langle \chi_{A_i(1)}T_{(4)}^{m_i} f, g \rangle &= d_{n+1} \sum_{l=1}^{h_n-p_i} f_n(l+p_i-1)\bar{g}_n(l) = \frac{3}{2}\langle T_{(4)}^{-1}Q'_i f, g \rangle, \\ \langle \chi_{A_i(4)}T_{(4)}^{m_i} f, g \rangle &= \langle \chi_{A_i(4)\cap O_i}T_{(4)}^{m_i} f, g \rangle + \langle \chi_{A_i(4)\setminus O_i}T_{(4)}^{m_i} f, g \rangle \\ &= \langle Q'_i f, g \rangle + \frac{1}{3}d_{n+1} \sum_{l=1}^{h_n-p_i} f_n(l+p_i-1)\bar{g}_n(l) \\ &= \langle Q'_i f, g \rangle + \frac{1}{2}\langle T_{(4)}^{-1}Q'_i f, g \rangle. \end{aligned}$$

For $j = 2, 3$, we get

$$\begin{aligned} \langle \chi_{A_i(j)}T_{(4)}^{m_i} f, g \rangle &= \frac{2}{3}d_{n+1} \sum_{l=1}^{h_n-p_i} f_n(l+p_i-1)\bar{g}_n(l) \\ &\quad + \frac{1}{3}d_{n+1} \sum_{l=1}^{h_n-p_i} f_n(l+p_i-2)\bar{g}_n(l) \\ &= \langle T_{(4)}^{-1}Q'_i f, g \rangle + \frac{1}{2}\langle T_{(4)}^{-2}Q'_i f, g \rangle. \end{aligned}$$

As before, for any j

$$\begin{aligned} \langle \chi_{B_i(j)} T_{(4)}^{m_i} f, g \rangle &= \frac{2}{3} d_{n+1} \sum_{l=h_n-p_i+2}^{h_n} f_n(l+p_i-h_n-1) \bar{g}_n(l) \\ &\quad + \frac{1}{3} d_{n+1} \sum_{l=h_n-p_i+3}^{h_n} f_n(l+p_i-h_n-2) \bar{g}_n(l) \\ &= \langle R'_i f, g \rangle + \frac{1}{2} \langle T_{(4)}^{-1} R'_i f, g \rangle. \end{aligned}$$

□

1. Fix $P = aE + bT_{(4)}$. Let $T_{(4)}^{m_i} \rightarrow P$ for some $m_i \rightarrow +\infty$. It is clear that $a, b \geq 0$, because $P \in \mathcal{L}_\mu^+$. Choose a subsequence of m_i (if necessary) such that

$$(2.4) \quad \begin{aligned} Q_i^{(x_i)} &\rightarrow Q, \\ R_i^{(x_i)} &\rightarrow R, \end{aligned}$$

where $S_i^{(x_i)}$ means S_i or S'_i for each i , and our choice is completely determined by m_i as in Lemma 2.2. Obviously, $Q, R \in \mathcal{L}_\mu^+$. By construction of $T_{(4)}$,

$$\|T_{(4)} \chi_{D_i} - \chi_{D_i}\|_\mu^2 \leq 2d_{n+1} \rightarrow 0,$$

where D_i is $A_i(i_0)$ or $A_i(i_0) \cap O_i$. Thus

$$\|(T_{(4)} Q_i^{(x_i)} - Q_i^{(x_i)} T_{(4)}) f\|_\mu^2 \leq \|T_{(4)} \chi_{D_i} - \chi_{D_i}\|_\mu \|T_{(4)}^2 f\|_\mu \rightarrow 0,$$

for any $f \in L_2(\mu)$. Therefore Q commutes with $T_{(4)}$. Analogously, we have that R commutes with $T_{(4)}$. Denote by $\mathcal{P}_1^+[x]$ the subset of $\mathcal{P}_1[x]$ with non-negative coefficients.

Next we will show that if $P = \sum_{i=1}^l S_i$, where $S_i \in \mathcal{L}_\mu^+$, and S_i commute with $T_{(4)}$, then $S_i \in \mathcal{P}_1^+[T_{(4)}]$. Indeed, measures $\phi^{-1} S_i$ are $T_{(4)} \times T_{(4)}$ -invariant and absolutely continuous with respect to the subpolymorphism $\phi^{-1} P = a\Delta_E + b\Delta_{T_{(4)}}$. The transformation $T_{(4)} \times T_{(4)}$ is ergodic for measures Δ_E and $\Delta_{T_{(4)}}$. Hence every $T_{(4)} \times T_{(4)}$ -invariant part of the measure Δ_E ($\Delta_{T_{(4)}}$) is $c\Delta_E$ ($c\Delta_{T_{(4)}}$) for some $c > 0$. This gives $\phi^{-1} S_i = a_i \Delta_E + b_i \Delta_{T_{(4)}}$ for some $a_i, b_i \geq 0$.

From Lemma 2.2 and (2.4) we have that

$$(2.5) \quad P = UQ + VR,$$

where (U, V) is at least one of the following pairs:

$$\begin{aligned} &(4/3T_{(4)}^{-1} + 8/3E, 1/3T_{(4)}^{-1} + 2E + 5/3T_{(4)}), \\ &(1/3T_{(4)}^{-2} + 2T_{(4)}^{-1} + 5/3E, 2/3T_{(4)}^{-1} + 8/3E + 2/3T_{(4)}), \\ &(T_{(4)}^{-2} + 4T_{(4)}^{-1} + E, 2T_{(4)}^{-1} + 4E). \end{aligned}$$

In any case P contains $c_1 Q$ and $c_2 R$ as parts. Thus $Q, R \in \mathcal{P}_1^+[T_{(4)}]$. Obviously,

$$(2.6) \quad 1 = \langle T_{(4)}^{m_i} \mathbf{1}, \mathbf{1} \rangle \rightarrow \langle P \mathbf{1}, \mathbf{1} \rangle.$$

This implies that $P \neq 0$. Therefore equality (2.5) is possible only when $P = (4/3T_{(4)}^{-1} + 8/3E)cT_{(4)}$. It remains to mention, using (2.6), that $c = 1/4$.

2. The proof that $1/3E + 2/3T_{(4)} \in \phi(LJ_+(T_{(4)}))$ is almost obvious. Namely, consider $m_i = h_n$. We obtain $i_0 = 2, p_i = 0, R_i = 0$. Thus (2.3) gives

$$\langle Q_i f, g \rangle = d_{n+1} \sum_{l=1}^{h_n} f_n(l) \bar{g}_n(l) = \frac{d_n}{4} \sum_{l=1}^{h_n} f_n(l) \bar{g}_n(l) = \frac{1}{4} \langle f, g \rangle.$$

This implies that

$$\langle P_i f, g \rangle = \langle (\frac{1}{3}T_{(4)}^{-1} + \frac{2}{3}E)f, g \rangle.$$

Thus Theorem 2.1 follows from Lemma 2.2 and Remark 1.9. □

Remark 2.3. By the same argument as in [3], it is not difficult to show that $T_{(k)}$ have minimal self-joinings. Then Q, R can be written in the following form:

$$\alpha \int + \sum_j a_j T_{(k)}^j,$$

where \int is the orthogonal projection onto the space of constants, and $0 \leq \alpha, 0 \leq a_i$. This gives that the first part of Theorem 2.1 also follows directly from (2.5).

Our main result is the following.

Theorem 2.4. $LJ_+(T_{(k)}) \neq LJ_-(T_{(k)})$ for $k > 3$.

Proof. Indeed, it is clear that

$$\phi(LJ_+(T^*)) = \{U_\nu^* : \nu \in LJ_+(T)\}.$$

Therefore, using Proposition 1.3, Remark 1.9, and Theorem 2.1, we have

$$\phi(LJ_-(T_{(k)})) \cap \mathcal{P}_1[T_{(k)}] = \{ \frac{k-2}{k-1}E + \frac{1}{k-1}T_{(k)} \},$$

and Theorem 2.4 is proved. □

3. CLOSING REMARKS

Proposition 3.1. For any $T \in \mathbf{Aut}(\mu)$

$$LJ_+(T) \cap LJ_-(T) \neq \emptyset.$$

This is an immediate consequence of the next proposition.

Proposition 3.2.

$$(3.1) \quad \phi(LJ_+(T)) \cap \phi(LJ_-(T)) \supseteq \{T_+T_- : T_\pm \in \phi(LJ_\pm(T))\},$$

and

$$\{T_+T_- : T_\pm \in \phi(LJ_\pm(T))\} = \{ \int \} \Leftrightarrow T \text{ is mixing},$$

where \int is defined as in Remark 2.3.

Proof. Fix $T_\pm \in \phi(LJ_\pm(T))$, and $n_i, k_i \rightarrow +\infty$ such that $T^{k_i} \rightarrow T_+, T^{-n_i} \rightarrow T_-$. Consider also a dense set of functions from $L_2(\mu)$, say f_l . For each $\epsilon > 0$ and $m \in \mathbb{N}$, choose i such that

$$|\langle T^{-n_i} f_{l_1}, T_+^* f_{l_2} \rangle - \langle T_- f_{l_1}, T_+^* f_{l_2} \rangle| < \epsilon,$$

for all $l_1, l_2 \leq m$. Finally choose $j = j(i)$ such that $k_{j(i)} - n_i > m$ and

$$|\langle T^{k_{j(i)}} T^{-n_i} f_{l_1}, f_{l_2} \rangle - \langle T_+ T^{-n_i} f_{l_1}, f_{l_2} \rangle| < \epsilon,$$

for all $l_1, l_2 \leq m$. Thus

$$|\langle T^{k_{j(i)}-n_i} f_{l_1}, f_{l_2} \rangle - \langle T_- f_{l_1}, T_+^* f_{l_2} \rangle| < 2\epsilon.$$

Therefore $T^{k_{j(i)}-n_i} \rightarrow T_+ T_-$, where $k_{j(i)} - n_i \rightarrow +\infty$.

Operators T_+ and T_- belong to the von Neumann algebra generated by T . Thus $T_+ T_- = T_- T_+$. This implies that arguing as above, we see that $T^{-n_{j'(i)}+k_i} \rightarrow T_+ T_-$, where $-n_{j'(i)} + k_i \rightarrow -\infty$.

The second part of the proof is more or less standard. Indeed, if T is not mixing, then there exists T_+ from $\phi(LJ_+(T)) \setminus \{f\}$. Thus $T_- = T_+^* \in \phi(LJ_-(T)) \setminus \{f\}$. Next for $S^* S = f$, where $S \in \phi(J(T, T))$, we have

$$\int f d\mu = 0 \Rightarrow \langle Sf, Sf \rangle = \langle S^* Sf, f \rangle = \langle \int f, f \rangle = 0.$$

This means that $Sf = 0$, and finally $S = f$. Therefore the operator $T_+ T_- = T_- T_+ = T_+^* T_+$ is not f . \square

Remark 3.3. Obviously, in (3.1) we have an exact equality if T is rigid or mixing. However, taking into account Remark 2.3, the operator $1/2E + 1/2T$ cannot be represented as $T_+ T_-$ for Chacon's transformation T . This yields that, in general, the left part of (3.1) is different from the right part.

REFERENCES

1. D. Rudolph, *An example of a measure preserving map with minimal self-joinings, and applications*, J. Analyse Math. **35** (1979), 97–122. MR **81e**:28011
2. M. Ratner, *Horocycle flows, joinings and rigidity of products*, Ann. of Math.(2) **118** (1983), 277–313. MR **85k**:58063
3. A del Junco, M. Rahe, and L. Swanson, *Chacon's automorphism has minimal self-joinings*, J. Analyse Math. **37** (1980), 276–284. MR **81j**:28027
4. J. King, *The commutant is the weak closure of the powers, for rank-1 transformations*, Erg. Theory and Dyn. Sys. **6** (1986), 363–384. MR **88a**:28021
5. B. Host, *Mixing of all orders and pairwise independent joinings of systems with singular spectrum*, Isr. J. Math. **76** (1991), 289–298. MR **93k**:28022
6. A del Junco and D. Rudolph, *On ergodic actions whose self-joinings are graphs*, Erg. Theory and Dyn. Sys. **7** (1987), 531–557. MR **89e**:28029
7. A. Vershik, *Dynamic theory of growth in groups: entropy, boundaries, examples*, Uspekhi Mat. Nauk **55** (2000), 59–128; English transl., Russian Math. Surv. **55** (2000), 667–733. MR **2001m**:37019
8. O. Ageev, *On ergodic transformations with homogeneous spectrum*, J. Dynam. Control Systems **5** (1999), 149–152. MR **99m**:28034
9. V. Ryzhikov, *Transformations having homogeneous spectra*, J. Dynam. Control Systems **5** (1999), 145–148. MR **99m**:28038
10. A. Katok, Ja. Sinai, and A. Stepin, *The theory of dynamical systems and general transformation groups with invariant measure*, Mathematical analysis, Vol. 13 (Russian), pp. 129–262. (errata insert) Akad. Nauk SSSR VINITI, Moscow, 1975. MR **58**:28430
11. G. Goodson, *A survey of recent results in the spectral theory of ergodic dynamical systems*, J. Dynam. Control Systems **5** (1999), 173–226. MR **2000f**:28021
12. E. Robinson, Jr., *Ergodic measure preserving transformations with arbitrary finite spectral multiplicities*, Invent. Math. **72** (1983), 299–314. MR **85a**:28014
13. O. Ageev, *The spectral multiplicity function and geometric representations of interval exchange transformations*, Math. Sb. **190** (1999), 3–28; English transl., Sb. Math. **190** (1999), 1–28. MR **2000m**:28015
14. D. Rudolph, *Fundamentals of measurable dynamics. Ergodic theory on Lebesgue spaces*, The Clarendon Press, Oxford University Press, New York, 1990. MR **92e**:28006
15. O. Ageev, *The spectrum of Cartesian powers of classical automorphisms*, Math. Notes **68** (2000), 547–551. MR **2001m**:37014

16. O. Ageev, C. Silva, *Genericity of rigidity and multiple recurrence for infinite measure preserving and nonsingular transformations*, preprint.
17. M. Lemańczyk, B. Host, J.-P. Thouvenot, *Gaussian automorphisms whose ergodic self-joinings are Gaussian*, *Fund. Math.* **164** (2000), 253–293. MR **2001h:37009**

DEPARTMENT OF MATHEMATICS, MOSCOW STATE TECHNICAL UNIVERSITY, 2ND BAUMANSKAYA ST. 5, 105005 MOSCOW, RUSSIA

E-mail address: `ageev@mx.bmstu.ru`