

## THE BANACH ALGEBRA INDUCED BY A DOUBLE CENTRALIZER

ETIENNE DESQUITH

(Communicated by David R. Larson)

ABSTRACT. Given a Banach algebra  $A$ , R. Larsen defined, in his book “*An introduction to the theory of multipliers*”, a Banach algebra  $A_T$  by means of a *multiplier*  $T$  on  $A$ , and essentially used it in the case of a *commutative semisimple* Banach algebra  $A$  to prove a result on multiplications which preserve regular maximal ideals. Here, we consider the analogue Banach algebra  $\mathcal{A}_R$  induced by a bounded *double centralizer*  $\langle L, R \rangle$  of a Banach algebra  $A$ . Then, our main concern is devoted to the relationships between  $L$ ,  $R$ , and the algebras of bounded double centralizers  $\mathcal{W}(A)$  and  $\mathcal{W}(\mathcal{A}_R)$  of  $A$  and  $\mathcal{A}_R$ , respectively. By removing the assumption of semisimplicity, we generalize some results proven by Larsen.

### 1. PRELIMINARIES

Unless otherwise stated, we shall adopt throughout the sequel the following conventions:  $A$  will be an arbitrary Banach algebra, and the Banach algebra of all (resp. bounded) linear operators on it will be denoted by  $\mathcal{L}(A)$  (resp.  $B(A)$ ). The maps  $\lambda$  and  $\mu$ , such that  $a \mapsto \lambda_a$  and  $a \mapsto \mu_a$  for  $a \in A$ , are the usual left and right regular representations of  $A$ . Composition of mappings will be denoted by simple juxtaposition.

**Definition 1.1.** A left (resp. right) *centralizer* of  $A$  is an element  $L \in \mathcal{L}(A)$  (resp.  $R \in \mathcal{L}(A)$ ), satisfying  $L(ab) = L(a)b$  (resp.  $R(ab) = aR(b)$ ),  $\forall a, b \in A$ .

A *double centralizer* of  $A$  is a pair  $\langle L, R \rangle$  where  $L$  (resp.  $R$ ) is a left (resp. right) centralizer, and which together satisfy the following *Double Centralizer Property (DC-Property)*:

$$(1) \quad aL(b) = R(a)b, \quad \forall a, b \in A.$$

The algebra  $\mathcal{W}(A)$  of *bounded double centralizers* of  $A$  is the set of double centralizers with pointwise linear operations and with product and norm given by

$$(2) \quad \begin{aligned} \langle L, R \rangle \langle T, S \rangle &= \langle LT, SR \rangle, \quad \|\langle L, R \rangle\| = \text{Max}\{\|L\|; \|R\|\}, \\ \forall \langle L, R \rangle, \langle T, S \rangle &\in \mathcal{W}(A). \end{aligned}$$

Relation (2) above ensures the automatic continuity of the elements  $L$  and  $R$  of any double centralizer  $\langle L, R \rangle$  of  $A$ . For the definitions of a *multiplier* and of the

---

Received by the editors November 28, 2001 and, in revised form, February 4, 2002.

2000 *Mathematics Subject Classification.* Primary 47D30.

*Key words and phrases.* Annihilator, Banach algebra, double centralizer, multiplier.

This work was supported by the Abdus Salam ICTP Associateship scheme (Trieste/Italy).

left annihilator  $LAn(A)$ , right annihilator  $RAn(A)$  and annihilator  $An(A)$  of  $A$ , the reader should refer to [3] and [4].  $A$  is said to have no annihilators if  $An(A) = LAn(A) = RAn(A) = \{0\}$ .

## 2. THE BANACH ALGEBRA $\mathcal{A}_R$

**Proposition 2.1.** *Let  $\langle L, R \rangle \in \mathcal{W}(A)$  such that  $\|R\| \leq 1$ , and define*

$$(3) \quad a \cdot b = R(a) b \quad , \quad \forall a, b \in A .$$

(i) *Relation (3) endows  $A$  with a Banach algebra structure denoted by  $\mathcal{A}_R$ .*

(ii)  *$R$  (resp.  $L$ ) is a homomorphism from  $\mathcal{A}_R$  onto the image  $Im(R)$  (resp.  $Im(L)$ ) of  $R$  (resp.  $L$ ).*

(iii) *If  $A$  has no annihilator, or if  $A$  admits a two-sided approximate identity, then the kernels of  $L$  and  $R$  satisfy  $Ker(R) = LAn(\mathcal{A}_R)$ ,  $Ker(L) = RAn(\mathcal{A}_R)$ , and hence  $An(\mathcal{A}_R) = Ker(L) \cap Ker(R)$ .*

*Proof.* (i) and (ii) are obtained by straightforward computations. It is also easy to show the relations  $R[LAn(\mathcal{A}_R)] \subset LAn(A)$  and  $L[RAn(\mathcal{A}_R)] \subset RAn(A)$ , from which (iii) follows, since  $A$  is without annihilators.  $\square$

**Remark 2.2.** It should be noticed that assuming  $\|L\| \leq 1$ , and defining the product in relation (3) as  $a \cdot b = a L(b)$ ,  $\forall a, b \in A$ , gives the same algebra  $\mathcal{A}_R$ . Therefore, when dealing in the sequel with the algebra  $\mathcal{A}_R$ , we shall say that  $\mathcal{A}_R$  is induced or defined by  $\langle L, R \rangle$ . Considering  $\mathcal{A}_R$  as a left Banach module over itself, the map  $R$  satisfying relation (3) is an example of a bounded module map from the module into the algebra (cf. [1]).

As an immediate consequence of Proposition 2.1 above, we have:

**Corollary 2.3.** *Let  $A$  have no annihilators. Then, for any  $\langle L, R \rangle \in \mathcal{W}(A)$ ,  $L$  (resp.  $R$ ) is one-to-one implies that  $An(\mathcal{A}_R) = \{0\}$ . Moreover,  $\mathcal{A}_R$  has no annihilators, if and only if  $L$  and  $R$  are both one-to-one.*  $\square$

**Remark 2.4.** If  $A$  is commutative and without annihilators, then  $\langle L, R \rangle$  is a multiplier  $L = R = T$  and therefore  $\mathcal{A}_R$  coincides with the algebra  $A_T$  of R. Larsen, and is also commutative.

For  $\langle L, R \rangle$  defining  $\mathcal{A}_R$ , direct computations show that the condition  $\langle T, S \rangle \in \mathcal{W}(\mathcal{A}_R)$  implies  $\langle LT, RS \rangle \in \mathcal{W}(A)$ .

As noted by T. W. Palmer in [4], page 26, the set of left (resp. right) centralizers is contained in the commutant  $(\mu_A)^c$  (resp.  $(\lambda_A)^c$ ) of  $\mu_A$  (resp.  $\lambda_A$ ). Moreover, as pointed out in [2], page 775, if  $A$  has no annihilators, then given two arbitrary elements  $\langle L, R \rangle$  and  $\langle T, S \rangle$  of  $\mathcal{W}(A)$ , their components cross-commute, i.e.  $TR = RT$  and  $LS = SL$ . As a consequence, any  $\langle L, R \rangle \in \mathcal{W}(A)$  satisfies  $LR = RL$ .

For every  $\langle L, R \rangle \in \mathcal{W}(A)$ , let  $Im(L)$  (resp.  $Im(R)$ ) denote the image of  $L$  (resp.  $R$ ), and define  $I_{LR} = Im(L) \cap Im(R)$ . Then we have:

**Proposition 2.5.** *Let  $A$  have no annihilators, and let  $\langle L, R \rangle \in \mathcal{W}(A) \setminus \{0\}$ . If  $\overline{Im(L)} \neq A$  and  $\overline{Im(R)} \neq A$ , then each of  $L$  and  $R$  admits a nontrivial closed invariant subspace (the bar denotes the closure in the norm topology of  $A$ ); moreover,  $I_{LR} = \{0\}$  implies that  $L$  (resp.  $R$ ) cannot be one-to-one.*

*Proof.* We shall write  $L[Im(R)]$  and  $R[Im(L)]$  to denote the restrictions of the maps  $L$  and  $R$ , respectively, on the subspaces  $Im(R)$  and  $Im(L)$  of  $A$ . According to the above comments, the components of  $\langle L, R \rangle \in \mathcal{W}(A)$  commute. Hence the following hold:  $L[Im(R)] \subset I_{LR}$  and  $R[Im(L)] \subset I_{LR}$ , from which we derive  $L[\overline{Im(R)}] \subset \overline{Im(R)}$ ,  $R[\overline{Im(L)}] \subset \overline{Im(L)}$ . Then, if the images of  $L$  and  $R$  are not dense in  $A$ , these inclusions prove that each of  $L$  and  $R$  admits a nontrivial closed invariant subspace.  $L$  and  $R$  being nonzero, the same inclusions imply that  $L$  (resp.  $R$ ) cannot be one-to-one if  $I_{LR} = \{0\}$ .  $\square$

**Theorem 2.6.** *Let  $A$  and  $\mathcal{A}_R$  be without annihilators and let  $\langle L, R \rangle \in \mathcal{W}(A)$  be such that  $Im(L) \subset R[Im(L)]$ . Then,  $R$  is bijective, and  $R^{-1}$  is a right centralizer. The result remains clearly true if the roles of  $R$  and  $L$  are interchanged, and the right centralizer is replaced by the left centralizer.*

*Proof.* We have, according to Corollary 2.3 and the proof of Proposition 2.5,  $R[Im(L)] \subset Im(L)$ , which, together with the assumption on  $Im(L)$  implies the equality  $Im(L) = R[Im(L)]$ . It then follows that  $R$  is onto  $A$ , according to the following result:  $L(x_0) \in R[Im(L)] \implies x_0 \in Im(R)$ . Indeed, let us assume by contradiction that  $x_0 \notin Im(R)$ . Then,  $L$  being one-to-one, we get  $L(x_0) \neq LR(a)$ ,  $\forall a \in A$ . But since  $L$  commutes with  $R$ , it also follows that  $L(x_0) \neq RL(a)$ ,  $\forall a \in A$ , which means that  $L(x_0) \notin R[Im(L)]$ . Hence, the desired result is established, and one easily checks that  $R^{-1}$  is a right centralizer.  $\square$

Let us define

- (4)  $\mathcal{C}_\ell(A) = \{L \in B(A) ; \exists R \in B(A) ; \langle L, R \rangle \in \mathcal{W}(A)\} ;$
- (5)  $\mathcal{C}_r(A) = \{S \in B(A) ; \exists T \in B(A) ; \langle T, S \rangle \in \mathcal{W}(A)\} ;$
- (6)  $\Phi(\langle L, R \rangle) = L \quad , \quad \Psi(\langle L, R \rangle) = R \quad , \quad \forall \langle L, R \rangle \in \mathcal{W}(A) .$

Then we get:

**Theorem 2.7.** *Let  $A$  be a Banach algebra with an approximate identity bounded by one. Then  $\mathcal{C}_\ell(A)$  and  $\mathcal{C}_r(A)$  are Banach subalgebras of  $B(A)$ , and  $\Phi$  is a faithful representation of  $\mathcal{W}(A)$  on  $A$ , with  $\Phi(\mathcal{W}(A)) = \mathcal{C}_\ell(A)$  (resp.  $\Psi$  is a one-to-one anti-homomorphism of  $\mathcal{W}(A)$  into  $B(A)$ , with  $\Psi(\mathcal{W}(A)) = \mathcal{C}_r(A)$ ).*

*Proof.*  $\mathcal{C}_\ell(A)$  is clearly a linear subspace of  $B(A)$ . Moreover, if  $L, T \in \mathcal{C}_\ell(A)$ , then by definition, there exist  $R, S \in B(A)$  such that  $\langle L, R \rangle \in \mathcal{W}(A)$ ,  $\langle T, S \rangle \in \mathcal{W}(A)$ , with  $L = \Phi(\langle L, R \rangle)$  and  $T = \Phi(\langle T, S \rangle)$ . Hence,  $\langle L, R \rangle \langle T, S \rangle = \langle L T, S R \rangle \in \mathcal{W}(A)$ , and therefore  $L T = \Phi(\langle L T, S R \rangle) \in \mathcal{C}_\ell(A)$ . So  $\mathcal{C}_\ell(A)$  is a subalgebra of  $B(A)$  and more precisely, a Banach algebra. Indeed, if  $(L_n)_{n \geq 1}$  is a sequence in  $\mathcal{C}_\ell(A)$  converging to some  $L \in B(A)$ , then  $L$  is a left centralizer. Moreover, the corresponding sequence  $(R_n)_{n \geq 1}$  in  $B(A)$  such that  $L_n = \Phi(\langle L_n, R_n \rangle)$ ,  $\forall n \geq 1$ , converges to some  $R \in B(A)$ . To see this, we proceed as follows: given  $x \in A$ , we let  $x \mapsto \lambda_x$  be the left regular representation of  $A$  onto itself, and for fixed  $n, m \in \mathbb{N}$  and  $a \in A$ , we consider the map

$$(7) \quad A \ni b \mapsto (R_n - R_m)(a) b = \lambda_{(R_n - R_m)(a)}(b) .$$

Then we get

$$(8) \quad \|(R_n - R_m)(a)\| = \|\lambda_{(R_n - R_m)(a)}\| = \text{Sup} \{ \|(R_n - R_m)(a)(b)\| ; \|b\| \leq 1 \},$$

so that the following holds:

$$(9) \quad \|R_n - R_m\| \leq \text{Sup} \left\{ \text{Sup} \{ \|(R_n - R_m)(a)(b)\| ; \|b\| \leq 1 \} ; \|a\| \leq 1 \right\}.$$

Since  $\langle L_n, R_n \rangle, \langle L_m, R_m \rangle \in \mathcal{W}(A)$ , we have, for each pair of elements  $a, b \in A$ ,

$$(10) \quad R_n(a)b = aL_n(b) \quad \text{and} \quad R_m(a)b = aL_m(b),$$

and the equality (8) hence gives

$$\begin{aligned} \|R_n - R_m\| &\leq \text{Sup} \left\{ \|a\| \text{Sup} \{ \|L_n - L_m\| \|b\| ; \|b\| \leq 1 \} ; \|a\| \leq 1 \right\} \\ &\leq \|L_n - L_m\|. \end{aligned}$$

$(R_n)_{n \geq 1}$  is therefore a Cauchy sequence in  $B(A)$ , and hence converges to some  $R \in B(A)$ .

Now, for each  $n \in \mathbb{N}$  and for all  $a, b \in A$  the following relations are satisfied:

$$(11) \quad (i) \quad R_n(ab) = aR_n(b) \quad \text{and} \quad (ii) \quad R_n(a)b = aL_n(b),$$

and we get, taking the limits in equalities (i) and (ii) when  $n$  tends to infinity:  $R(ab) = aR(b)$  and  $R(a)b = aL(b)$ , that is,  $R$  is a right centralizer and  $\langle L, R \rangle \in \mathcal{W}(A)$ . Therefore  $L = \Phi(\langle L, R \rangle) \in \mathcal{C}_\ell(A)$ , and  $\mathcal{C}_\ell(A)$  is hence norm-closed in  $B(A)$ . That  $\mathcal{C}_r(A)$  is a Banach subalgebra of  $B(A)$  is obtained by quite similar arguments as in the case of  $\mathcal{C}_\ell(A)$ . That  $\Phi$  is a faithful representation of  $\mathcal{W}(A)$  on  $A$ , and  $\Psi$  a one-to-one anti-homomorphism of  $\mathcal{W}(A)$  into  $A$ , are obtained by routine computations.  $\square$

Given  $R \in B(A)$ , let  $\{R\}^c = \{S \in B(A) ; RS = SR\}$  denote the *commutant* of  $\{R\}$ . According to the comments following Remark 2.4, if  $\langle L, R \rangle \in \mathcal{W}(A)$  defines  $\mathcal{A}_R$  and if  $S \in \{R\}^c$  is such that  $\langle T, S \rangle \in \mathcal{W}(\mathcal{A}_R)$ , then  $\langle LT, RS \rangle = \langle LT, SR \rangle \in \mathcal{W}(A)$ , and it is “tempting” in this situation to write  $\langle LT, SR \rangle = \langle L, R \rangle \langle T, S \rangle$ . But, the product in the right-hand side of this equation would not make sense, unless  $\mathcal{W}(A)$  and  $\mathcal{W}(\mathcal{A}_R)$  are subalgebras of some larger Banach algebra. We construct in what follows such an algebra.

Consider the set  $\mathcal{D}_{P_0}(A)$  of all pairs  $\langle L, R \rangle$  with  $L \in B(A)$ ,  $R \in B(A)$ , made into a linear algebra under pointwise linear operations and with the same product as in  $\mathcal{W}(A)$ . Then  $\mathcal{D}_{P_0}(A)$  is a normed algebra when it is endowed with the same norm as in  $\mathcal{W}(A)$ . Now, since the elements of  $\mathcal{D}_{P_0}(A)$  are not assumed to satisfy the *DC-Property*,  $\mathcal{D}_{P_0}(A)$  may not be a Banach algebra. So we just have to consider the norm completion  $\mathcal{D}_P(A)$  of  $\mathcal{D}_{P_0}(A)$  to have our needed Banach algebra. The notation  $\mathcal{D}_P(A)$ , for the algebra just constructed, is motivated by the fact that we may think of its elements as bounded *Double Pre-centralizers* of  $A$ .

Proposition 2.8 below follows from the above comments and the previous results.

**Proposition 2.8.**  *$\mathcal{W}(A)$  and  $\mathcal{W}(\mathcal{A}_R)$  are Banach subalgebras of the Banach algebra  $\mathcal{D}_P(A)$ , with nonvoid intersection, since  $\langle L, R \rangle \in \mathcal{W}(A) \cap \mathcal{W}(\mathcal{A}_R)$ . Moreover,  $\langle T, S \rangle \in \mathcal{W}(\mathcal{A}_R)$  if and only if  $\langle LT, RS \rangle \in \mathcal{W}(A)$ .  $\square$*

N.B. Throughout the remainder of the sequel, and unless otherwise stated,  $A$  will always be a Banach algebra without annihilators and  $\mathcal{A}_R$  will be the Banach algebra in Proposition 2.1, induced by some double centralizer  $\langle L, R \rangle \in \mathcal{W}(A)$ .

Given  $a \in A$ , let us set

$$(12) \quad \lambda_a^R(x) = R(a) x = \lambda_{R(a)}(x) = [(\lambda R)(a)](x) \quad , \quad \forall x \in A \quad ,$$

$$(13) \quad \mu_a^R(x) = R(x) a = x L(a) = [(\mu L)(a)](x) \quad , \quad \forall x \in A \quad ,$$

and let  $u_R$  be defined by  $u_R(a) = \langle \lambda_a^R, \mu_a^R \rangle, \forall a \in \mathcal{A}_R$ . Then we have:

**Proposition 2.9.** *Let  $\langle L, R \rangle \in \mathcal{W}(A)$  define  $\mathcal{W}(\mathcal{A}_R)$ . Then*

(i)  $u_R$  is the regular homomorphism of  $\mathcal{A}_R$  into  $\mathcal{W}(\mathcal{A}_R)$ , and if there exists  $z \in A$  such that  $L(z) = R(z)$ , we have in the algebra  $\mathcal{D}_P(A)$ :

$$(14) \quad \left\{ \langle \lambda_z, \mu_z \rangle, \langle \lambda_z^R, \mu_z^R \rangle \right\} \subset \mathcal{W}(A) \cap \mathcal{W}(\mathcal{A}_R).$$

(ii) If  $\langle L, R \rangle$  is a multiplier ( $T = L = R$ ), then  $\mathcal{W}(A)$  is a Banach subalgebra of  $\mathcal{W}(\mathcal{A}_R)$ .

*Proof.* (i) Short computations using the definition of the product in  $\mathcal{W}(\mathcal{A}_R)$  show that  $a \mapsto u_R(a) = \langle \lambda_a^R, \mu_a^R \rangle$  is a homomorphism into  $\mathcal{W}(\mathcal{A}_R)$  and that  $\lambda_a^R$  and  $\mu_a^R$  satisfy the DC-Property in  $\mathcal{W}(\mathcal{A}_R)$ . The regular homomorphism  $u : a \mapsto \langle \lambda_a, \mu_a \rangle$ , being from  $A$  into  $\mathcal{W}(A)$ ,  $\langle \lambda_z, \mu_z \rangle$  and  $\langle \lambda_{R(z)}, \mu_{R(z)} \rangle$ , belong to  $\mathcal{W}(A)$ , for each  $z \in A$ . Let us show that these two elements belong to  $\mathcal{W}(\mathcal{A}_R)$ , whenever  $z \in A$  fulfills  $L(z) = R(z)$ . Indeed, for each  $a \in A$ , we have

$$(15) \quad x \cdot \lambda_a(y) = R(x) a y = x L(a) y \quad , \quad \forall x, y \in A,$$

$$(16) \quad \mu_a(x) \cdot y = R(x a) y = x R(a) y \quad , \quad \forall x, y \in A.$$

Hence, if  $z \in A$  satisfies  $L(z) = R(z)$ , and since  $A$  is without annihilators, relations (15) and (16) above yield  $\langle \lambda_z, \mu_z \rangle \in \mathcal{W}(\mathcal{A}_R)$ . For  $z$  with the above property, we have

$$\begin{aligned} x \cdot \lambda_{R(z)}(y) &= R(x) R(z) y = x L R(z) y = x R L(z) y \quad (L \text{ and } R \text{ commute}) \\ &= R(x L(z)) y = (x L(z)) \cdot y = (x R(z)) \cdot y \quad (L(z) = R(z)) \\ &= \mu_{R(z)}(x) \cdot y \quad , \end{aligned}$$

that is,  $\langle \lambda_{R(z)}, \mu_{R(z)} \rangle = \langle \lambda_z^R, \mu_z^R \rangle \in \mathcal{W}(\mathcal{A}_R)$ , and the desired result,  $\langle \lambda_z^R, \mu_z^R \rangle \in \mathcal{W}(\mathcal{A}_R) \cap \mathcal{W}(A)$ , hence follows.

(ii) Let  $\langle L, R \rangle \in \mathcal{W}(A)$  be such that  $L = R$ , so that  $R$  satisfies  $aR(b) = R(a)b$  for all  $a, b \in A$ . Then given any  $\langle T, S \rangle \in \mathcal{W}(A)$ , we have

$$\begin{aligned} a \cdot T(b) &= R(a) T(b) = a R(T(b)) = R(a T(b)) \\ &= R(S(a) b) = R(S(a)) b = S(a) \cdot b \quad , \end{aligned}$$

which proves that  $\langle T, S \rangle \in \mathcal{W}(\mathcal{A}_R)$ . □

*Remark 2.10.* There exists a homomorphism  $\varphi$  from  $\mathcal{W}(\mathcal{A}_R)$  into  $\mathcal{W}(A)$  which makes the following diagram commutative:

$$\begin{array}{ccc} R : \mathcal{A}_R & \longrightarrow & \text{Im}(R) \subset A \\ u_R \downarrow & & \downarrow u \\ \varphi : \mathcal{W}(\mathcal{A}_R) & \longrightarrow & \mathcal{W}(A) \end{array}$$

where  $u$  and  $u_R$  denote respectively the regular homomorphisms of  $A$  into  $\mathcal{W}(A)$  and of  $\mathcal{A}_R$  into  $\mathcal{W}(\mathcal{A}_R)$ . Indeed, the homomorphisms  $uR$  and  $u_R$  are defined from  $\mathcal{A}_R$ , respectively into  $\mathcal{W}(A)$  and  $\mathcal{W}(\mathcal{A}_R)$ , and satisfy  $\text{Ker}(u_R) = \{0\} \subset \text{Ker}(uR)$ . Then there exists a map  $\varphi : \mathcal{W}(\mathcal{A}_R) \rightarrow \mathcal{W}(A)$  such that we have  $\varphi u_R = uR$ . Moreover,  $\varphi$  is clearly a homomorphism from  $\mathcal{W}(\mathcal{A}_R)$  into  $\mathcal{W}(A)$ .  $\square$

Under the hypotheses and notations of Remark 2.10, short computations lead to the following relations, for each  $a \in A$ , and each  $\langle T, S \rangle \in \mathcal{W}(\mathcal{A}_R)$ :

$$(17) \quad \langle T, S \rangle u_R(a) = \langle \lambda_{RT(a)}, \mu_{LT(a)} \rangle = \langle \lambda_{T(a)}^R, \mu_{T(a)}^R \rangle = u_R(T(a)),$$

$$(18) \quad u_R(a) \langle T, S \rangle = \langle \lambda_{RS(a)}, \mu_{LS(a)} \rangle = \langle \lambda_{S(a)}^R, \mu_{S(a)}^R \rangle = u_R(S(a)).$$

Relations (17) and (18) show that for each  $a \in A$ ,  $\varphi$  is still defined on  $\mathcal{W}(\mathcal{A}_R) u_R(a)$  and on  $u_R(a) \mathcal{W}(\mathcal{A}_R)$  as follows:

$$(19) \quad \varphi(\langle T, S \rangle u_R(a)) = \varphi(u_R(T(a))) = u[RT(a)] = \langle \lambda_{RT(a)}, \mu_{RT(a)} \rangle,$$

$$(20) \quad \varphi(u_R(a) \langle T, S \rangle) = \varphi(u_R(S(a))) = u[RS(a)] = \langle \lambda_{RS(a)}, \mu_{RS(a)} \rangle.$$

The next result gives the conditions under which  $\varphi$  extends to the whole space  $\mathcal{W}(\mathcal{A}_R)$ .

**Theorem 2.11.** *If for each  $\langle T, S \rangle \in \mathcal{W}(\mathcal{A}_R)$  there exists  $\langle P, Q \rangle \in \mathcal{W}(A)$  such that the system below is satisfied:*

$$(*) \quad \begin{cases} P R = R T, \\ Q R = R S. \end{cases}$$

*Then the map  $\Theta : \mathcal{W}(\mathcal{A}_R) \rightarrow \mathcal{W}(A)$ , defined by  $\Theta(\langle T, S \rangle) = \langle P, Q \rangle$ , extends the homomorphism  $\varphi : \mathcal{W}(\mathcal{A}_R) \rightarrow \mathcal{W}(A)$ . Furthermore,  $\Theta$  is an isomorphism, provided  $R$  is one-to-one.*

*Proof.* If  $\varphi$  extends to the whole of  $\mathcal{W}(\mathcal{A}_R)$ , then for each  $a \in A$  and each  $\langle T, S \rangle \in \mathcal{W}(\mathcal{A}_R)$ , the element  $\langle P, Q \rangle = \varphi(\langle T, S \rangle)$  would satisfy, according to (19) and (20),

$$\begin{aligned} \varphi(\langle T, S \rangle u_R(a)) &= \langle P, Q \rangle \varphi(u_R(a)) = \langle P, Q \rangle \langle \lambda_{R(a)}, \mu_{R(a)} \rangle \\ &= \langle \lambda_{RT(a)}, \mu_{RT(a)} \rangle, \\ \varphi(u_R(a) \langle T, S \rangle) &= \varphi(u_R(a)) \langle P, Q \rangle = \langle \lambda_{R(a)}, \mu_{R(a)} \rangle \langle P, Q \rangle \\ &= \langle \lambda_{RS(a)}, \mu_{RS(a)} \rangle. \end{aligned}$$

But we have

$$(21) \quad \langle P, Q \rangle \langle \lambda_{R(a)}, \mu_{R(a)} \rangle = \langle P \lambda_{R(a)}, \mu_{R(a)} Q \rangle = \langle \lambda_{PR(a)}, \mu_{PR(a)} \rangle,$$

$$(22) \quad \langle \lambda_{R(a)}, \mu_{R(a)} \rangle \langle P, Q \rangle = \langle \lambda_{R(a)} P, Q \mu_{R(a)} \rangle = \langle \lambda_{QR(a)}, \mu_{QR(a)} \rangle.$$

Hence, if  $\varphi$  extends to  $\mathcal{W}(\mathcal{A}_R)$ , then we must have

$$\begin{cases} \langle \lambda_{RT(a)}, \mu_{RT(a)} \rangle = \langle \lambda_{PR(a)}, \mu_{PR(a)} \rangle, \\ \langle \lambda_{RS(a)}, \mu_{RS(a)} \rangle = \langle \lambda_{QR(a)}, \mu_{QR(a)} \rangle, \end{cases}$$

from which the system (\*) follows. So, if for each  $\langle T, S \rangle \in \mathcal{W}(\mathcal{A}_R)$  there exists  $\langle P, Q \rangle \in \mathcal{W}(A)$  such that the system (\*) is satisfied, we define the map  $\Theta : \mathcal{W}(\mathcal{A}_R) \rightarrow \mathcal{W}(A)$  by

$$(23) \quad \Theta(\langle T, S \rangle) = \langle P, Q \rangle.$$

The restriction of  $\Theta$  coincides with  $\varphi$  on  $u_R(\mathcal{A}_R)$  since for each  $u_R(a) = \langle \lambda_{R(a)}, \mu_{R(a)} \rangle$ , the element  $\langle P, Q \rangle = \langle \lambda_{R(a)}, \mu_{R(a)} \rangle \in \mathcal{W}(A)$  clearly fulfills the system (\*).

We need to show that  $\Theta$  as defined above is a homomorphism. So let  $\langle T, S \rangle$  and  $\langle T', S' \rangle$  belong to  $\mathcal{W}(\mathcal{A}_R)$ , and let  $\langle P, Q \rangle$  and  $\langle P', Q' \rangle$  be elements of  $\mathcal{W}(A)$  satisfying

$$(24) \quad \Theta(\langle T, S \rangle) = \langle P, Q \rangle \quad \text{and} \quad \Theta(\langle T', S' \rangle) = \langle P', Q' \rangle.$$

Then  $\langle T T', S' S \rangle \in \mathcal{W}(\mathcal{A}_R)$ ,  $\langle P P', Q' Q \rangle \in \mathcal{W}(A)$ , and we have

$$\begin{aligned} (P P') R &= P (P' R) = P (R T') = (P R) T' = R (T T'), \\ (Q' Q) R &= Q' (Q R) = Q' (R S) = (Q' R) S = R (S' S). \end{aligned}$$

So,  $\langle P P', Q' Q \rangle \in \mathcal{W}(A)$  fulfills the system (\*). Therefore, by definition of  $\Theta$ , we get

$$\begin{aligned} \Theta(\langle T T', S' S \rangle) &= \langle P P', Q' Q \rangle = \langle P, Q \rangle \langle P', Q' \rangle \\ &= \Theta(\langle T, S \rangle) \Theta(\langle T', S' \rangle) \end{aligned}$$

and  $\Theta$  is hence a homomorphism from  $\mathcal{W}(\mathcal{A}_R)$  into  $\mathcal{W}(A)$ , as desired.

Now, if  $\langle T, S \rangle \in \mathcal{W}(\mathcal{A}_R)$  is such that  $\langle P, Q \rangle = \Theta(\langle T, S \rangle) = 0$ , then by virtue of the system (\*), the following holds:  $Im(T) \cup Im(S) \subset Ker(R)$ , and if  $R$  is a one-to-one map, then  $T$  and  $S$  must be identically null.  $\square$

We next deal with the uniqueness of the above extension of the homomorphism  $\varphi$ .

**Theorem 2.12.** *Let us assume that there exists another extension  $\Sigma$  of the homomorphism  $\varphi$  to the entire  $\mathcal{W}(\mathcal{A}_R)$ . Then,  $\Sigma$  “fulfills” the system (\*), and in the particular case where  $R$  is bijective, there exists a unique way of extending  $\varphi$  to  $\mathcal{W}(\mathcal{A}_R)$ .*

*Proof.*  $\Sigma$  must coincide with  $\varphi$  on the image  $u_R(\mathcal{A}_R)$  of the regular homomorphism from  $\mathcal{A}_R$  into  $\mathcal{W}(\mathcal{A}_R)$ , that is,

$$(25) \quad \Sigma(u_R(a)) = \Sigma(\langle \lambda_{R(a)}, \mu_{L(a)} \rangle) = \langle \lambda_{R(a)}, \mu_{R(a)} \rangle,$$

and since  $u_R(A)$  is a two-sided ideal in  $\mathcal{W}(\mathcal{A}_R)$ ,  $\Sigma$  must also satisfy

$$(26) \quad \Sigma(\langle T, S \rangle u_R(a)) = \Sigma(\langle \lambda_{RT(a)}, \mu_{LT(a)} \rangle) = \langle \lambda_{RT(a)}, \mu_{RT(a)} \rangle,$$

$$(27) \quad \Sigma(u_R(a) \langle T, S \rangle) = \Sigma(\langle \lambda_{RS(a)}, \mu_{LS(a)} \rangle) = \langle \lambda_{RS(a)}, \mu_{RS(a)} \rangle.$$

So, if  $\Sigma(\langle T, S \rangle) = \langle U, V \rangle$ , then we must have

$$\begin{aligned} \Sigma(\langle T, S \rangle u_R(a)) &= \langle U, V \rangle \langle \lambda_{R(a)}, \mu_{R(a)} \rangle = \langle \lambda_{UR(a)}, \mu_{UR(a)} \rangle \\ &= \langle \lambda_{RT(a)}, \mu_{RT(a)} \rangle, \\ \Sigma(u_R(a) \langle T, S \rangle) &= \langle \lambda_{R(a)}, \mu_{R(a)} \rangle \langle U, V \rangle = \langle \lambda_{VR(a)}, \mu_{VR(a)} \rangle \\ &= \langle \lambda_{RS(a)}, \mu_{RS(a)} \rangle. \end{aligned}$$

The above relations show that  $\Sigma(\langle T, S \rangle) = \langle U, V \rangle$  satisfies the system (\*) as well, and therefore the following system (\*\*) is also fulfilled by  $\langle P, Q \rangle$  and  $\langle U, V \rangle$ :

$$(**) \quad \begin{cases} P R = U R, \\ Q R = V R. \end{cases}$$

But the conditions in (\*\*) above may also be interpreted in the following way, in terms of restrictions of maps:  $P|_{Im(R)} = U|_{Im(R)}$  and  $Q|_{Im(R)} = V|_{Im(R)}$ , which, in the case where  $R$  is bijective (and hence onto  $A$ ), gives

$$(28) \quad \Theta(\langle T, S \rangle) = \langle P, Q \rangle = \langle U, V \rangle = \Sigma(\langle T, S \rangle),$$

that is,  $\Theta = \Sigma$ , which proves the uniqueness of the extension of  $\varphi$ . In such a case where the map  $R$  is bijective,  $R$  admits a well-defined inverse  $R^{-1} \in B(A)$  which is a right centralizer, and therefore, the system (\*) gives, for  $\langle P, Q \rangle = \Theta(\langle T, S \rangle)$ ,

$$(29) \quad P = R T R^{-1} \quad \text{and} \quad Q = R S R^{-1},$$

which completes the proof.  $\square$

Let us denote by  $W^R(A)$ , the normed algebra of bounded double centralizers of  $Im(R) \subseteq A$ . Then it is clear that  $\mathcal{W}(A) \subseteq W^R(A)$ , and that equality holds whenever  $R$  is onto  $A$ . The next result links  $W^R(A)$  with the space of elements  $P, Q \in B(A)$  which satisfy the system (\*) with the elements of  $\mathcal{W}(\mathcal{A}_R)$ .

**Proposition 2.13.** *Let  $P, Q \in B(A)$  satisfy the system (\*) with some  $\langle T, S \rangle \in \mathcal{W}(\mathcal{A}_R)$ . Then  $\langle P, Q \rangle \in W^R(A)$ .*

*Proof.* For all  $a, b \in Im(R)$ , we have  $\alpha, \beta \in A$ , such that  $a = R(\alpha)$ ;  $b = R(\beta)$ , and then, for each  $\langle P, Q \rangle \in B(A)$  satisfying the system (\*) with some  $\langle T, S \rangle \in \mathcal{W}(\mathcal{A}_R)$ , the following holds:

$$\begin{aligned} a P(b) &= R(\alpha) (P R)(\beta) = R(\alpha) (R T)(\beta) \quad (\text{since } (*) \text{ holds}) \\ &= R(R(\alpha) T(\beta)) \\ &= R(\alpha \cdot T(\beta)) = R(S(\alpha) \cdot \beta) \quad (\text{since } \langle T, S \rangle \in \mathcal{W}(\mathcal{A}_R)) \\ &= R(R S(\alpha) \beta) \\ &= R S(\alpha) R(\beta) = Q R(\alpha) R(\beta) \quad (\text{since } (*) \text{ holds}) \\ &= Q(a) b, \end{aligned}$$

which is the *DC-Property* in  $W^R(A)$ , and the proof is complete.  $\square$

As is well known, the theory of double centralizers is a helpful device in the study of extensions of algebras. Let us recall the following definition (cf. [4], pp. 33 - 34).

**Definition 2.14.** Let  $A$  and  $B$  be Banach algebras. An *extension of  $A$  by  $C$*  is a short exact sequence  $0 \rightarrow A \xrightarrow{\rho} B \xrightarrow{\psi} C \rightarrow 0$  of Banach algebras. The extension is called a *semidirect product* if there is a continuous homomorphism  $\chi : C \rightarrow B$ , such that  $\psi \chi$  is the identity map on  $C$ . The short exact sequence is then said to *split*, with splitting homomorphism  $\chi$ .

**Theorem 2.15.** Let  $A$  be a Banach algebra with a left approximate identity bounded by one. Let  $\mathcal{A}_R$  be induced by  $\langle L, R \rangle \in \mathcal{W}(A)$  with  $R$  bijective. Then there exists an extension of  $A$  by  $\mathcal{W}(\mathcal{A}_R)$ :

$$(30) \quad 0 \rightarrow A \xrightarrow{\rho} B \xrightarrow{\psi} \mathcal{W}(\mathcal{A}_R) \rightarrow 0,$$

and a continuous homomorphism  $\Theta : B \rightarrow \mathcal{W}(A)$ , such that the following diagram commutes:

$$\begin{array}{ccccccc} 0 & \longrightarrow & A & \xrightarrow{u} & \mathcal{W}(A) & \xrightarrow{\pi} & \mathcal{W}(A)/A \longrightarrow 0 \\ & & \uparrow I_A & & \uparrow \Theta & & \uparrow \tau \\ 0 & \longrightarrow & A & \xrightarrow{\rho} & B & \xrightarrow{\psi} & \mathcal{W}(\mathcal{A}_R) \longrightarrow 0. \end{array}$$

This extension, which is unique up to equivalence, is moreover a semidirect product.

*Proof.* Identifying  $A$  with  $u(A)$  under the regular homomorphism  $u$  of  $A$  into  $\mathcal{W}(A)$ , consider the short exact sequence  $0 \rightarrow A \xrightarrow{u} \mathcal{W}(A) \xrightarrow{\pi} C \rightarrow 0$ , where  $\pi$  is the natural map. Since  $R$  is bijective, Theorem 2.12 asserts that  $\Theta$  is a homeomorphic isomorphism of  $\mathcal{W}(\mathcal{A}_R)$  onto  $\mathcal{W}(A)$ . The map  $\tau : \mathcal{W}(\mathcal{A}_R) \rightarrow \mathcal{W}(A)/A$ , defined by

$$(31) \quad \tau(\langle T, S \rangle) = \Theta(\langle T, S \rangle) + A, \quad \forall \langle T, S \rangle \in \mathcal{W}(\mathcal{A}_R),$$

is hence a continuous homomorphism satisfying:  $\tau = \pi \Theta$ . Then, the hypotheses of Theorem 1.2.10 in [4], page 34, are satisfied, and its conclusion provides the existence of the extension of  $A$  given by relation (30), and unique up to equivalence. Furthermore, in virtue of Theorem 1.2.11 in [4], page 35, the existence of  $\tau$  satisfying  $\tau = \pi \Theta$  shows that the extension is a semidirect product.  $\square$

Now, let us assume that  $A$  is commutative, and let us denote by  $\mathcal{M}(A)$  the algebra of multipliers of  $A$ , which necessarily coincides with  $\mathcal{W}(A)$ . In [3], Corollary 1.3.1, page 23, R. Larsen proved that for a commutative semisimple Banach algebra  $A$ , the following equality holds:  $\mathcal{M}(A) = \mathcal{M}(\mathcal{A}_R)$ . In what follows, we still obtain the same result when the assumption of semisimplicity on  $A$  is removed. We shall need the following:

**Lemma 2.16.** Let  $A$  be a commutative Banach algebra and let  $R \in \mathcal{M}(A)$ . Then the following conditions are equivalent:

- (i)  $\mathcal{A}_R$  has no annihilators;
- (ii)  $A$  has no annihilators and  $R$  is one-to-one.

*Proof.* Assume that (i) is fulfilled, and let  $a \in A$  be given. Then,  $A$  being commutative and  $R$  being a multiplier of  $A$ , we get for all  $x \in A$ :

$$(32) \quad R(x a) = R(a x) = R(a) x = a \cdot x = x \cdot a \quad (\mathcal{A}_R \text{ is commutative}),$$

which, since  $\mathcal{A}_R$  is without annihilators, implies that  $A$  has no annihilators. Moreover, due to Corollary 2.3,  $R$  is one-to-one, if  $\mathcal{A}_R$  has no annihilators. Hence (i) implies (ii).

Conversely if (ii) is satisfied, then Corollary 2.3 again implies that  $\mathcal{A}_R$  has no annihilators, and the equivalence of (i) and (ii) is therefore established.  $\square$

We can now prove:

**Theorem 2.17.** *Let  $A$  be a commutative Banach algebra and let  $R \in \mathcal{M}(A)$ . Then  $\mathcal{M}(A) = \mathcal{M}(\mathcal{A}_R)$  if and only if one of the equivalent conditions in Lemma 2.16 is satisfied.*

*Proof.* We assume that condition (i) of Lemma 2.16 holds. Let  $T \in \mathcal{M}(A)$ ; then for all  $a, b \in A$ , we get  $T(a \cdot b) = T[R(a) b] = R(a) T(b) = a \cdot T(b)$  and

$$\begin{aligned} T(a \cdot b) &= R(a) T(b) = T(b) R(a) = R[T(b) a] = R[b T(a)] \\ &= R(T(a) b) = R[T(a)] b = T(a) \cdot b, \end{aligned}$$

which means that  $T \in \mathcal{M}(A)$ , and consequently,  $\mathcal{M}(A) \subseteq \mathcal{M}(\mathcal{A}_R)$ .

Conversely, let  $T \in \mathcal{M}(\mathcal{A}_R)$ . Then, for all  $a, b, x \in A$ , we get on one hand

$$\begin{aligned} R(x) T(a b) &= x \cdot T(a b) = T(x) \cdot a b = a b T(x) = R(ab) T(x) \\ &= [R(a) b] T(x) = R(a) T(x) b = [a \cdot T(x)] b = [T(a) \cdot x] b \\ &= (R[T(a)] x) = T(a) R(x) b = R(x)[T(a) b], \end{aligned}$$

which implies, since  $\mathcal{A}_R$  is without annihilators,  $T(a b) = T(a) b$ .

On the other hand, let  $a, b, x \in A$ ; then

$$R(x) T(a b) = R(a b) T(x) = a [b \cdot T(x)] = a [T(b) R(x)] = R(x) [a T(b)],$$

and once again, the fact that  $\mathcal{A}_R$  is without annihilators, leads to  $T(a b) = a T(b)$ . Therefore,  $T \in \mathcal{M}(A)$ , and this provides the reverse inclusion  $\mathcal{M}(\mathcal{A}_R) \subseteq \mathcal{M}(A)$ , and hence the desired equality  $\mathcal{M}(\mathcal{A}_R) = \mathcal{M}(A)$ .  $\square$

The following result also appears in [3], Corollary 1.3.2, page 23:

**Proposition 2.18.** *Let  $A$  be a commutative semisimple Banach algebra which admits factorization, i.e.: For every  $a \in A$ , there exist  $x, y \in A$ , such that  $a = xy$ . Then the following conditions are equivalent:*

- (i)  $A_T$  admits factorization;
- (ii)  $T$  is invertible.

Dropping the assumption of semisimplicity on  $A$ , we can prove the following:

**Theorem 2.19.** *Let  $A$  be a commutative Banach algebra which admits factorization, and let  $R \in \mathcal{M}(A)$  be such that  $\mathcal{A}_R$  is without annihilators. Then  $\mathcal{A}_R$  admits factorization, if and only if  $R$  is invertible.*

*Proof.* Assume that  $\mathcal{A}_R$  admits factorization. Then, since  $\mathcal{A}_R$  is without annihilators,  $R$  is one-to-one, according to Lemma 2.16. Moreover, since  $\mathcal{A}_R$  admits factorization, for each  $a \in \mathcal{A}_R$  there exist  $\alpha, \beta \in \mathcal{A}_R$ , such that  $a = \alpha \cdot \beta = R(\alpha) \beta = R(\alpha \beta)$ , which means that  $R$  is onto  $A$ , whence  $R$  is bijective and  $R^{-1}$  exists.

Conversely, let  $R$  be invertible; then  $R$  is one-to-one, and since it is also onto  $A$ , we get for each  $a \in A$  some  $\alpha \in \mathcal{A}_R$  satisfying  $a = R(\alpha)$ . But,  $A$  admits factorization, and therefore  $\alpha$  has the following decomposition:  $\alpha = u v$ ,  $u, v \in A$ .

Hence we get  $a = R(\alpha) = R(uv) = R(u)v = u \cdot v$ , so that  $\mathcal{A}_R$  also admits factorization, and the proof is complete.  $\square$

Coming back to the general case where  $A$  is neither commutative nor semisimple, we give the analogous version of Theorem 2.17, when the multiplier algebra is replaced by the double centralizer algebra, as follows:

**Theorem 2.20.** *Let  $A$  be without annihilators, and let  $\langle L, R \rangle \in \mathcal{W}(A)$  define  $\mathcal{A}_R$  and be such that  $R$  is bijective. Then  $\mathcal{W}(\mathcal{A}_R)$  and  $\mathcal{W}(A)$  are homeomorphically isomorphic.*

*Proof.* According to Theorems 2.11 and 2.12, there exists a homomorphism  $\Theta$  which extends in a unique way the homomorphism  $\varphi$  constructed in Remark 2.10 to the algebra  $\mathcal{W}(\mathcal{A}_R)$ . Moreover, according to Theorem 2.11, each element  $\langle T, S \rangle \in \text{Ker}(\Theta)$  satisfies  $\text{Im}(T) \cup \text{Im}(S) \subset \text{Ker}(R)$ . So, if  $R$  is bijective and hence one-to-one,  $\text{Ker}(R) = \{0\}$ , and  $\Theta$  is also one-to-one. Therefore,  $\Theta$  is a bijection of  $\mathcal{W}(\mathcal{A}_R)$  onto  $\mathcal{W}(A)$ . But, according to relation (29), the positive constant  $K_R = \|R^{-1}\|\|R\|$  (depending only on  $R$ ) satisfies  $\|\Theta(\langle T, S \rangle)\| \leq K_R \|\langle T, S \rangle\|$ ;  $\forall \langle T, S \rangle \in \mathcal{W}(\mathcal{A}_R)$ , so that  $\Theta$  is a bounded linear map from  $\mathcal{W}(\mathcal{A}_R)$  onto the Banach space  $\mathcal{W}(A)$ , that is, a homeomorphism, and the proof is complete.  $\square$

#### REFERENCES

1. Desquith, E.: Banach algebras associated to bounded module maps; Preprint ICTP N<sup>o</sup> IC 98/194; (1998).
2. Doran, R.S. & Fell, J.M.G.: Representations of \*-algebras, locally compact groups, and Banach \*-algebraic Bundles, Vol. 2; Academic Press (1988). MR **90c**:46002
3. Larsen, R.: Introduction to the theory of multipliers; Springer-Verlag (1971). MR **55**:8695
4. Palmer, T.W.: Banach algebras and the general theory of Banach\*-algebras, Vol. I: Algebras and Banach algebras; Cambridge University Press (1994). MR **95c**:46002

INSTITUT DE RECHERCHES MATHÉMATIQUES (IRMA), 08 BP 2030 ABIDJAN 08, IVORY COAST  
*E-mail address:* [desquith@hotmail.com](mailto:desquith@hotmail.com)