

TRANSFINITE SEQUENCES OF CONTINUOUS AND BAIRE CLASS 1 FUNCTIONS

MÁRTON ELEKES AND KENNETH KUNEN

(Communicated by Alan Dow)

ABSTRACT. The set of continuous or Baire class 1 functions defined on a metric space X is endowed with the natural pointwise partial order. We investigate how the possible lengths of well-ordered monotone sequences (with respect to this order) depend on the space X .

INTRODUCTION

Any set \mathcal{F} of real valued functions defined on an arbitrary set X is partially ordered by the pointwise order; that is, $f \leq g$ iff $f(x) \leq g(x)$ for all $x \in X$. Then, $f < g$ iff $f \leq g$ and $g \not\leq f$; equivalently, $f(x) \leq g(x)$ for all $x \in X$ and $f(x) < g(x)$ for at least one $x \in X$. Our aim will be to investigate the possible lengths of the increasing or decreasing well-ordered sequences of functions in \mathcal{F} with respect to this order. A classical theorem (see Kuratowski [7], §24.III, Theorem 2') asserts that if \mathcal{F} is the set of Baire class 1 functions (that is, pointwise limits of continuous functions) defined on a Polish space X (that is, a complete separable metric space), then there exists a monotone sequence of length ξ in \mathcal{F} iff $\xi < \omega_1$. P. Komjáth [5] proved that the corresponding question concerning Baire class α functions for $2 \leq \alpha < \omega_1$ is independent of *ZFC*.

In the present paper we investigate what happens if we replace the Polish space X by an arbitrary metric space.

Section 1 considers chains of continuous functions. We show that for any metric space X , there exists a chain in $C(X, \mathbb{R})$ of order type ξ iff $|\xi| \leq d(X)$. Here, $|A|$ denotes the cardinality of the set A , while $d(X)$ denotes the density of the space X , that is,

$$d(X) = \max(\min\{|D| : D \subseteq X \text{ \& } \overline{D} = X\}, \omega).$$

In particular, for separable X , every well-ordered chain has countable length, just as for Polish spaces.

Section 2 considers chains of Baire class 1 functions on separable metric spaces. Here, the situation is entirely different from the case of Polish spaces, since on some

Received by the editors December 11, 2001 and, in revised form, March 13, 2002.
2000 *Mathematics Subject Classification.* Primary 26A21; Secondary 03E17, 54C30.
Key words and phrases. Baire class 1, separable metric space, transfinite sequence of functions.
The second author was partially supported by NSF Grant DMS-0097881.

separable metric spaces there are well-ordered chains of every order type less than ω_2 . Furthermore, the existence of chains of type ω_2 and longer is independent of $ZFC + \neg CH$. Under MA , there are chains of all types less than \mathfrak{c}^+ , whereas in the Cohen model, all chains have type less than ω_2 .

We note here that instead of examining well-ordered sequences, which is a classical problem, we could try to characterize all the possible order types of linearly ordered subsets of the partially ordered set \mathcal{F} . This problem was posed by M. Laczko, and is considered in detail in [3].

1. SEQUENCES OF CONTINUOUS FUNCTIONS

Lemma 1.1. *For any topological space X : If there is a well-ordered sequence of length ξ in $C(X, \mathbb{R})$, then $\xi < d(X)^+$.*

Proof. Let $\{f_\alpha : \alpha < \xi\}$ be an increasing sequence in $C(X, \mathbb{R})$, and let $D \subseteq X$ be a dense subset of X such that $d(X) = \max(|D|, \omega)$. By continuity, the $f_\alpha \upharpoonright D$ are all distinct; so, for each $\alpha < \xi$, choose a $d_\alpha \in D$ such that $f_\alpha(d_\alpha) < f_{\alpha+1}(d_\alpha)$. For each $d \in D$ the set $E_d = \{\alpha : d_\alpha = d\}$ is countable, because every well-ordered subset of \mathbb{R} is countable. Since $\xi = \bigcup_{d \in D} E_d$, we have $|\xi| \leq \max(|D|, \omega) = d(X)$. \square

The converse implication is not true in general. For example, if X has the countable chain condition (ccc), then every well-ordered chain in $C(X, \mathbb{R})$ is countable (because $X \times \mathbb{R}$ is also ccc). However, the converse is true for metric spaces:

Lemma 1.2. *If (X, ρ) is any non-empty metric space and \prec is any total order of the cardinal $d(X)$, then there is a chain in $C(X, \mathbb{R})$ which is isomorphic to \prec .*

Proof. First, note that every countable total order is embeddable in \mathbb{R} , so if $d(X) = \omega$, then the result follows trivially using constant functions. In particular, we may assume that X is infinite, and then fix $D \subseteq X$ which is dense and of size $d(X)$. For each $n \in \omega$, let D_n be a subset of D which is maximal with respect to the property $\forall d, e \in D_n [d \neq e \rightarrow \rho(d, e) \geq 2^{2^{-n}}]$. Then $\bigcup_n D_n$ is also dense, so we may assume that $\bigcup_n D_n = D$. We may also assume that \prec is a total order of the set D . Now, we shall produce $f_d \in C(X, \mathbb{R})$ for $d \in D$ such that $f_d < f_e$ whenever $d \prec e$.

For each n , if $c \in D_n$, define $\varphi_c^n(x) = \max(0, 2^{-n} - \rho(x, c))$. For each $d \in D$, let $\psi_d^n = \sum \{\varphi_c^n : c \in D_n \text{ \& } c \prec d\}$. Since every $x \in X$ has a neighborhood on which all but at most one of the φ_c^n vanish, we have $\psi_d^n \in C(X, [0, 2^{-n}])$, and $\psi_d^n \leq \psi_e^n$ whenever $d \prec e$. Thus, if we let $f_d = \sum_{n < \omega} \psi_d^n$, we have $f_d \in C(X, [0, 2])$, and $f_d \leq f_e$ whenever $d \prec e$. But also, if $d \in D_n$ and $d \prec e$, then $\psi_d^n(d) = 0 < 2^{-n} = \psi_e^n(d)$, so actually $f_d < f_e$ whenever $d \prec e$. \square

Putting these lemmas together, we have:

Theorem 1.3. *Let (X, ρ) be a metric space. Then there exists a well-ordered sequence of length ξ in $C(X, \mathbb{R})$ iff $\xi < d(X)^+$.*

Corollary 1.4. *A metric space (X, ρ) is separable iff every well-ordered sequence in $C(X, \mathbb{R})$ is countable.*

2. SEQUENCES OF BAIRE CLASS 1 FUNCTIONS

If we replace continuous functions by Baire class 1 functions, then Corollary 1.4 becomes false, since on some separable metric spaces we can get well-ordered sequences of every type less than ω_2 . To prove this, we shall apply some basic facts

about \subset^* on $\mathcal{P}(\omega)$. As usual, for $x, y \subseteq \omega$, we say that $x \subseteq^* y$ iff $x \setminus y$ is finite. Then $x \subset^* y$ iff $x \setminus y$ is finite and $y \setminus x$ is infinite. This \subset^* partially orders $\mathcal{P}(\omega)$.

Lemma 2.1. *If $X \subset \mathcal{P}(\omega)$ is a chain in the order \subset^* , then on X (viewed as a subset of the Cantor set $2^\omega \cong \mathcal{P}(\omega)$) there is a chain of Baire class 1 functions which is isomorphic to (X, \subset^*) .*

Proof. Note that for each $x \in X$,

$$\{y \in X : y \subseteq^* x\} = \bigcup_{m \in \omega} \{y \in X : \forall n \geq m [y(n) \leq x(n)]\},$$

which is an F_σ set in X . Likewise, the sets $\{y \in X : y \supseteq^* x\}$, $\{y \in X : y \subset^* x\}$, and $\{y \in X : y \supset^* x\}$, are all F_σ sets in X , and hence also G_δ sets. It follows that if $f_x : X \rightarrow \{0, 1\}$ is the characteristic function of $\{y \in X : y \subset^* x\}$, then $f_x : X \rightarrow \mathbb{R}$ is a Baire class 1 function. Then $\{f_x : x \in X\}$ is the required chain. \square

Lemma 2.2. *For any infinite cardinal κ , suppose that $(\mathcal{P}(\omega), \subset^*)$ contains a chain $\{x_\alpha : \alpha < \kappa\}$ (i.e., $\alpha < \beta \rightarrow x_\alpha \subset^* x_\beta$). Then $(\mathcal{P}(\omega), \subset^*)$ contains a chain X of size κ such that every ordinal $\xi < \kappa^+$ is embeddable into X .*

Proof. Let $S = \bigcup_{1 \leq n < \omega} \kappa^n$. For $s = (\alpha_1, \dots, \alpha_{n-1}, \alpha_n) \in S$, let

$$s^+ = (\alpha_1, \dots, \alpha_{n-1}, \alpha_n + 1).$$

Starting with the $x_{(\alpha)} = x_\alpha$, choose $x_s \in \mathcal{P}(\omega)$ by induction on $\text{length}(s)$ so that

$$x_s = x_{s \smallfrown 0} \subset^* x_{s \smallfrown \alpha} \subset^* x_{s \smallfrown \beta} \subset^* x_{s^+}$$

whenever $s \in S$ and $0 < \alpha < \beta < \kappa$. Let $X = \{x_s : s \in S\}$. Then, whenever $x, y \in X$ with $x \subset^* y$, the ordinal κ is embeddable in $(x, y) = \{z \in X : x \subset^* z \subset^* y\}$. From this, one easily proves by induction on $\xi < \kappa^+$ (using $\text{cf}(\xi) \leq \kappa$) that ξ is embeddable in each such interval (x, y) . \square

Since $\mathcal{P}(\omega)$ certainly contains a chain of type ω_1 , these two lemmas yield:

Theorem 2.3. *There is a separable metric space X on which, for every $\xi < \omega_2$, there is a well-ordered chain of length ξ of Baire class 1 functions.*

Under CH , this is best possible, since there will be only $2^\omega = \omega_1$ Baire class 1 functions on a separable metric space, so there could not be a chain of length ω_2 . Under $\neg CH$, the existence of longer chains of Baire class 1 functions depends on the model of set theory. It is consistent with $\mathfrak{c} = 2^\omega$ being arbitrarily large that there is a chain in $(\mathcal{P}(\omega), \subset^*)$ of type \mathfrak{c} ; for example, this is true under MA (see [2]). In this case, there will be a separable X with well-ordered chains of all lengths less than \mathfrak{c}^+ . However, in the Cohen model, where \mathfrak{c} can also be made arbitrarily large, we never get chains of type ω_2 . We shall prove this by using the following lemma, which relates it to the rectangle problem:

Lemma 2.4. *Suppose that there is a separable metric space Y with an ω_2 -chain of Borel subsets, $\{B_\alpha : \alpha < \omega_2\}$ (so, $\alpha < \beta \rightarrow B_\alpha \subsetneq B_\beta$). Then in $\omega_2 \times \omega_2$, the well-order relation $<$ is in the σ -algebra generated by the set of all rectangles, $\{S \times T : S, T \in \mathcal{P}(\omega_2)\}$.*

Proof. Each B_α has some countable Borel rank. Since there are only ω_1 ranks, we may, by passing to a subsequence, assume that the ranks are bounded. Say each B_α is a Σ_μ^0 set for some fixed $\mu < \omega_1$.

Let $J = \omega^\omega$, and let $A \subseteq Y \times J$ be a universal Σ_μ^0 set; that is, A is Σ_μ^0 in $Y \times J$ and every Σ_μ^0 subset of Y is of the form $A^j = \{y : (y, j) \in A\}$ for some $j \in J$ (see [7], §31). Now, for $\alpha, \beta < \omega_2$, fix $y_\alpha \in B_{\alpha+1} \setminus B_\alpha$, and fix $j_\beta \in J$ such that $A^{j_\beta} = B_\beta$. Then $\alpha < \beta$ iff $(y_\alpha, j_\beta) \in A$. Thus, $\{(y_\alpha, j_\beta) : \alpha < \beta < \omega_2\}$ is a Borel subset of $\{y_\alpha : \alpha < \omega_2\} \times \{j_\beta : \beta < \omega_2\}$, and is hence in the σ -algebra generated by open rectangles, so $<$, as a subset of $\omega_2 \times \omega_2$, is in the σ -algebra generated by rectangles. \square

Theorem 2.5. *Assume that the well-order relation $<$ on ω_2 is not in the σ -algebra generated by the set of all rectangles. Then no separable metric space can have a chain of length ω_2 of Baire class 1 functions.*

Proof. Suppose that $\{f_\alpha : \alpha < \omega_2\}$ is a chain of Baire class one functions on the separable metric space X . Let $B_\alpha = \{(x, r) \in X \times \mathbb{R} : r \leq f_\alpha(x)\}$. Then the B_α form an ω_2 -chain of Borel subsets of the separable metric space $X \times \mathbb{R}$, so we have a contradiction by Lemma 2.4. \square

Finally, we point out that the hypothesis of this theorem is consistent, since it holds in the extension $V[G]$ formed by adding $\geq \omega_2$ Cohen reals to a ground model V which satisfies *CH*. This fact was first proved in [6]. It also follows from the more general principle $HP_2(\omega_2)$ of Brendle, Fuchino, and Soukup [1]. They define this principle, prove that it holds in Cohen extensions (and in a number of other forcing extensions), and show the following:

Lemma 2.6. *$HP_2(\kappa)$ implies that if R is any relation on $\mathcal{P}(\omega)$ which is first-order definable over $H(\omega_1)$ from a fixed element of $H(\omega_1)$, then there is no $X \subseteq \mathcal{P}(\omega)$ such that $(X; R)$ is isomorphic to $(\kappa; <)$.*

These matters are also discussed in [4], which indicates how such statements are verified in Cohen extensions. Here, $H(\omega_1)$ denotes the set of hereditarily countable sets.

Lemma 2.7. *$HP_2(\omega_2)$ implies that in $\omega_2 \times \omega_2$, the well-order relation $<$ is not in the σ -algebra generated by the set of all rectangles, $\{S \times T : S, T \in \mathcal{P}(\omega_2)\}$.*

Proof. Suppose that $<$ were in this σ -algebra. Then we would have fixed $K_n \subseteq \omega_2$ for $n < \omega$ such that $<$ is in the σ -algebra generated by all the $K_m \times K_n$. For each α , let $u_\alpha = \{n \in \omega : \alpha \in K_n\}$. There is then a formula $\varphi(x, y, z)$ and a fixed $w \in H(\omega_1)$ such that for all $\alpha, \beta < \omega_2$, $\alpha < \beta$ iff $H(\omega_1) \models \varphi(u_\alpha, u_\beta, w)$; here, w encodes the particular countable boolean combination used to get $<$ from the K_n . Now, if $X = \{u_\alpha : \alpha < \omega_2\}$, then φ defines a relation R on $H(\omega_1)$ such that $(X; R)$ is isomorphic to $(\omega_2; <)$, contradicting Lemma 2.6. \square

REFERENCES

- [1] J. Brendle, S. Fuchino, and L. Soukup, Coloring ordinals by reals, to appear.
- [2] E. K. van Douwen, The integers and topology, in *Handbook of Set-Theoretic Topology*, North-Holland, Amsterdam, 1984, pp. 111-167. MR **87f**:54008
- [3] M. Elekes, Linearly ordered families of Baire 1 functions, *Real Analysis Exchange*, **27** (2001/02), 49–63.

- [4] I. Juhász and K. Kunen, The power set of ω , *Fundamenta Mathematicae*, Vol **170** (2001), 257-265.
- [5] P. Komjáth, Ordered families of Baire-2-functions, *Real Analysis Exchange*, Vol **15** (1989-90), 442-444. MR **91d**:26007
- [6] K. Kunen, *Inaccessibility Properties of Cardinals*, Doctoral Dissertation, Stanford, 1968.
- [7] K. Kuratowski, *Topology, Vol. 1*, Academic Press, 1966. MR **36**:840

DEPARTMENT OF ANALYSIS, EÖTVÖS LORÁND UNIVERSITY, BUDAPEST, PÁZMÁNY PÉTER SÉTÁNY
1/C, 1117, HUNGARY

E-mail address: `emarci@cs.elte.hu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN, MADISON, WISCONSIN 53706

E-mail address: `kunen@math.wisc.edu`

URL: `http://www.math.wisc.edu/~kunen`