

BELL REPRESENTATIONS OF FINITELY CONNECTED PLANAR DOMAINS

MOONJA JEONG AND MASAHIKO TANIGUCHI

(Communicated by Mei-Chi Shaw)

ABSTRACT. In this paper, we solve a conjecture of S. Bell (1992) affirmatively. Actually, we prove that every non-degenerate n -connected planar domain Ω , where $n > 1$ is representable as $\Omega = \{|f| < 1\}$ with a suitable rational function f of degree n . This result is considered as a natural generalization of the classical Riemann mapping theorem for simply connected planar domains.

1. INTRODUCTION AND THE MAIN THEOREM

Recently, S. Bell posed the following problem ([B1] and [B2]).

Problem 1.1. Can every non-degenerate n -connected planar domain with $n > 1$ be mapped biholomorphically onto a domain of the form

$$\left\{ \left| z + \sum_{k=1}^{n-1} \frac{a_k}{z - b_k} \right| < r \right\}$$

with complex numbers a_k and b_k , and a positive r ?

Here and in the sequel, a *non-degenerate n -connected planar domain* is a subdomain Ω of the Riemann sphere $\hat{\mathbb{C}}$ such that $\hat{\mathbb{C}} - \Omega$ consists of exactly n connected components each of which contains more than one point. In this note, we solve this problem affirmatively. Actually, we give a proof of the following assertion.

Theorem 1.2. *Every non-degenerate n -connected planar domain with $n > 1$ is mapped biholomorphically onto a domain defined by*

$$\left\{ \left| z + \sum_{k=1}^{n-1} \frac{a_k}{z - b_k} \right| < 1 \right\}$$

with suitable complex numbers a_k and b_k .

Recall that every domain defined as in Theorem 1.2 has algebraic kernel functions. See Theorem 4.4 in [B1]. This is one of the reasons why we consider such domains.

Received by the editors March 15, 2002.

2000 *Mathematics Subject Classification.* Primary 32G10, 32G15; Secondary 30C20, 30F60.

Key words and phrases. Conformal representation, Ahlfors maps.

The second author was supported in part by Grant-in-Aid for Scientific Research (B)(2) 2001-13440047.

Remark 1.3. It is well known (cf. for instance [IT]) that the reduced Teichmüller space $T(\Omega)$ of a non-degenerate n -connected planar domain Ω can be identified with the Fricke space of a Fuchsian model G of Ω . Since G is a free real Möbius group with $n - 1$ hyperbolic generators, $T(\Omega)$ is real $(3n - 6)$ -dimensional.

Such a Bell representation as in Theorem 1.2 contains $2n - 2$ complex, i.e. $4n - 4$ real, parameters. The reason why we need many more number of parameters in a Bell representation than Teichmüller parameters for $T(\Omega)$ is that every Bell representation of a domain is actually associated with an n -sheeted branched covering of the unit disk by Ω .

Remark 1.4. Such a space as $H_{0,n}$ consisting of all branched coverings of $\hat{\mathbb{C}}$ induced by rational functions of degree n with $n > 1$ is called a *Hurwitz space* ([N]). This space $H_{0,n}$ is parametrized by $2n - 2$ critical values (the images of critical points), and hence is complex $(2n - 2)$ -dimensional.

Actually, we show that every Ahlfors map on Ω can be considered as the restriction of a rational function of degree n to a suitable domain which can be identified with Ω . On the other hand, the set of all branched coverings of $\hat{\mathbb{C}}$ induced from Bell representations can be considered as a subdomain of $H_{0,n}$. Thus to solve the following problem would be interesting.

Problem 1.5. Find the sublocus $A_{0,n}$ of $H_{0,n}$ which corresponds to the set of all Bell representations of non-degenerate n -connected planar domains such that the restrictions of the associated rational functions give Ahlfors maps.

The authors also thank the referee for his/her valuable suggestions and comments.

2. PROOF OF THEOREM 1.2

First let a non-degenerate n -connected planar domain Ω be given. Then by using the classical Riemann mapping theorem n times if necessary, we can assume that the boundary of Ω consists of exactly n smooth simple closed curves. Fix a point a in Ω , and let f_a be the Ahlfors map associated to the pair (Ω, a) . Here for the definition and properties of the Ahlfors maps, see for instance, [B]. In particular, f_a maps Ω properly and holomorphically onto the unit disk U . Moreover, f_a can be extended to a continuous map of the closure $\bar{\Omega}$ of Ω onto the closed unit disk so that every component γ_j of the boundary of Ω , where $j = 1, \dots, n$, is mapped homeomorphically onto the unit circle.

Lemma 2.1. *There is a compact Riemann surface R (without boundary) of genus 0 and a holomorphic injection ι of Ω into R such that*

$$f_a \circ \iota^{-1}$$

can be extended to a meromorphic function, say F , on R .

Proof. Since there are only a finite number of zeros of f'_a , there is a positive constant ρ such that $\rho < 1$ and that

$$D = \{\rho < |\zeta| < 1\},$$

where ζ is the complex coordinate on the target plane of the map f_a , and contains no critical values (i.e. no images of the zeros of f'_a by f_a). Hence every component

W_j , where $j = 1, \dots, n$, of $f_a^{-1}(D)$ is mapped biholomorphically onto D by the restriction $f_a|_{W_j}$, of f_a to W_j .

Now we construct a compact Riemann surface R by using the Ahlfors map f_a to attach disks to the exterior of Ω along each boundary curve. More precisely, we consider the disjoint union \mathbf{R} of Ω and n copies V_j ($j = 1, \dots, n$) of

$$V = \{\rho < |\zeta|\} \cup \{\infty\}.$$

Identify every subdomain W_j of Ω with the subdomain D_j of V_j corresponding to D by the biholomorphic map corresponding to $f_a|_{W_j}$. Then the resulting set, which we denote by $R = \mathbf{R}/f_a$, has a natural complex structure induced from those on Ω and on every V_j , and hence is a Riemann surface. Here the natural inclusion map ι of Ω into R is a holomorphic injection, and using the complex coordinate ζ_j on the copy V_j corresponding to ζ on V , we have

$$f_a \circ \iota^{-1}(\zeta_j) = \zeta$$

on D_j by the definition.

Now, since topologically R is obtained from Ω by attaching a disk along each boundary curve of Ω , R is a simply connected compact Riemann surface without boundary, and hence in particular, is of genus 0. Also we can extend $F = f_a \circ \iota^{-1}$ to a meromorphic function on the whole R by setting $F(\zeta_j) = \zeta$ and $F(\infty) = \infty$ on the whole V_j for every j . □

Here the following uniformization theorem (which is also called the generalized Riemann mapping theorem) is classical and well-known. As references, we cite for instance [FK] and [IT].

Proposition 2.2 (Klein, Koebe and Poincaré). *Every simply connected Riemann surface is mapped biholomorphically onto one of*

- the unit disk U ,
- the complex plane \mathbb{C} , and
- the Riemann sphere $\hat{\mathbb{C}}$.

Corollary 2.3. *There is a biholomorphic map h of the above Riemann surface R onto the Riemann sphere $\hat{\mathbb{C}}$, and hence $F \circ h^{-1}$ is a rational function.*

Proof. Since R is compact, R is mapped by a biholomorphic map h onto the Riemann sphere. Set $f = F \circ h^{-1}$. Then f is meromorphic on the whole $\hat{\mathbb{C}}$, which implies that f is a rational function. □

Here and in the sequel, we may assume that

$$f(\infty) = \infty$$

by applying to f the pre-composition of a Möbius transformation S which sends ∞ to a pole of f , i.e. by replacing h by $S^{-1} \circ h$, if necessary.

Lemma 2.4. *Let w be the complex variable of the above rational function f . Then f has the following partial fraction decomposition:*

$$f(w) = Cw + D + \sum_{k=1}^{n-1} \frac{A_k}{w - B_k}.$$

Here A_k, B_k, C and D are complex constants, every A_k and C are non-zero, and $\{B_k\}$ are mutually distinct.

Proof. Since f has exactly n simple poles, as is seen from the construction, and one of them is ∞ by the above assumption, f is of degree exactly n and has $n - 1$ finite, mutually distinct, simple poles, say B_1, \dots, B_{n-1} . Hence we can write $f(w)$ as

$$f(w) = \frac{P(w)}{Q(w)}$$

with polynomials $P(w)$ of degree exactly n and

$$Q(w) = (w - B_1) \cdots (w - B_{n-1}).$$

Thus it is easy to see that the partial fraction decomposition of f is as claimed. \square

Proof of Theorem 1.2. We replace the complex variable w of f by

$$z = T(w) = Cw + D$$

by applying to f the precomposition by an affine transformation T . Further set

$$a_k = CA_k, \quad b_k = CB_k + D$$

for every k . Then we conclude that

$$f \circ T^{-1}(z) = z + \sum_{k=1}^{n-1} \frac{a_k}{z - b_k}.$$

Thus the Bell representation

$$\left\{ \left| z + \sum_{k=1}^{n-1} \frac{a_k}{z - b_k} \right| < 1 \right\}$$

is the subdomain

$$\{|f \circ T^{-1}(z)| < 1\} = \{|F \circ h^{-1} \circ T^{-1}(z)| < 1\} = (T \circ h \circ \iota)(\Omega)$$

of $\hat{\mathbb{C}}$, which is mapped biholomorphically onto Ω by the holomorphic injection $(T \circ h \circ \iota)^{-1}$. \square

REFERENCES

- [B] S. R. Bell, *The Cauchy Transform, Potential Theory, and Conformal Mapping*, CRC Press, Boca Raton, 1992. MR **94k**:30013
- [B1] S. R. Bell, *Finitely generated function fields and complexity in potential theory in the plane*, Duke Math. J., **98** (1999), 187–207. MR **2000i**:30015
- [B2] S. R. Bell, *A Riemann surface attached to domains in the plane and complexity in potential theory*, Houston J. Math., **26**, (2000), 277–297. MR **2001m**:30009
- [FK] H. M. Farkas and I. Kra, *Riemann Surfaces*, Springer-Verlag, 1980. MR **82c**:30067
- [IT] Y. Iwayoshi and M. Taniguchi, *An Introduction to Teichmüller Spaces*, Springer-Verlag, Tokyo, 1992. MR **94b**:32031
- [N] S. M. Natanzon, *Hurwitz spaces*, Topics on Riemann surfaces and Fuchsian Groups, LMS Lecture Notes, **287** (2001), 165–177.

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF SUWON, SUWON P.O. BOX 77, KYUNG-KIDO, 440-600, KOREA

E-mail address: mjeong@mail.suwon.ac.kr

DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE, KYOTO UNIVERSITY, KYOTO 606, JAPAN

E-mail address: tanig@kusm.kyoto-u.ac.jp