

ESTIMATES ON THE MEAN GROWTH OF H^p FUNCTIONS IN CONVEX DOMAINS OF FINITE TYPE

HONG RAE CHO

(Communicated by Mei-Chi Shaw)

ABSTRACT. Let D be a bounded convex domain of finite type in \mathbb{C}^n with smooth boundary. In this paper, we prove the following inequality:

$$\left(\int_0^{\delta_0} \mathcal{M}_q^\lambda(f; t) t^{\lambda n(1/p-1/q)-1} dt \right)^{1/\lambda} \leq C_{p,q} \|f\|_{p,0},$$

where $1 < p < q < \infty$, $f \in H^p(D)$, and $p \leq \lambda < \infty$. This is a generalization of some classical result of Hardy-Littlewood for the case of the unit disc. Using this inequality, we can embed the H^p space into a weighted Bergman space in a convex domain of finite type.

1. INTRODUCTION AND STATEMENT OF RESULTS

Let D be a bounded domain in \mathbb{C}^n with smooth boundary. For $z \in D$ let $\delta(z)$ denote the distance from z to ∂D . For $\alpha > 0$, we define a measure dV_α on D by $dV_\alpha = C_\alpha \delta^{\alpha-1} dV$ where dV is the volume element and C_α is chosen so that dV_α is a probability measure. As $\alpha \rightarrow 0^+$, the measures dV_α converge as measures on ∂D to the normalized surface measure on ∂D which we denote dV_0 (or sometimes $d\sigma$). We will denote the L^p space with respect to dV_α by L_α^p , and the associated norm by $\|\cdot\|_{p,\alpha}$. We will denote by $A_\alpha^p(D) = L_\alpha^p(D) \cap \mathcal{O}(D)$ the subspace of $L_\alpha^p(D)$ consisting of functions which are holomorphic on D . In particular, $A_0^p(D)$ is the Hardy class usually denoted by $H^p(D)$, which we identify in the usual way with a subspace of $L_0^p(D) = L^p(\partial D; d\sigma)$.

Let \vec{N} be a real vector field in a neighborhood of ∂D which agree with the outward unit normal vector field on ∂D . For $z \in \partial D$ and $t > 0$ sufficiently small, say $0 < t < \delta_0$, the integral curve of \vec{N} through z has a unique intersection point with the hypersurface $\{\delta = t\}$. We call this intersection point z_t . For any function f on D we define f_t on ∂D by $f_t(z) = f(z_t)$ for $z \in \partial D$, and we define means of f by

$$\mathcal{M}_p(f; t) = \left(\int_{\partial D} |f_t|^p d\sigma \right)^{1/p}.$$

Received by the editors March 4, 2002.

2000 *Mathematics Subject Classification*. Primary 32A35, 32A26; Secondary 32T25.

Key words and phrases. Mean growth of H^p functions, convex domains of finite type, reproducing kernel.

The author was supported by grant No. R01-2000-000-00001-0 from the Basic Research Program of the Korea Science & Engineering Foundation. The author thanks the referee for helpful suggestions.

It follows from Fubini’s theorem and elementary estimates for Jacobians that for $f \in \mathcal{O}(D)$ we have $f \in A_0^p(D)$ if and only if $\sup_{0 < t < \delta_0} \mathcal{M}_p(f; t) < \infty$, and for $\alpha > 0$ we have $f \in A_\alpha^p(D)$ if and only if

$$\int_0^{\delta_0} \mathcal{M}_p^p(f; t) t^{\alpha-1} dt < \infty.$$

In this paper we get a sharp estimate on the mean growth of H^p functions on convex domains of finite type.

Theorem 1.1. *Let D be a bounded convex domain of finite type in \mathbb{C}^n with smooth boundary. If $1 < p < q < \infty, f \in H^p(D)$, and $p \leq \lambda < \infty$, then*

$$\left(\int_0^{\delta_0} \mathcal{M}_q^\lambda(f; t) t^{\lambda n(1/p-1/q)-1} dt \right)^{1/\lambda} \leq C_{p,q} \|f\|_{p,0}.$$

In the present setting we do not know whether the estimate in Theorem 1.1 remains valid when $0 < p \leq 1$. If we apply Theorem 1.1 with $\lambda = q$, then we get the following result.

Theorem 1.2. *Let D be as in Theorem 1.1 and assume that $\alpha \geq 0, 1 < p \leq q < \infty$, and $n/p = (n + \alpha)/q$. Then $H^p(D) \subset A_\alpha^q(D)$ and the inclusion is continuous.*

These results were first proved by Hardy-Littlewood for the case of the unit disc ([6], p. 87). When D is the unit ball and the strictly pseudoconvex domain, the results were proved by Beatrous-Burbea [2] and Beatrous [1]. The key point is the reproducing kernel with right estimate matching quasimetric on ∂D . For the case of convex domains of finite type we use the holomorphic support function with best possible non-isotropic estimates constructed by Diederich-Fornaess [4].

2. REPRODUCING KERNELS

Throughout, $D = \{z \in \mathbb{C}^n : \rho(z) < 0\}$ is a smoothly bounded, convex domain of finite type m defined by a real-valued function ρ with convex infralevel sets. The defining function ρ can be chosen in such a way that there exists a neighborhood U of ∂D such that $|\partial\rho(z)| > 1$ for all $z \in U$. Diederich-Fornaess [4] constructed a good C^∞ -family $S(z, \zeta)$ of support functions on D , holomorphic in $z \in \bar{D}$ and C^∞ in ζ chosen in a suitable neighborhood U of ∂D with the following estimates. For $\zeta \in U$ let \vec{n}_ζ denote the outer unit vector normal to the level set $\{\rho = \rho(\zeta)\}$ at ζ and let \vec{v} be any unit vector complex tangential to this level set at ζ . Define

$$a_{\alpha\beta}(\zeta, \vec{v}) := \frac{\partial^{\alpha+\beta}}{\partial \lambda^\alpha \partial \bar{\lambda}^\beta} \rho(\zeta + \lambda \vec{v})|_{\lambda=0}.$$

Then there are constants $K, c, d > 0$, such that one has for all points z written as $z = \zeta + \mu \vec{n}_\zeta + \lambda \vec{v}$ with $\mu, \lambda \in \mathbb{C}$ the estimate

$$(2.1) \quad \begin{aligned} 2 \operatorname{Re} S(z, \zeta) &\leq -|\operatorname{Re} \mu| - K(\operatorname{Im} \mu)^2 \\ &- c \sum_{j=2}^m \sum_{\alpha+\beta=j} |a_{\alpha\beta}(\zeta, \vec{v})| |\lambda|^j + d \sup\{0, \rho(z) - \rho(\zeta)\}. \end{aligned}$$

In (5) and (6) of [3] C^∞ functions $Q_j(z, \zeta), j = 1, \dots, n$, holomorphic in z , were defined, such that

$$S(z, \zeta) = \sum_{j=1}^n Q_j(z, \zeta)(z_j - \zeta_j).$$

Henkin [7] proved the integral representation

$$f(z) = c \int_{\zeta \in \partial D} f(\zeta) \frac{Q \wedge (\bar{\partial}^T Q)^{n-1}}{S(z, \zeta)^n}, \quad z \in D,$$

for $f \in L^1(\partial D) \cap \mathcal{O}(D)$, where $\bar{\partial}^T Q$ means the tangential components of $\bar{\partial}Q$. We define the reproducing kernel

$$K(z, \zeta) = c \frac{Q(z, \zeta) \wedge (\bar{\partial}^T Q(z, \zeta))^{n-1}}{S(z, \zeta)^n}, \quad z \in D, \zeta \in \partial D.$$

3. INTEGRAL ESTIMATES FOR THE REPRODUCING KERNELS

For $z \in U$ and $0 < \epsilon < \epsilon_0$ we define some sort of boundary distances by

$$\tau(z, \vec{v}, \epsilon) := \sup\{r > 0 : |\rho(z + \lambda \vec{v}) - \rho(z)| < \epsilon, |\lambda| \leq r, \lambda \in \mathbb{C}\}.$$

The quantity τ measures the size of the largest complex disc centered at z lying on the line spanned by \vec{v} that fits in the domain $\{\zeta : \rho(\zeta) < \rho(z) + \epsilon\}$. Next we define the ϵ -extremal basis $(\vec{v}_1, \dots, \vec{v}_n)$ centered at z of McNeal [8]. The first vector \vec{v}_1 is the unit vector in the direction of $\partial\rho(z)$; chosen $\vec{v}_1, \dots, \vec{v}_{i-1}, \vec{v}_i$ is a unit vector orthogonal to $\vec{v}_1, \dots, \vec{v}_{i-1}$. In this way we obtain a basis $(\vec{v}_1, \dots, \vec{v}_n)$ depending on both z and $\epsilon > 0$. We denote the i -th component of the coordinates with respect to this basis by w_i . We call the coordinates ϵ -extremal coordinates at z . We write $\tau_i(z, \epsilon) := \tau(z, \vec{v}_i, \epsilon)$. Since D is finite type m , we have (see [8])

$$\tau_1(z, \epsilon) \sim \epsilon \quad \text{and} \quad \epsilon^{1/2} \lesssim \tau_i(z, \epsilon) \lesssim \epsilon^{1/m} \quad \text{for} \quad 2 \leq i \leq n.$$

We can now define the non-isotropic polydiscs at z with radius ϵ by

$$P_\epsilon(z) := \left\{ \zeta = z + \sum_{i=1}^n w_i \vec{v}_i : |w_i| \leq \tau_i(z, \epsilon), i = 1, \dots, n \right\}.$$

We transform the forms $Q(z, \zeta)$ by pulling back to the ζ -variable, obtaining

$$Q^*(z, w) := \sum_{i=1}^n Q_i(z, \zeta(w)) d\zeta_i(w).$$

We write

$$Q^*(z, w) = \sum_{i=1}^n Q_i^*(z, w) dw_i.$$

Then $|Q \wedge (\bar{\partial}^T Q)^{n-1}|$ is bounded from above by the sum over all terms of the form

$$(3.1) \quad |Q_{i_1}^*(z, w)| \prod_{l=2}^n \left| \frac{\partial Q_{i_l}^*(z, w)}{\partial \bar{w}_{j_l}} \right|$$

where (i_1, \dots, i_n) is a multi-index with values from $\{1, \dots, n\}$ and $i_k \neq i_l$ for $k \neq l$ and (j_2, \dots, j_n) is a multi-index with values in $\{2, \dots, n\}$ with $j_k \neq j_l$ for $k \neq l$.

According to Lemma 5.1 of [3] we have

$$(3.2) \quad |Q_{i_1}^*(z, w)| \lesssim \frac{\epsilon}{\tau_{i_1}(z, \epsilon)}, \quad \zeta(w) \in P_\epsilon(z)$$

and

$$(3.3) \quad \left| \frac{\partial Q_{i_l}^*(z, w)}{\partial \bar{w}_{j_l}} \right| \lesssim \frac{\epsilon}{\tau_{i_l}(z, \epsilon) \tau_{j_l}(z, \epsilon)}, \quad \zeta(w) \in P_\epsilon(z).$$

By (3.2) and (3.3) each term in (3.1) is bounded by

$$\frac{\epsilon}{\tau_{i_1}(z, \epsilon)} \prod_{l=2}^n \frac{\epsilon}{\tau_{i_l}(z, \epsilon)\tau_{j_l}(z, \epsilon)}.$$

If we put this together into $|Q \wedge (\bar{\partial}^T Q)^{n-1}|$ we get

$$(3.4) \quad |Q \wedge (\bar{\partial}^T Q)^{n-1}| \lesssim \frac{\epsilon^{n-1}}{\prod_{j=2}^n \tau_j(z, \epsilon)^2}.$$

For integral estimates we define a family of polyannuli based on non-isotropic polydiscs. We choose a constant $C_1 > 0$ such that $C_1 P_{\epsilon/2}(z) \supset \frac{1}{2} P_{\epsilon}(z)$ for $\epsilon > 0$ and put for integer i

$$P_{\epsilon}^i(z) := C_1 P_{2^i \epsilon}(z) \setminus \frac{1}{2} P_{2^{i+1} \epsilon}(z).$$

Then we see that

$$\bigcup_{i=0}^{\infty} P_{\epsilon}^{-i}(z) \supset P_{\epsilon}(z) \setminus \{z\}$$

and for $\epsilon < \epsilon_0$

$$\bigcup_{i=0}^{\infty} P_{\epsilon}^i(z) \supset P_{\epsilon_0}(z) \setminus P_{\epsilon}(z).$$

Lemma 3.1 ([5]). *For integer i we have*

$$(3.5) \quad |S(z, \zeta)| \gtrsim 2^i \epsilon$$

uniformly in $z \in D \cap U$, $\zeta \in P_{\epsilon}^i(z) \cap \partial D$ (or uniformly in $\zeta \in \partial D$, $z \in P_{\epsilon}^i(\zeta) \cap D$).

Let $K_t(z, \zeta)$ be the kernel on $\partial D \times \partial D$ defined by $K_t(z, \zeta) = K(z_t, \zeta)$. Then we get the following estimates.

Proposition 3.2. *Let $s > 1$. Then we have*

$$(3.6) \quad \int_{\zeta \in \partial D} |K_t(z, \zeta)|^s d\sigma(\zeta) \lesssim t^{-n(s-1)} \quad \text{uniformly in } z \in \partial D,$$

$$(3.7) \quad \int_{z \in \partial D} |K_t(z, \zeta)|^s d\sigma(z) \lesssim t^{-n(s-1)} \quad \text{uniformly in } \zeta \in \partial D.$$

Proof. We treat only the case (3.6), since the other case (3.7) is similar. We denote

$$I(X) = \int_{\partial D \cap X} |K_t(z, \zeta)|^s d\sigma(\zeta).$$

For $z \in \partial D$ we write $z_t = \zeta + \mu \bar{n}_{z_t} + \lambda \bar{v}$. Then we have from (2.1) that

$$-2 \operatorname{Re} S(z_t, \zeta) \geq |\operatorname{Re} \mu| + K(\operatorname{Im} \mu)^2 + c \sum_{j=2}^m \sum_{\alpha+\beta=j} |a_{\alpha\beta}(\zeta, \bar{v})| |\lambda|^j.$$

Since D is finite type m , it follows that

$$\sum_{j=2}^m \sum_{\alpha+\beta=j} |a_{\alpha\beta}(\zeta, \bar{v})| |\lambda|^j \gtrsim |\lambda|^m.$$

Since $m \geq 2$, we have $|z_t - \zeta|^m = |\mu \bar{n}_{z_t} + \lambda \bar{v}|^m \lesssim |\operatorname{Re} \mu| + K(\operatorname{Im} \mu)^2 + |\lambda|^m$. Thus it follows that

$$-2 \operatorname{Re} S(z_t, \zeta) \gtrsim |z_t - \zeta|^m$$

and hence the only singularity of $K_t(z, \zeta)$ occurs for $\zeta = z_t$. Thus $I(\partial D) \leq C$ if $t \geq \delta_0$ and $I(\partial D \setminus V) \leq C$ for some small neighborhood V of z_t . Hence it is enough to prove that for fixed $\epsilon_0 > 0$ and small $t > 0$

$$(3.8) \quad I(P_{\epsilon_0}(z_t)) \lesssim t^{-n(s-1)}.$$

For fixed $z \in \partial D$ we define $\rho = |\rho(z_t)| \sim t$ and then split the polydisc $P_{\epsilon_0}(z_t)$ into the two parts $P_\rho(z_t)$ and $P_{\epsilon_0}(z_t) \setminus P_\rho(z_t)$. Remember that $P_\rho(z_t) \setminus \{z_t\}$ can be covered by $\bigcup_{i=0}^{\infty} P_\rho^{-i}(z_t)$. By (3.4), it follows that

$$\begin{aligned} I(P_\rho^{-i}(z_t)) &= \int_{\partial D \cap P_\rho^{-i}(z_t)} |K_t(z, \zeta)|^s d\sigma(\zeta) \\ &\lesssim \int_{\partial D \cap P_\rho^{-i}(z_t)} \frac{1}{|S(z_t, \zeta)|^{ns}} \frac{(2^{-i}\rho)^{(n-1)s}}{\prod_{j=2}^n \tau_j(z_t, 2^{-i}\rho)^{2s}} d\sigma(\zeta). \end{aligned}$$

Since $\rho = |\rho(z_t)| = |\rho(z_t) - \rho(\zeta)| \lesssim |\operatorname{Re} \mu|$, it follows that $|S(z_t, \zeta)| \gtrsim \rho$ and hence we have

$$\begin{aligned} I(P_\rho^{-i}(z_t)) &\lesssim \frac{(2^{-i})^{(n-1)s}}{\rho^s} \int_{|v_1| < \tau_1(z_t, 2^{-i}\rho)} dv_1 \prod_{j=2}^n \int_{|w_j| < \tau_j(z_t, 2^{-i}\rho)} \frac{du_j dv_j}{\tau_j(z_t, 2^{-i}\rho)^{2s}} \\ &\lesssim \frac{(2^{-i})^{(n-1)s}}{\rho^s} \tau_1(z_t, 2^{-i}\rho) \prod_{j=2}^n \tau_j(z_t, 2^{-i}\rho)^{-2(s-1)} \\ &\lesssim \frac{(2^{-i})^{(n-1)s}}{\rho^s} (2^{-i}\rho)^{-(s-1)(n-1)} = \rho^{-n(s-1)} (2^{-i})^n. \end{aligned}$$

Thus we get

$$(3.9) \quad \begin{aligned} I(P_\rho(z_t)) &\leq \sum_{i=0}^{\infty} I(P_\rho^{-i}(z_t)) \\ &\lesssim \sum_{i=0}^{\infty} \rho^{-n(s-1)} (2^{-i})^n \lesssim \rho^{-n(s-1)} \sim t^{-n(s-1)}. \end{aligned}$$

To estimate the integral over $P_{\epsilon_0}(z_t) \setminus P_\rho(z_t)$ we use the covering by $\bigcup_{i=0}^{\infty} P_\rho^i(z_t)$. By (3.4) and (3.5), we have

$$\begin{aligned} I(P_\rho^i(z_t)) &\lesssim \int_{\partial D \cap P_\rho^i(z_t)} \frac{1}{|S(z_t, \zeta)|^{ns}} \frac{(2^i\rho)^{(n-1)s}}{\prod_{j=2}^n \tau_j(z_t, 2^i\rho)^{2s}} d\sigma(\zeta) \\ &\lesssim \frac{1}{(2^i\rho)^s} \tau_1(z_t, 2^i\rho) \prod_{j=2}^n \tau_j(z_t, 2^i\rho)^{-2(s-1)} \\ &\lesssim \rho^{-n(s-1)} (2^i)^{-n(s-1)}. \end{aligned}$$

Thus we get

$$(3.10) \quad \begin{aligned} I(P_{\epsilon_0}(z_t) \setminus P_\rho(z_t)) &\leq \sum_{i=0}^{\infty} I(P_\rho^i(z_t)) \\ &\lesssim \sum_{i=0}^{\infty} \rho^{-n(s-1)} (2^i)^{-n(s-1)} \lesssim \rho^{-n(s-1)} \sim t^{-n(s-1)}. \end{aligned}$$

By (3.9) and (3.10), we proved the required estimate (3.8). \square

4. PROOFS OF MAIN RESULTS

For $f \in L^1(\partial D; d\sigma)$ we define

$$\mathcal{K}f(z) = \int_{\zeta \in \partial D} f(\zeta)K(z, \zeta), \quad z \in D.$$

By using Proposition 3.2, we can prove as in ([1], Lemma 2.8) that

$$(4.1) \quad \left(\int_0^{\delta_0} \mathcal{M}_q^\lambda(\mathcal{K}f; t) t^{\lambda n(1/p-1/q)-1} dt \right)^{1/\lambda} \leq C_{p,q} \|f\|_{p,0},$$

where $1 < p < q < \infty$ and $q \leq \lambda < \infty$. Since $K(z, \zeta)$ is the reproducing kernel for holomorphic functions, we get Theorem 1.1 from (4.1). Moreover, for $\alpha > 0$, $1 < p \leq q < \infty$, and $n/p = (n + \alpha)/q$, if we apply (4.1) with $\lambda = q$ we have

$$\int_D |\mathcal{K}f|^q dV_\alpha \lesssim \|f\|_{p,0}^q.$$

Thus the integral operator $\mathcal{K} : L^p(\partial D; d\sigma) \rightarrow A_\alpha^q(D)$ is bounded and so we get Theorem 1.2.

REFERENCES

1. F. Beatrous, *Estimates for derivatives of holomorphic functions in pseudoconvex domains*, Math. Z. **191** (1986), 91–116. MR **87b**:32033
2. F. Beatrous and J. Burbea, *Holomorphic Sobolev spaces on the ball*, Dissertationes Math. **256** (1989), 1–57. MR **90k**:32010
3. K. Diederich, B. Fischer and J.E. Fornæss, *Hölder estimates on convex domains of finite type*, Math. Z. **232** (1999), 43–61. MR **2001f**:32065
4. K. Diederich and J.E. Fornæss, *Support functions for convex domains of finite type*, Math. Z. **230** (1999), 145–164. MR **2000b**:32024
5. K. Diederich and E. Mazzilli, *Zero varieties for the Nevanlinna class on all convex domains of finite type*, Nagoya Math. J. **163** (2001), 215–227. MR **2003b**:32006
6. P. Duren, *Theory of H^p spaces*, Academic Press, New York, 1970. MR **42**:3552
7. G. Henkin, *Integral representations of functions holomorphic in strictly pseudoconvex domains and some applications*, Math. USSR, Sb. **7** (1969), 597–616. MR **40**:2902
8. J.D. McNeal, *Estimates on the Bergman kernels of convex domains*, Adv. in Math. **109** (1994), 108–139. MR **95k**:32023

DEPARTMENT OF MATHEMATICS EDUCATION, ANDONG NATIONAL UNIVERSITY, ANDONG 760-749, SOUTH KOREA

E-mail address: chohr@anu.ac.kr

Current address: Department of Mathematics, Pusan National University, Pusan 609-735, Korea

E-mail address: chohr@pusan.ac.kr