

RECONSTRUCTION OF FUNCTIONS IN SPLINE SUBSPACES FROM LOCAL AVERAGES

WENCHANG SUN AND XINGWEI ZHOU

(Communicated by David R. Larson)

ABSTRACT. In this paper, we study the reconstruction of functions in spline subspaces from local averages. We present an average sampling theorem for shift invariant subspaces generated by cardinal B-splines and give the optimal upper bound for the support length of averaging functions. Our result generalizes an earlier result by Aldroubi and Gröchenig.

1. INTRODUCTION AND THE MAIN RESULT

The sampling theorem is one of the most powerful results in signal analysis. It states that if a signal (function) $f(x)$ satisfies certain conditions, then it is uniquely determined by its sampled values at some discrete points $\{x_k : k \in \mathbb{Z}\}$ and

$$f(x) = \sum_{k \in \mathbb{Z}} f(x_k) S_k(x)$$

for some sampling functions $\{S_k(x)\}$. For example, every $f \in B_\Omega := \{f : \text{supp } \hat{f} \subset [-\Omega, \Omega]\}$ can be reconstructed by the formula $f(x) = \sum_{k \in \mathbb{Z}} f\left(\frac{k\pi}{\Omega}\right) \frac{\sin(\Omega x - k\pi)}{\Omega x - k\pi}$. This is the classical Shannon sampling theorem. Although the assumption that a signal is band-limited is eminently useful, it is not always realistic since a band-limited signal is of infinite duration. Thus, it is natural to investigate other signal classes for which a sampling theorem holds. A simple model is to consider shift-invariant subspaces, e.g., wavelet subspaces, which generalize the space of band-limited functions and are of the form

$$V = \overline{\text{span}}\{\varphi(\cdot - k) : k \in \mathbb{Z}\}$$

for some kernel function $\varphi(x)$. In fact, there have been many results concerning the sampling in shift-invariant subspaces for both regular and irregular sampling. For details, see [1]–[7], [10]–[19], [22]–[25] and [27].

Received by the editors May 11, 2001.

2000 *Mathematics Subject Classification*. Primary 94A20; Secondary 42C40, 42C15.

Key words and phrases. Average sampling, sampling theorems, spline subspaces.

This work was supported by the Research Fund for the Doctoral Program of Higher Education, the National Natural Science Foundation of China (Grant Nos. 10171050 and 10201014), the Mathematical Tianyuan Foundation(Grant No. TY10126007), and the Liuhui Center for Applied Mathematics.

In particular, for the spline subspace

$$V_m = \left\{ \sum_{k \in \mathbb{Z}} c_k \varphi_m(\cdot - k) : \{c_k\} \in \ell^2 \right\}$$

generated by the cardinal B-spline

$$\varphi_m = \chi_{[0,1]} * \cdots * \chi_{[0,1]} \quad (m + 1 \text{ terms}), \quad m \geq 1,$$

Liu [16] proved that every $f \in V_m$ is uniquely determined and can be reconstructed by an iterative algorithm from its samples $f(x_k)$ if $x_{k+1} - x_k$ is small enough. Furthermore, Aldroubi and Gröchenig [1] proved that if $\alpha \leq x_{k+1} - x_k \leq \beta$ for some constants $0 < \alpha < \beta < 1$, then every $f \in V_m$ is uniquely determined by $f(x_k)$.

For physical reasons, e.g., the inertia of the measurement apparatus, measured sampled values obtained in practice may not be values of a function f precisely at times x_k , but only local averages of f near x_k . Specifically, measured sampled values are

$$\langle f, u_k \rangle = \int f(x) u_k(x) dx$$

for some collection of averaging functions $u_k(x)$, $k \in \mathbb{Z}$, which satisfy the following properties:

$$\text{supp } u_k \subset \left[x_k - \frac{\delta}{2}, x_k + \frac{\delta}{2} \right], \quad u_k(x) \geq 0, \quad \text{and} \quad \int u(x) dx = 1.$$

Observe that the averaging procedure is allowed to vary from point to point.

It is clear that from local averages one should obtain at least a good approximation of the original signal if δ is small enough. Wiley, Butzer and Lei [5, 6, 24] studied the approximation error when local averages are used as sampled values. Furthermore, Gröchenig [14] proved that if sampling points $\{x_k : k \in \mathbb{Z}\}$ satisfy $0 < x_{k+1} - x_k \leq \delta < \frac{1}{\sqrt{2\Omega}}$, then every $f \in B_\Omega$ is uniquely determined by local averages $\langle f, u_k \rangle$ around x_k and can be reconstructed by an iteration scheme. In [13], Feichtinger and Gröchenig proved that if $\delta := \sup_{k \in \mathbb{Z}} (x_{k+1} - x_k) < \frac{\pi}{\Omega}$, then every $f \in B_\Omega$ is uniquely determined by $\frac{1}{y_k - y_{k-1}} \int_{y_{k-1}}^{y_k} f(x) dx$ with $y_k = \frac{x_k + x_{k+1}}{2}$.

In [20], we gave an average sampling theorem for shift invariant subspaces with equally spaced sampling points and arbitrary averaging functions. For the case of band-limited signals, we gave an analogy of Kadec's 1/4-theorem. In [21], we studied average sampling theorems for spline subspaces with standard averaging functions $u_k = \chi_{[x_k - 1/2, x_k + 1/2]}$.

In this paper, we study the reconstruction of functions in spline subspaces from local averages with arbitrary averaging functions and irregular sampling points. We present an average sampling theorem and give the optimal upper bound for the support length of averaging functions. Our result generalizes Aldroubi and Gröchenig's irregular sampling theorem. In fact, we prove the following.

Theorem 1.1. *Let $\{x_k : k \in \mathbb{Z}\}$ be a real sequence such that $\lim_{k \rightarrow \pm\infty} x_k = \pm\infty$ and*

$$0 < x_{k+1} - x_k \leq \beta < 1, \quad k \in \mathbb{Z},$$

for some constant β . Then for any $0 < \delta < 1 - \beta$ and averaging functions $u_k(x)$ with $\text{supp } u_k \subset [x_k - \frac{\delta}{2}, x_k + \frac{\delta}{2}]$, there is a frame $\{S_k(x) : k \in \mathbb{Z}\}$ for V_m such that

for any $f \in V_m$,

$$(1.1) \quad f(x) = \sum_{k \in \mathbb{Z}} \left(\frac{x_{k+1} - x_{k-1}}{2} \right)^{1/2} \langle f, u_k \rangle S_k(x),$$

where the convergence is both in $L^2(\mathbb{R})$ and uniform on \mathbb{R} .

Furthermore, the conclusion fails if $\delta \geq 1 - \beta$.

Notation and definition. The Fourier transform and the Zak transform of $f \in L^2(\mathbb{R})$ is defined by $\hat{f}(\omega) = \int_{\mathbb{R}} f(x)e^{-ix\omega} dx$ and $Zf(x, \omega) = \sum_{k \in \mathbb{Z}} f(x+k)e^{-ik\omega}$ respectively.

We call $u(x)$ an averaging function if $u(x) \geq 0, u(x) \in L^1(\mathbb{R})$ and $\int_{\mathbb{R}} u(x)dx = 1$.

2. SOME BASIC FACTS IN FRAME THEORY

In this section, we introduce some basic facts in frame theory.

A family of functions $\{\varphi_k : k \in \mathbb{Z}\}$ belonging to a separable Hilbert space \mathcal{H} is said to be a frame if there exist positive constants A and B such that $A\|f\|^2 \leq \sum_{k \in \mathbb{Z}} |\langle f, \varphi_k \rangle|^2 \leq B\|f\|^2$ for every $f \in \mathcal{H}$. The constants A and B are called the lower and upper frame bounds respectively.

A frame that ceases to be a frame when any one of its elements is removed is said to be an exact frame. It is well known that exact frames and Riesz bases are identical (see [26]).

Let $\{\varphi_k : k \in \mathbb{Z}\}$ be a frame for some Hilbert space \mathcal{H} with bounds A and B . Define the frame operator T as follows: $Tf = \sum_{n \in \mathbb{Z}} \langle f, \varphi_n \rangle \varphi_n, \forall f \in \mathcal{H}$. Then T is bounded and invertible and $\{T^{-1}\varphi_k : k \in \mathbb{Z}\}$ is also a frame for \mathcal{H} with bounds $\frac{1}{B}$ and $\frac{1}{A}$, called the dual frame of $\{\varphi_k : k \in \mathbb{Z}\}$. It can be shown that for any $f \in \mathcal{H}$,

$$f = \sum_{n \in \mathbb{Z}} \langle f, T^{-1}\varphi_n \rangle \varphi_n = \sum_{n \in \mathbb{Z}} \langle f, \varphi_n \rangle T^{-1}\varphi_n.$$

For the general theory on frames and Riesz bases, see [9, 26].

For frames in shift invariant subspaces, we have the following.

Lemma 2.1. Let $\{\varphi(\cdot - k) : k \in \mathbb{Z}\}$ be a frame for some closed subspace V of $L^2(\mathbb{R})$. Suppose that φ is continuous and $\sum_{k \in \mathbb{Z}} |\varphi(x - k)|^2 \leq L < +\infty$. Then for any frame $\{S_k : k \in \mathbb{Z}\}$ of V_0 , $\sum_{k \in \mathbb{Z}} |S_k(x)|^2$ is bounded on \mathbb{R} .

Proof. Let $\{\tilde{\varphi}(\cdot - k) : k \in \mathbb{Z}\}$ be the dual frame of $\{\varphi(\cdot - k) : k \in \mathbb{Z}\}$. Then for any $f \in V, f(x) = \sum_{k \in \mathbb{Z}} \langle f, \tilde{\varphi}(\cdot - k) \rangle \varphi(x - k)$. Therefore,

$$\|f\|_{\infty}^2 \leq \sup_x \sum_{k \in \mathbb{Z}} |\langle f, \tilde{\varphi}(\cdot - k) \rangle|^2 \sum_{k \in \mathbb{Z}} |\varphi(x - k)|^2 \leq \frac{L}{A} \|f\|_2^2.$$

It follows that for any $x \in \mathbb{R}$,

$$\sum_{k \in \mathbb{Z}} |S_k(x)|^2 = \sup_{\|c\|_2=1} \left| \sum_{k \in \mathbb{Z}} c_k S_k(x) \right|^2 \leq \sup_{\|c\|_2=1} \frac{L}{A} \left\| \sum_{k \in \mathbb{Z}} c_k S_k \right\|_2^2 \leq \frac{LM}{A},$$

where M is the upper frame bound of $\{S_k(x) : k \in \mathbb{Z}\}$. □

Lemma 2.2. *Suppose that $\{u_k : k \in \mathbb{Z}\}$ is a sequence of compactly supported averaging functions and there exist positive constants A and B such that*

$$(2.1) \quad A\|f\|^2 \leq \sum_{k \in \mathbb{Z}} |\langle f, u_k \rangle|^2 \leq B\|f\|^2, \quad \forall f \in V_m.$$

Then there is a frame $\{S_k : k \in \mathbb{Z}\}$ for V_m such that $f(x) = \sum_{k \in \mathbb{Z}} \langle f, u_k \rangle S_k(x), \forall f \in V_m$, where the convergence is both in $L^2(\mathbb{R})$ and uniform on \mathbb{R} .

Proof. By [8, Theorem 4.5], $\{\varphi_m(\cdot - n) : n \in \mathbb{Z}\}$ is a Riesz basis for V_m . Let $\{\tilde{\varphi}_m(\cdot - n) : n \in \mathbb{Z}\}$ be the dual Riesz basis. For any $k \in \mathbb{Z}$, let

$$h_k = \sum_{n \in \mathbb{Z}} \langle u_k, \varphi_m(\cdot - n) \rangle \tilde{\varphi}_m(\cdot - n).$$

Since both φ_m and u_k are compactly supported, the above series contains only finite non-zero terms. Hence $h_k \in V_m$. For any $f \in V_m$, we have

$$\begin{aligned} \langle f, h_k \rangle &= \sum_{n \in \mathbb{Z}} \langle \varphi_m(\cdot - n), u_k \rangle \cdot \langle f, \tilde{\varphi}_m(\cdot - n) \rangle \\ &= \left\langle \sum_{n \in \mathbb{Z}} \langle f, \tilde{\varphi}_m(\cdot - n) \rangle \varphi_m(\cdot - n), u_k \right\rangle \\ &= \langle f, u_k \rangle. \end{aligned}$$

Now (2.1) shows that $\{h_k : k \in \mathbb{Z}\}$ is a frame for V_m . Let $\{S_k : k \in \mathbb{Z}\}$ be the dual frame. Then for any $f \in V_m$, $f(x) = \sum_{k \in \mathbb{Z}} \langle f, h_k \rangle S_k(x) = \sum_{k \in \mathbb{Z}} \langle f, u_k \rangle S_k(x)$. The uniform convergence follows by Lemma 2.1. \square

3. PROOF OF THEOREM 1.1

First, we introduce a result by Aldroubi and Gröchenig. The following proposition is a special case of [1, Theorem 1].

Proposition 3.1. *Let $\{x_k : k \in \mathbb{Z}\}$ be a real sequence such that*

$$0 < \alpha < x_{k+1} - x_k < \beta < 1, \quad \forall k \in \mathbb{Z},$$

for some constants α and β . Then there exist positive constants C_1 and C_2 depending only on m, α , and β such that

$$C_1\|f\|_2^2 \leq \sum_{k \in \mathbb{Z}} |f(x_k)|^2 \leq C_2\|f\|_2^2, \quad \forall f \in V_m.$$

Lemma 3.2. $\inf_{f \in V_m} \frac{1}{\|f\|_2^2} \sum_{k \in \mathbb{Z}} |f(\frac{m}{2} + k)|^2 = 0$.

Proof. Since $\varphi_m(x)$ is symmetric with respect to $x = \frac{m+1}{2}$ and $\sum_{k \in \mathbb{Z}} \varphi_m(x-k) = 1$ for any $x \in \mathbb{R}$, it is easy to check that $Z\varphi_m(\frac{m}{2}, \pi) = 0$.

For any $\varepsilon > 0$, by the continuity of $Z\varphi_m(x, \omega)$, there exists some $0 < \delta < \pi$ such that

$$|Z\varphi_m(\frac{m}{2}, \omega)| \leq \varepsilon, \quad |\omega - \pi| \leq \delta.$$

Let $C(\omega) = \sum_{k \in \mathbb{Z}} c_k e^{-ik\omega}$ be a 2π -periodic function such that $C(\omega) = 1$ for $\omega \in [\pi - \delta, \pi + \delta]$ and $C(\omega) = 0$ for $\omega \in [0, \pi - \delta] \cup [\pi + \delta, 2\pi]$. Define $\hat{f}(\omega) = C(\omega)\hat{\varphi}_m(\omega)$.

Then we have

$$\begin{aligned} \sum_{k \in \mathbb{Z}} |f(\frac{m}{2} + k)|^2 &= \frac{1}{2\pi} \int_0^{2\pi} |Zf(\frac{m}{2}, \omega)|^2 d\omega \\ &= \frac{1}{2\pi} \int_0^{2\pi} |C(\omega)Z\varphi_m(\frac{m}{2}, \omega)|^2 d\omega \leq \frac{\varepsilon^2}{2\pi} \int_{\pi-\delta}^{\pi+\delta} |C(\omega)|^2 d\omega = \varepsilon^2 \sum_{k \in \mathbb{Z}} |c_k|^2. \end{aligned}$$

Since $\{\varphi_m(\cdot - n) : n \in \mathbb{Z}\}$ is a Riesz basis for V_m with lower bound $A_m > 0$, the above inequalities show that $\sum_{k \in \mathbb{Z}} |f(\frac{m}{2} + k)|^2 < \frac{\varepsilon^2}{A_m}$. Since ε is arbitrary, the conclusion follows. \square

On the other hand, since $\varphi_m(x)$ is compactly supported and absolutely continuous for any $m \geq 1$, we also have

Lemma 3.3. *For any $f \in V_m, f' \in L^2(\mathbb{R})$.*

We are now ready to give the proof of the main result.

Proof of Theorem 1.1. For any $f \in V_m$, let

$$f_1(x) = \frac{f(x) + \overline{f(x)}}{2}, \quad f_2(x) = \frac{f(x) - \overline{f(x)}}{2i}.$$

Then $f_1, f_2 \in V_m$ and $f(x) = f_1(x) + if_2(x)$.

Since $f_p(x)$ is continuous on \mathbb{R} and $u_k(x) \geq 0$, by the first mean value theorem, there are some $x_{p,k} \in [x_k - \frac{\delta}{2}, x_k + \frac{\delta}{2}]$ such that

$$\langle f_p, u_k \rangle = f_p(x_{p,k}), \quad p = 1, 2.$$

Put $0 < \alpha < \frac{1}{3}(1 - \beta - \delta)$. Let

$$\begin{aligned} n_0 &= 0, \\ n_k &= \begin{cases} \min\{n : n > n_{k-1}, x_n - x_{n_{k-1}} \geq \alpha\}, & k > 0, \\ \max\{n : n < n_{k+1}, x_{n_{k+1}} - x_n \geq \alpha\}, & k < 0. \end{cases} \end{aligned}$$

Then

$$\alpha \leq x_{n_k} - x_{n_{k-1}} \leq \beta + \alpha, \quad k \in \mathbb{Z}.$$

For any $k \in \mathbb{Z}$, let j_k and j'_k be the minimum and maximum of $\{j : \frac{x_{n_{k-1}} + x_{n_k}}{2} < x_j \leq \frac{x_{n_k} + x_{n_{k+1}}}{2}\}$ respectively. Then $\{j : j_k \leq j \leq j'_k\}_{k \in \mathbb{Z}}$ is a partition of \mathbb{Z} .

For any $n_k < j \leq j'_k$, since $j'_k < n_{k+1}$, by the definition of n_k ,

$$\min\{x_j - x_{n_k}, x_{n_{k+1}} - x_j\} < \alpha.$$

But $x_{n_{k+1}} - x_j \geq \frac{x_{n_{k+1}} - x_{n_k}}{2} \geq x_j - x_{n_k} > 0$, so $x_j - x_{n_k} < \alpha$. A similar argument shows that $x_{n_k} - x_j < \alpha$ for $j_k \leq j < n_k$. Hence for any $j_k \leq j \leq j'_k, |x_j - x_{n_k}| < \alpha$. Therefore,

$$x_{p,j} \in [x_{n_k} - \alpha - \frac{\delta}{2}, x_{n_k} + \alpha + \frac{\delta}{2}], \quad j_k \leq j \leq j'_k.$$

It follows from the continuity of $f_p(x)$ and $\sum_{j_k \leq j \leq j'_k} (x_{j+1} - x_{j-1}) = x_{j'_k+1} - x_{j_k-1} + x_{j'_k} - x_{j_k}$ that there is some $y_{p,k} \in [x_{n_k} - \alpha - \frac{\delta}{2}, x_{n_k} + \alpha + \frac{\delta}{2}]$ such that

$$\begin{aligned}
 (3.1) \quad & \sum_{j_k \leq j \leq j'_k} \frac{x_{j+1} - x_{j-1}}{2} |f_p(x_{p,j})|^2 \\
 &= \frac{x_{j'_k+1} - x_{j_k-1} + x_{j'_k} - x_{j_k}}{2} \sum_{j_k \leq j \leq j'_k} \frac{x_{j+1} - x_{j-1}}{x_{j'_k+1} - x_{j_k-1} + x_{j'_k} - x_{j_k}} |f_p(x_{p,j})|^2 \\
 &= \frac{x_{j'_k+1} - x_{j_k-1} + x_{j'_k} - x_{j_k}}{2} |f_p(y_{p,k})|^2.
 \end{aligned}$$

Let $\{z_{p,k} : k \in \mathbb{Z}\}$ be a rearrangement of $\{y_{p,k} : k \in \mathbb{Z}\}$ such that $z_{p,k} \leq z_{p,k+1}$. Since $0 < x_{n_{k+1}} - x_{n_k} \leq \beta + \alpha$ and $|y_{p,k} - x_{n_k}| \leq \alpha + \frac{\delta}{2}$, it is easy to check that every interval with length $\beta + \delta + 3\alpha$ must contain at least one point of $\{y_{p,k} : k \in \mathbb{Z}\}$. Hence

$$0 \leq z_{p,k+1} - z_{p,k} \leq \beta + \delta + 3\alpha.$$

Since $\beta + \delta + 3\alpha < 1$, similar to the choice of x_{n_k} we can prove that there is a subsequence $\{z_{p,i_k} : k \in \mathbb{Z}\}$ of $\{z_{p,k} : k \in \mathbb{Z}\}$ such that

$$\varepsilon \leq z_{p,i_{k+1}} - z_{p,i_k} \leq \beta + \delta + 3\alpha + \varepsilon < 1$$

for some $\varepsilon > 0$. By Proposition 3.1, there is a positive constant C_1 depending only on m, ε and $\beta + \delta + 3\alpha + \varepsilon$ such that

$$\sum_{k \in \mathbb{Z}} |f(z_{p,i_k})|^2 \geq C_1 \|f\|_2^2, \quad \forall f \in V_m.$$

Hence

$$(3.2) \quad \sum_{k \in \mathbb{Z}} |f_p(y_{p,k})|^2 = \sum_{k \in \mathbb{Z}} |f_p(z_{p,k})|^2 \geq C_1 \|f_p\|_2^2.$$

Noting that

$$\frac{x_{j'_k+1} - x_{j_k-1} + x_{j'_k} - x_{j_k}}{2} \geq \frac{x_{j'_k+1} - x_{j_k-1}}{2} \geq \frac{x_{n_{k+1}} - x_{n_{k-1}}}{4} \geq \frac{\alpha}{2},$$

we derive from (3.1) and (3.2) that

$$\begin{aligned}
 (3.3) \quad \sum_{k \in \mathbb{Z}} \frac{x_{k+1} - x_{k-1}}{2} |\langle f_p, u_k \rangle|^2 &= \sum_{k \in \mathbb{Z}} \frac{x_{k+1} - x_{k-1}}{2} |f_p(x_{p,k})|^2 \\
 &= \sum_{k \in \mathbb{Z}} \sum_{j_k \leq j \leq j'_k} \frac{x_{j+1} - x_{j-1}}{2} |f_p(x_{p,j})|^2 \\
 &= \sum_{k \in \mathbb{Z}} \frac{x_{j'_k+1} - x_{j_k-1} + x_{j'_k} - x_{j_k}}{2} |f_p(y_{p,k})|^2 \\
 &\geq \frac{\alpha C_1}{2} \|f_p\|_2^2.
 \end{aligned}$$

On the other hand, it can be shown (see Section 4) that there is some constant C_2 such that

$$(3.4) \quad \sum_{k \in \mathbb{Z}} \frac{x_{k+1} - x_{k-1}}{2} |\langle f_p, u_k \rangle|^2 = \sum_{k \in \mathbb{Z}} \frac{x_{k+1} - x_{k-1}}{2} |f_p(x_{p,k})|^2 \leq C_2 \|f_p\|_2^2.$$

Since $f(x) = f_1(x) + if_2(x)$ and f_1, f_2, u_k are real functions, we have $\|f\|_2^2 = \|f_1\|_2^2 + \|f_2\|_2^2$ and $|\langle f, u_k \rangle|^2 = |\langle f_1, u_k \rangle|^2 + |\langle f_2, u_k \rangle|^2$. Therefore, (3.3) and (3.4) hold if we substitute f for f_p . Now the first part of this theorem follows by Lemma 2.2.

Next we are to prove the second part. We only need to show that the conclusion fails if $\delta = 1 - \beta$.

First, we consider the case of $\frac{1}{2} \leq \beta < 1$.

Let $x_{2k} = k + \frac{m-1}{2} - \frac{\beta}{2}$ and $x_{2k+1} = k + \frac{m-1}{2} + \frac{\beta}{2}$. Then $1 - \beta \leq x_{k+1} - x_k \leq \beta$. Put $\varepsilon_k = \frac{1-\beta}{2+|k|}$. Choose some averaging functions $u_k(x)$ such that

$$\begin{aligned} \text{supp } u_{2k} &\subset [k - 1 + \frac{m}{2}, k - 1 + \frac{m}{2} + \varepsilon_k], \\ \text{supp } u_{2k+1} &\subset [k + \frac{m}{2} - \varepsilon_k, k + \frac{m}{2}]. \end{aligned}$$

Then $\text{supp } u_k \subset [x_k - \frac{\delta}{2}, x_k + \frac{\delta}{2}]$.

For any $\varepsilon > 0$, there exists some $f \in V_m$ such that

$$(3.5) \quad \sum_{k \in \mathbb{Z}} |f(\frac{m}{2} + k)|^2 \leq \varepsilon \|f\|_2^2,$$

thanks to Lemma 3.2. Since $\varphi_m(x)$ is real-valued, we can assume that $f(x)$ is also real-valued. Let $f_n(x) = f(x - n), n \geq 1$. Then $f_n \in V_m$. By the first mean value theorem, there are some $x_{n,2k} \in [k - 1 + \frac{m}{2}, k - 1 + \frac{m}{2} + \varepsilon_k]$ and $x_{n,2k+1} \in [k + \frac{m}{2} - \varepsilon_k, k + \frac{m}{2}]$ such that

$$\langle f_n, u_k \rangle = f_n(x_{n,k}).$$

For any $p > 0$, we have

$$\begin{aligned} &\sum_{|k| \geq p} |f_n(x_{n,2k}) - f_n(k - 1 + \frac{m}{2})|^2 \\ &= \sum_{|k| \geq p} \left| \int_0^{x_{n,2k} - k + 1 - \frac{m}{2}} f'_n(k - 1 + \frac{m}{2} + t) dt \right|^2 \\ &\leq \sum_{|k| \geq p} \varepsilon_k \int_0^{\varepsilon_k} |f'_n(k - 1 + \frac{m}{2} + t)|^2 dt \\ &\leq \varepsilon_p \int_0^{\varepsilon_p} \sum_{k \in \mathbb{Z}} |f'_n(k - 1 + \frac{m}{2} + t)|^2 dt \\ &\leq \varepsilon_p \|f'_n\|_2^2 = \varepsilon_p \|f'\|_2^2. \end{aligned}$$

It follows from (3.5) that

$$(3.6) \quad \sum_{|k| \geq p} |f_n(x_{n,2k})|^2 \leq (\varepsilon_p^{1/2} \|f'\|_2 + \varepsilon^{1/2} \|f\|_2)^2.$$

On the other hand, since

$$\begin{aligned}
 \sum_{|k| \leq n/2} |f_n(x_{n,2k}) - f_n(k - 1 + \frac{m}{2})|^2 &\leq \sum_{|k| \leq n/2} \varepsilon_k \int_0^{\varepsilon_k} |f'_n(k - 1 + \frac{m}{2} + t)|^2 dt \\
 &\leq \int_0^1 \sum_{|k| \leq n/2} |f'_n(k - 1 + \frac{m}{2} + t)|^2 dt \\
 &\leq \int_0^1 \sum_{|l| \geq n/2} |f'(l - 1 + \frac{m}{2} + t)|^2 dt \\
 &\leq \int_{|t| \geq \frac{n-m}{2}} |f'(t)|^2 dt \\
 &:= \|T_{\frac{n-m}{2}} f'\|_2^2,
 \end{aligned}$$

we have

$$\begin{aligned}
 (3.7) \quad \left(\sum_{|k| \leq n/2} |f_n(x_{n,2k})|^2 \right)^{1/2} &\leq \left(\sum_{k \in \mathbb{Z}} |f_n(k - 1 + \frac{m}{2})|^2 \right)^{1/2} + \|T_{\frac{n-m}{2}} f'\|_2 \\
 &\leq \varepsilon^{1/2} \|f\|_2 + \|T_{\frac{n-m}{2}} f'\|_2.
 \end{aligned}$$

Putting (3.6) and (3.7) together, we have

$$\begin{aligned}
 (3.8) \quad \sum_{k \in \mathbb{Z}} |f_n(x_{n,2k})|^2 &= \sum_{|k| > n/2} |f_n(x_{n,2k})|^2 + \sum_{|k| \leq n/2} |f_n(x_{n,2k})|^2 \\
 &\leq (\varepsilon_n^{1/2} \|f'\|_2 + \varepsilon^{1/2} \|f\|_2)^2 + (\varepsilon^{1/2} \|f\|_2 + \|T_{\frac{n-m}{2}} f'\|_2)^2.
 \end{aligned}$$

A similar argument shows that (3.8) holds if we substitute $f_n(x_{n,2k+1})$ for $f_n(x_{n,2k})$. Therefore,

$$\sum_{k \in \mathbb{Z}} |\langle f_n, u_k \rangle|^2 \leq 2(\varepsilon_n^{1/2} \|f'\|_2 + \varepsilon^{1/2} \|f\|_2)^2 + 2(\varepsilon^{1/2} \|f\|_2 + \|T_{\frac{n-m}{2}} f'\|_2)^2.$$

Noting that $\|f_n\|_2 = \|f\|_2$ for any $n \geq 1$, we have by the above inequality that

$$\limsup_{n \rightarrow \infty} \frac{1}{\|f_n\|_2^2} \sum_{k \in \mathbb{Z}} |\langle f_n, u_k \rangle|^2 \leq 4\varepsilon.$$

Since ε is arbitrary, we have $\inf_{f \in V_m} \frac{1}{\|f\|_2^2} \sum_{k \in \mathbb{Z}} |\langle f, u_k \rangle|^2 = 0$ and hence

$$\inf_{f \in V_m} \frac{1}{\|f\|_2^2} \sum_{k \in \mathbb{Z}} \frac{x_{k+1} - x_{k-1}}{2} |\langle f, u_k \rangle|^2 = 0.$$

For the general case, suppose that $\frac{1}{r+1} \leq \beta < \frac{1}{r}$ for some integer $r \geq 1$. Let $x_{(r+1)k+l} = k + \frac{m-1}{2} + l\beta - \frac{\beta}{2}$, $0 \leq l \leq r$. Then $1 - r\beta \leq x_{k+1} - x_k \leq \beta$. Define $\varepsilon_k = \frac{1-\beta}{2+|k|}$. Choose averaging functions $u_k(x)$ such that

$$\begin{aligned}
 \text{supp } u_{(r+1)k+l} &\subset [k + \frac{m}{2} - \varepsilon_k, k + \frac{m}{2}], \quad 1 \leq l \leq \frac{r+1}{2}, \\
 \text{supp } u_{(r+1)k+l} &\subset [k + \frac{m}{2}, k + \frac{m}{2} + \varepsilon_k], \quad \frac{r+1}{2} < l \leq r+1.
 \end{aligned}$$

It is easy to check that $\text{supp } u_k \subset [x_k - \frac{\delta}{2}, x_k + \frac{\delta}{2}]$. Similar to the case $\frac{1}{2} \leq \beta < 1$ we can prove that

$$(3.9) \quad \inf_{f \in V_m} \frac{1}{\|f\|_2^2} \sum_{k \in \mathbb{Z}} \frac{x_{k+1} - x_{k-1}}{2} |\langle f, u_k \rangle|^2 = 0.$$

If there is a frame $\{S_k : k \in \mathbb{Z}\}$ for V_m such that $f = \sum_{k \in \mathbb{Z}} (\frac{x_{k+1} - x_{k-1}}{2})^{1/2} \langle f, u_k \rangle S_k$ holds for any $f \in V_m$, by [9, Proposition 3.2.4], we have

$$\sum_{k \in \mathbb{Z}} \frac{x_{k+1} - x_{k-1}}{2} |\langle f, u_k \rangle|^2 \geq \sum_{k \in \mathbb{Z}} |\langle f, \tilde{S}_k \rangle|^2 \geq \tilde{A} \|f\|_2^2,$$

where $\{\tilde{S}_k(x) : k \in \mathbb{Z}\}$ is the dual frame of $\{S_k(x) : k \in \mathbb{Z}\}$ and \tilde{A} is the lower frame bound. This is a contradiction to (3.9), which completes the proof. \square

4. ITERATIVE RECONSTRUCTION ALGORITHM

By Theorem 1.1, we can reconstruct any $f \in V_m$ from local averages by (1.1). However, it may be difficult to find the frame $\{S_k(x)\}$ in practice. In this case, the reconstruction of $f \in V_m$ from local averages can be performed by applying the following Feichtinger-Gröchenig algorithm.

Proposition 4.1 (see [13]). *Suppose that $\{\varphi_k : k \in \mathbb{Z}\}$ is a frame with bounds A and B for some Hilbert space \mathcal{H} . Let λ be a constant such that $0 < \lambda < \frac{2}{A+B}$. For any $f \in \mathcal{H}$, define*

$$\begin{aligned} Sf &= \lambda \sum_{k \in \mathbb{Z}} \langle f, \varphi_k \rangle \varphi_k, \\ f_0 &= Sf, \\ f_{k+1} &= f_k + S(f - f_k), \quad k \geq 0. \end{aligned}$$

Then $\lim_{k \rightarrow \infty} f_k = f$ and $\|f - f_k\| \leq \gamma^{k+1} \|f\|$, where $\gamma = \max\{|1 - \lambda A|, |1 - \lambda B|\} < 1$.

For Theorem 1.1, the operator S is of a slightly different form:

$$Sf := \lambda \sum_{k \in \mathbb{Z}} \frac{x_{k+1} - x_{k-1}}{2} \langle f, u_k \rangle h_k,$$

where h_k is defined as in Lemma 2.2.

In the above iterative reconstruction algorithm, the relaxation parameter λ plays an important role. If the exact value of the frame bounds are known, then $\lambda = \frac{2}{A+B}$ will be the best one since γ will be minimized in this case. Unfortunately, we have only some estimates for the frame bounds in most cases. Whereas it is not difficult to get an estimate for the upper bound of the frame $\{(\frac{x_{k+1} - x_{k-1}}{2})^{1/2} h_k : k \in \mathbb{Z}\}$ (see below), for the lower bound only existence is known. To ensure the convergence of the iterative reconstruction algorithm, we have to choose a small λ , which in turn may be responsible for slow convergence.

To apply Proposition 4.1, an upper frame bound must be found. Let $\{x_k\}$, δ , β , and $\{u_k\}$ be defined as in Theorem 1.1. Define $y_k = \frac{x_k + x_{k+1}}{2}$. We have

$$\begin{aligned} & \left\| f - \sum_{k \in \mathbb{Z}} \langle f, u_k \rangle \chi_{[y_{k-1}, y_k]} \right\|_2^2 = \sum_{k \in \mathbb{Z}} \int_{y_{k-1}}^{y_k} |f(x) - \langle f, u_k \rangle|^2 dx \\ &= \sum_{k \in \mathbb{Z}} \int_{y_{k-1}}^{y_k} \left| \int_{x_k - \frac{\delta}{2}}^{x_k + \frac{\delta}{2}} [f(x) - f(t)] u_k(t) dt \right|^2 dx \\ &\leq \sum_{k \in \mathbb{Z}} \int_{y_{k-1}}^{y_k} \int_{x_k - \frac{\delta}{2}}^{x_k + \frac{\delta}{2}} |f(x) - f(t)|^2 u_k(t) dt dx \\ &= \sum_{k \in \mathbb{Z}} \int_{y_{k-1}}^{y_k} \int_{x_k - \frac{\delta}{2}}^{x_k + \frac{\delta}{2}} \left| \int_0^{t-x} f'(x+s) ds \right|^2 u_k(t) dt dx \\ &\leq \sum_{k \in \mathbb{Z}} \int_{y_{k-1}}^{y_k} \int_{x_k - \frac{\delta}{2}}^{x_k + \frac{\delta}{2}} \frac{\delta + \beta}{2} \int_{-\frac{\delta+\beta}{2}}^{\frac{\delta+\beta}{2}} |f'(x+s)|^2 u_k(t) ds dt dx \\ &= \frac{\delta + \beta}{2} \int_{-\frac{\delta+\beta}{2}}^{\frac{\delta+\beta}{2}} \sum_{k \in \mathbb{Z}} \int_{y_{k-1}}^{y_k} |f'(x+s)|^2 dx ds = \frac{(\delta + \beta)^2}{2} \|f'\|_2^2. \end{aligned}$$

On the other hand, it can be shown (e.g., see [16]) that $\|f'\|_2^2 \leq \frac{M_m}{A_m} \|f\|_2^2$ with

$$\begin{aligned} M_m &= \sup_{|\omega| \leq \frac{\pi}{2}} \sum_{k \in \mathbb{Z}} |2\omega + 2k\pi|^2 \left| \frac{\sin \omega}{\omega + k\pi} \right|^{2m+2} \leq M_1 = 4, \\ A_m &= \inf_{|\omega| \leq \frac{\pi}{2}} \sum_{k \in \mathbb{Z}} \left| \frac{\sin \omega}{\omega + k\pi} \right|^{2m+2} = \left(\frac{2}{\pi} \right)^{2m+2} \sum_{k \in \mathbb{Z}} \frac{1}{(2k+1)^{2m+2}}. \end{aligned}$$

Hence

$$\sum_{k \in \mathbb{Z}} \frac{x_{k+1} - x_{k-1}}{2} |\langle f, u_k \rangle|^2 \leq \left(1 + \frac{(\delta + \beta) \sqrt{M_m}}{\sqrt{2A_m}} \right)^2 \|f\|_2^2 := B \|f\|_2^2.$$

Therefore, we can choose $0 < \lambda < \frac{2}{B}$ to ensure Proposition 4.1 works. However, we cannot give the convergence rate due to the absence of the lower frame bound.

REFERENCES

1. A.Aldroubi and K.Gröchenig, *Beurling-Landau-type theorems for non-uniform sampling in shift invariant spline spaces*, J. Fourier Anal. Appl., **6** (2000), 93–103. MR **2001i**:42057
2. A.Aldroubi and M.Unser, *Families of wavelet transforms in connection with Shannon's sampling theory and the Gabor transform*, Wavelets: A tutorial in Theory and Applications (C.K.Chui, Ed.), Academic Press, Boston, 1992, pp.509–528. MR **93a**:94003
3. A.Aldroubi and M.Unser, *Sampling procedures in function spaces and asymptotic equivalence with Shannon's sampling theory*, Numer. Funct. Anal. and Optimiz., **15** (1994), 1–21. MR **95a**:94002
4. J.J.Benedetto, *Irregular sampling and frames*, Wavelets: A Tutorial in Theory and Applications (C.K.Chui, Ed.), Academic Press, Boston, 1992, pp.445–507. MR **93c**:42030
5. P.L.Butzer and J.Lei, *Errors in truncated sampling series with measured sampled values for non-necessarily bandlimited functions*, Functiones et Approximation, **26** (1998), 25–39. MR **99m**:42003
6. P.L.Butzer and J.Lei, *Approximation of signals using measured sampled values and error analysis*, Comm. Appl. Anal., **4** (2000), 245–255. MR **2001i**:94033

7. W.Chen, S.Itoh, and J.Shiki, *Irregular sampling theorems for wavelet subspaces*, IEEE Trans. Inform. Theory, **44** (1998), 1131–1142. MR **99e**:94008
8. C.K.Chui, *An Introduction to Wavelets*, Academic Press, New York, 1992. MR **93f**:42055
9. I.Daubechies, *Ten Lectures on Wavelets*, SIAM, Philadelphia, 1992. MR **93e**:42045
10. H.Feichtinger and K.Gröchenig, *Iterative reconstruction of multivariate band-limited functions from irregular sampling values*, SIAM J. Math. Anal., **23** (1992), 244–261. MR **93a**:94006
11. H.Feichtinger and K.Gröchenig, *Irregular sampling theorems and series expansions of band-limited functions*, J. Math. Anal. Appl., **167** (1992), 530–556. MR **93g**:41003
12. H.Feichtinger and K.Gröchenig, *Error analysis in regular and irregular sampling theory*, Appl. Anal., **50** (1992), 167–189. MR **95b**:65160
13. H.Feichtinger and K.Gröchenig, *Theory and practice of irregular sampling*, Wavelets: Mathematics and Applications (J.Benedetto and M.Frazier, Eds.), CRC Press Inc., 1994, pp.305–363. MR **94i**:94008
14. K.Gröchenig, *Reconstruction algorithms in irregular sampling*, Math. Comput., **59** (1992), 181–194. MR **93a**:41025
15. A.J.E.M.Janssen, *The Zak transform and sampling theorem for wavelet subspaces*, IEEE Trans. Signal processing, **41** (1993), 3360–3364.
16. Y.Liu, *Irregular sampling for spline wavelet subspaces*, IEEE Trans. Inform. Theory, **42** (1996), 623–627. MR **97b**:94006
17. Y.Liu and G.Walter, *Irregular sampling in wavelet subspaces*, J. Fourier Anal. Appl., **2** (1995), 181–189. MR **96k**:42033
18. W.Sun and X.Zhou, *Frames and sampling theorem*, Science in China, Series A, **41** (1998), 606–612. MR **99m**:94015
19. W.Sun and X.Zhou, *Sampling theorem for wavelet subspaces: error estimate and irregular sampling*, IEEE Trans. Signal Processing, **48** (2000), 223–226.
20. W.Sun and X.Zhou, *Average sampling theorems for shift invariant subspaces*, Science in China, Series E, **43** (2000), 524–530. MR **2001m**:94030
21. W.Sun and X.Zhou, *Average sampling in spline subspaces*, Applied Mathematics Letters, **15**(2002), 233–237. MR **2003a**:94026
22. M.Unser and A.Aldroubi, *A general sampling theory for nonideal acquisition devices*, IEEE Trans. Signal Processing, **42** (1994), 2915–2925.
23. G.Walter, *A sampling theorem for wavelet subspaces*, IEEE Trans. Inform. Theory, **38** (1992), 881–884. MR **93e**:94006
24. R.G.Wiley, *Recovery of band-limited signals from unequally spaced samples*, IEEE Trans. Comm., **26** (1978), 135–137.
25. X.G.Xia and Z.Zhang, *On sampling theorem, wavelets, and wavelet transforms*, IEEE Transactions on Signal Processing, **41** (1993), 2524–2535.
26. R.M.Young, *An Introduction to Non-Harmonic Fourier Series*, Academic, New York, 1980. MR **81m**:42027
27. X.Zhou, W.Sun, *On the sampling theorem for wavelet subspaces*, J. Fourier Anal. Appl., **5** (1999), 347–354. MR **2001i**:42025

DEPARTMENT OF MATHEMATICS, NANKAI UNIVERSITY, TIANJIN 300071, PEOPLE'S REPUBLIC OF CHINA

E-mail address: sunwch@nankai.edu.cn

DEPARTMENT OF MATHEMATICS, NANKAI UNIVERSITY, TIANJIN 300071, PEOPLE'S REPUBLIC OF CHINA

E-mail address: xwzhou@nankai.edu.cn