

PARTIAL SUMS OF HYPERGEOMETRIC SERIES OF UNIT ARGUMENT

WOLFGANG BÜHRING

(Communicated by Carmen C. Chicone)

ABSTRACT. The asymptotic behaviour of partial sums of generalized hypergeometric series of unit argument is investigated.

1. INTRODUCTION

This paper deals with finite sums

$$(1) \quad \sum_{l=0}^{m-1} \frac{\Gamma(a_1+l)\Gamma(a_2+l)\cdots\Gamma(a_{p+1}+l)}{\Gamma(b_1+l)\cdots\Gamma(b_p+l)\Gamma(1+l)}$$

which are partial sums of the first m terms of hypergeometric series ${}_{p+1}F_p$ of unit argument $z = 1$ multiplied by appropriate gamma function factors,

$$(2) \quad \frac{\Gamma(a_1)\Gamma(a_2)\cdots\Gamma(a_{p+1})}{\Gamma(b_1)\cdots\Gamma(b_p)} {}_{p+1}F_p \left(\begin{matrix} a_1, a_2, \dots, a_{p+1} \\ b_1, \dots, b_p \end{matrix} \middle| z \right) \\ = \sum_{l=0}^{\infty} \frac{\Gamma(a_1+l)\Gamma(a_2+l)\cdots\Gamma(a_{p+1}+l)}{\Gamma(b_1+l)\cdots\Gamma(b_p+l)\Gamma(1+l)} z^l.$$

An important quantity in this context is

$$(3) \quad s_p = b_1 + \cdots + b_p - a_1 - a_2 - \cdots - a_{p+1},$$

the non-trivial characteristic exponent of the underlying differential equation at $z = 1$.

If $\Re(s_p) > 0$, then the hypergeometric series converges at $z = 1$, and its value is given by the Gaussian summation formula for $p = 1$ or by the corresponding generalized formulas [5] for $p > 1$. The terms of the series behave asymptotically as

$$l^{a_1+a_2+\cdots+a_{p+1}-b_1-\cdots-b_p-1} = l^{-s_p-1}$$

for $l \rightarrow \infty$, and if we consider the partial sum of the first m terms, then the contribution of the missing tail is $O(m^{-s_p})$ as $m \rightarrow \infty$. Therefore we have

Received by the editors June 28, 2002 and, in revised form, September 25, 2002.

2000 *Mathematics Subject Classification*. Primary 33C20.

Key words and phrases. Partial sums, generalized hypergeometric series.

Theorem 1. *If $\Re(s_1) > 0$, then*

$$(4) \quad \sum_{l=0}^{m-1} \frac{\Gamma(a_1+l)\Gamma(a_2+l)}{\Gamma(b_1+l)\Gamma(1+l)} = \frac{\Gamma(a_1)\Gamma(a_2)\Gamma(s_1)}{\Gamma(a_1+s_1)\Gamma(a_2+s_1)} + O(m^{-s_1})$$

as $m \rightarrow \infty$.

Or more generally,

Theorem 2. *If $\Re(s_p) > 0$, then*

$$(5) \quad \sum_{l=0}^{m-1} \frac{\Gamma(a_1+l)\Gamma(a_2+l)\cdots\Gamma(a_{p+1}+l)}{\Gamma(b_1+l)\cdots\Gamma(b_p+l)\Gamma(1+l)} = g_0(0) + O(m^{-s_p})$$

as $m \rightarrow \infty$, where

$$(6) \quad g_0(0) = \frac{\Gamma(a_1)\Gamma(a_2)\Gamma(s_p)}{\Gamma(a_1+s_p)\Gamma(a_2+s_p)} \sum_{k=0}^{\infty} \frac{(s_p)_k}{(a_1+s_p)_k(a_2+s_p)_k} A_k^{(p)}$$

and the sum converges provided that $\Re(a_j) > 0$ for $j = 3, \dots, p+1$.

Here use is made of the Pochhammer symbol

$$(x)_n = x(x+1)\cdots(x+n-1) = \Gamma(x+n)/\Gamma(x),$$

and the coefficients $A_k^{(p)}$ for $p = 2, 3, \dots$ are given in [5], but a few of them are displayed here again for convenience:

$$(7) \quad A_k^{(2)} = \frac{(b_2 - a_3)_k (b_1 - a_3)_k}{k!},$$

$$(8) \quad A_k^{(3)} = \sum_{k_2=0}^k \frac{(b_3 + b_2 - a_4 - a_3 + k_2)_{k-k_2} (b_1 - a_3)_{k-k_2} (b_3 - a_4)_{k_2} (b_2 - a_4)_{k_2}}{(k - k_2)! k_2!},$$

$$(9) \quad A_k^{(4)} = \sum_{k_2=0}^k \frac{(b_4 + b_3 + b_2 - a_5 - a_4 - a_3 + k_2)_{k-k_2} (b_1 - a_3)_{k-k_2}}{(k - k_2)!} \\ \times \sum_{k_3=0}^{k_2} \frac{(b_4 + b_3 - a_5 - a_4 + k_3)_{k_2-k_3} (b_2 - a_4)_{k_2-k_3}}{(k_2 - k_3)!} \\ \times \frac{(b_4 - a_5)_{k_3} (b_3 - a_5)_{k_3}}{k_3!}.$$

For $p = 3, 4, \dots$ several other representations are possible [5].

In the case $p = 2$, the sum in (6) is an ${}_3F_2$ and so (5)–(6) reduce, when $m \rightarrow \infty$, to a well-known transformation formula for ${}_3F_2$ of unit argument [1], [10], [14].

The simple formula (4) above corresponding to $p = 1$ can be recovered from the general formula (5)–(6) if we define

$$(10) \quad A_0^{(1)} = 1, \quad A_k^{(1)} = 0 \quad \text{for } k > 0,$$

so that then the sum over k is equal to 1 and disappears. In a similar way the formulas in the other theorems below simplify for $p = 1$.

It is the purpose of this work to investigate the asymptotic behaviour of the partial sums (1) for situations when $\Re(s_p) \leq 0$ and to find a more detailed formula in cases like $s_p = 1$, when a few additional terms of higher order might be desirable.

2. THE CASE WHEN s_p IS NOT EQUAL TO AN INTEGER

If $\Re(s_p) \leq 0$, then the hypergeometric series diverges at $z = 1$, and the contribution of the missing tail of the series does not asymptotically vanish. Responsible for this behaviour are the singular terms of the hypergeometric function at $z = 1$. We have to explicitly subtract the contribution from the most important singular terms in order to get the tail to vanish as $m \rightarrow \infty$. The behaviour of the hypergeometric function at $z = 1$ is given by the continuation formula

$$(11) \quad \frac{\Gamma(a_1)\Gamma(a_2)\cdots\Gamma(a_{p+1})}{\Gamma(b_1)\cdots\Gamma(b_p)} {}_{p+1}F_p \left(\begin{matrix} a_1, a_2, \dots, a_{p+1} \\ b_1, \dots, b_p \end{matrix} \middle| z \right) \\ = \sum_{n=0}^{\infty} g_n(0) (1-z)^n + (1-z)^{s_p} \sum_{n=0}^{\infty} g_n(s_p) (1-z)^n,$$

which is valid provided that s_p is not equal to an integer. Here the coefficient $g_0(0)$ defined in (6) above enters, and, while the other $g_n(0)$ are not needed here, the coefficients $g_n(s_p)$ of the singular term are [5]

$$(12) \quad g_n(s_p) = (-1)^n \frac{(a_1 + s_p)_n (a_2 + s_p)_n \Gamma(-s_p - n)}{(1)_n} \\ \times \sum_{k=0}^n \frac{(-n)_k}{(a_1 + s_p)_k (a_2 + s_p)_k} A_k^{(p)}.$$

The singular terms on the right-hand side of (11) have, with $x = s_p + n$, the z -dependence

$$(13) \quad (1-z)^x = \sum_{l=0}^{\infty} \frac{(-x)_l}{\Gamma(1+l)} z^l.$$

With $z = 1$, we shall need the partial sum of this series,

$$(14) \quad \sum_{l=0}^{m-1} \frac{(-x)_l}{\Gamma(1+l)} = -\frac{1}{x} \frac{(-x)_m}{\Gamma(m)} \quad (m = 1, 2, \dots \quad x \neq 0).$$

Treating the first $N + 1$ singular terms on the right-hand side of (11) in this way, we may obtain

$$(15) \quad \sum_{l=0}^{m-1} \frac{\Gamma(a_1+l)\Gamma(a_2+l)\cdots\Gamma(a_{p+1}+l)}{\Gamma(b_1+l)\cdots\Gamma(b_p+l)\Gamma(1+l)} = g_0(0) \\ - \sum_{n=0}^N g_n(s_p) \frac{1}{s_p+n} \frac{(-s_p-n)_m}{\Gamma(m)} + T_1 + T_2,$$

where

$$(16) \quad T_1 = - \sum_{l=m}^{\infty} \frac{\Gamma(a_1+l)\Gamma(a_2+l)\cdots\Gamma(a_{p+1}+l)}{\Gamma(b_1+l)\cdots\Gamma(b_p+l)\Gamma(1+l)} \\ + \sum_{n=0}^N g_n(s_p) \sum_{l=m}^{\infty} \frac{(-s_p-n)_l}{\Gamma(1+l)},$$

$$(17) \quad T_2 = - \lim_{m \rightarrow \infty} \sum_{n=N+1}^{\infty} \frac{g_n(s_p)}{(s_p+n)\Gamma(-s_p-n)} \frac{\Gamma(-s_p-n+m)}{\Gamma(m)}.$$

We now have to show that the contributions T_1 and T_2 of the tails of the series are small. The terms in the series in T_2 are $O(m^{-s_p-n})$ with $n \geq N+1$ as $m \rightarrow \infty$, and so the series, in the worst case when $n = N+1$, converges provided that $\Re(-s_p - N) < 0$. The sum of the series then is $O(m^{-s_p-N})$, and T_2 exists and vanishes. The term T_1 may be written

$$(18) \quad T_1 = - \sum_{l=m}^{\infty} \frac{\Gamma(-s_p+l)}{\Gamma(1+l)} S_l,$$

where

$$(19) \quad S_l = \frac{\Gamma(a_1+l)\Gamma(a_2+l)\cdots\Gamma(a_{p+1}+l)}{\Gamma(b_1+l)\cdots\Gamma(b_p+l)\Gamma(-s_p+l)} \\ - \sum_{n=0}^N (-1)^n g_n(s_p) \frac{1}{\Gamma(-s_p-n)(1+s_p-l)_n}.$$

In the second term, use has been made of the reflection formula of the gamma function. It can be shown [6] that S_l is $O(l^{-N-1})$ as $l \rightarrow \infty$, so the terms of the sum are $O(l^{-s_p-N-2})$, and T_1 is $O(m^{-s_p-N-1})$ as $m \rightarrow \infty$. In this way we arrive at

Theorem 3. *If s_p is not equal to an integer and $N = 0, 1, 2, \dots$ is chosen to be greater than $\Re(-s_p)$, then*

$$(20) \quad \sum_{l=0}^{m-1} \frac{\Gamma(a_1+l)\Gamma(a_2+l)\cdots\Gamma(a_{p+1}+l)}{\Gamma(b_1+l)\cdots\Gamma(b_p+l)\Gamma(1+l)} = g_0(0) \\ - \sum_{n=0}^N (-1)^n \frac{1}{s_p+n} \frac{(a_1+s_p)_n (a_2+s_p)_n}{(1)_n} \frac{\Gamma(-s_p-n+m)}{\Gamma(m)} \\ \times \sum_{k=0}^n \frac{(-n)_k}{(a_1+s_p)_k (a_2+s_p)_k} A_k^{(p)} + O(m^{-s_p-N-1})$$

as $m \rightarrow \infty$.

This theorem is applicable in the case of $\Re(s_p) > 0$, too, and then may give more details of the asymptotic behaviour than Theorem 2.

3. THE CASE WHEN s_p IS EQUAL TO AN INTEGER

If s_p is equal to an integer, then the procedure of proof remains essentially the same, except that the starting point is a more complicated continuation formula, in place of (11), containing logarithmic terms which have to be expanded, too.

3.1. The case when s_p is equal to zero. If s_p is equal to zero, the hypergeometric series is called zero-balanced. It is this case which seems to be most interesting. While a few early results can be found in the monographs [1], [14], [10], the topic had received renewed attention, with emphasis on the zero-balanced series, in connection with Ramanujan's notebooks [11], [2], [3], [9], [8], [4], [15], [12], [13], [7].

The required continuation formula for the zero-balanced hypergeometric function is [5]

$$(21) \quad \frac{\Gamma(a_1)\Gamma(a_2)\cdots\Gamma(a_{p+1})}{\Gamma(b_1)\cdots\Gamma(b_p)} {}_{p+1}F_p \left(\begin{matrix} a_1, a_2, \dots, a_{p+1} \\ b_1, \dots, b_p \end{matrix} \middle| z \right) \\ = \sum_{n=0}^{\infty} d_n(1-z)^n + \sum_{n=0}^{\infty} e_n(1-z)^n \ln(1-z),$$

where

$$(22) \quad e_n = -\frac{(a_1)_n(a_2)_n}{(1)_n(1)_n} \sum_{k=0}^n \frac{(-n)_k}{(a_1)_k(a_2)_k} A_k^{(p)},$$

$$(23) \quad d_0 = 2\psi(1) - \psi(a_1) - \psi(a_2) + \sum_{k=1}^{\infty} \frac{\Gamma(k)}{(a_1)_k(a_2)_k} A_k^{(p)},$$

and the other coefficients are not needed here. Again, we have to expand the singular terms on the right-hand side,

$$(24) \quad (1-z)^n \ln(1-z) = \sum_{l=1}^{\infty} c_l^{(n)} z^l,$$

where

$$(25) \quad c_l^{(n)} = -\frac{1}{l}(-1)^n \frac{\Gamma(1+n)\Gamma(l-n)}{\Gamma(l)}$$

for $l > n$, while the coefficients for $l \leq n$ are

$$(26) \quad c_1^{(1)} = -1, \quad c_1^{(2)} = -1, \quad c_2^{(2)} = \frac{3}{2}.$$

With $z = 1$, we need the partial sums of these series,

$$(27) \quad \sum_{l=1}^{m-1} c_l^{(0)} = \psi(1) - \psi(m),$$

$$(28) \quad \sum_{l=1}^{m-1} c_l^{(1)} = -\frac{1}{m-1},$$

$$(29) \quad \sum_{l=1}^{m-1} c_l^{(2)} = \frac{1}{(m-1)(m-2)}.$$

Keeping the contribution from the singular terms up to and including $n = 2$ together with the constant term on the right-hand side of (21) we arrive at

Theorem 4. *If s_p is equal to zero, then*

$$(30) \quad \sum_{l=0}^{m-1} \frac{\Gamma(a_1+l)\Gamma(a_2+l)\cdots\Gamma(a_{p+1}+l)}{\Gamma(b_1+l)\cdots\Gamma(b_p+l)\Gamma(1+l)} = \sum_{k=1}^{\infty} \frac{\Gamma(k)}{(a_1)_k(a_2)_k} A_k^{(p)} \\ + \psi(1) - \psi(a_1) - \psi(a_2) + \psi(m) + [a_1 a_2 - A_1^{(p)}](m-1)^{-1} \\ - \frac{1}{4}[a_1(a_1+1)a_2(a_2+1) - 2(a_1+1)(a_2+1)A_1^{(p)} + 2A_2^{(p)}] \\ \times [(m-1)(m-2)]^{-1} + O(m^{-3})$$

as $m \rightarrow \infty$, where the infinite sum over k converges if $\Re(a_j) > 0$ for $j = 3, \dots, p+1$.

This theorem gives more details of the asymptotic behaviour than our earlier formula [7]. If desired, equation (30) may be rewritten using

$$(31) \quad \psi(m) = \ln(m) - \frac{1}{2}m^{-1} - \frac{1}{12}m^{-2} + O(m^{-3})$$

in order to exhibit the logarithmic dependence on m .

3.2. The case when s_p is equal to a positive integer. For s_p equal to a positive integer t , Theorems 1 or 2 are applicable, but for s_p equal to 1 or 2 a few more terms of the asymptotic expansion might be desirable. In these cases the required continuation formula reads [5]

$$(32) \quad \frac{\Gamma(a_1)\Gamma(a_2)\cdots\Gamma(a_{p+1})}{\Gamma(b_1)\cdots\Gamma(b_p)} {}_{p+1}F_p \left(\begin{matrix} a_1, a_2, \dots, a_{p+1} \\ b_1, \dots, b_p \end{matrix} \middle| z \right) \\ = \sum_{n=0}^{t-1} l_n (1-z)^n + (1-z)^t \sum_{n=0}^{\infty} [w_n + q_n \ln(1-z)] (1-z)^n,$$

where

$$(33) \quad q_n = -(-1)^t \frac{(a_1+t)_n (a_2+t)_n}{\Gamma(1+t+n)} \sum_{k=0}^n \frac{(-n)_k}{(a_1+t)_k (a_2+t)_k} A_k^{(p)},$$

$$(34) \quad l_0 = \frac{\Gamma(a_1)\Gamma(a_2)\Gamma(t)}{\Gamma(a_1+t)\Gamma(a_2+t)} \sum_{k=0}^{\infty} \frac{(t)_k}{(a_1+t)_k (a_2+t)_k} A_k^{(p)},$$

and the other coefficients are not needed here. Using the expansions of the logarithmic terms and their partial sums from above and keeping the contributions up to and including $n = 1$ in case of $t = 1$ or $n = 0$ in case of $t = 2$, we may arrive at the following theorems.

Theorem 5. *If s_p is equal to 1, then*

$$(35) \quad \sum_{l=0}^{m-1} \frac{\Gamma(a_1+l)\Gamma(a_2+l)\cdots\Gamma(a_{p+1}+l)}{\Gamma(b_1+l)\cdots\Gamma(b_p+l)\Gamma(1+l)} \\ = \frac{1}{a_1 a_2} \sum_{k=0}^{\infty} \frac{(1)_k}{(a_1+1)_k (a_2+1)_k} A_k^{(p)} - (m-1)^{-1} \\ + \frac{1}{2} [(a_1+1)(a_2+1) - A_1^{(p)}] [(m-1)(m-2)]^{-1} + O(m^{-3})$$

as $m \rightarrow \infty$, where the infinite sum converges if $\Re(a_j) > 0$ for $j = 3, \dots, p+1$.

Theorem 6. *If s_p is equal to 2, then*

$$(36) \quad \sum_{l=0}^{m-1} \frac{\Gamma(a_1+l)\Gamma(a_2+l)\cdots\Gamma(a_{p+1}+l)}{\Gamma(b_1+l)\cdots\Gamma(b_p+l)\Gamma(1+l)} \\ = \frac{1}{(a_1)_2 (a_2)_2} \sum_{k=0}^{\infty} \frac{(2)_k}{(a_1+2)_k (a_2+2)_k} A_k^{(p)} - \frac{1}{2} [(m-1)(m-2)]^{-1} + O(m^{-3})$$

as $m \rightarrow \infty$, where the infinite sum converges if $\Re(a_j) > 0$ for $j = 3, \dots, p+1$.

We may observe that Theorem 5 and Theorem 6 are not unexpected, but (35) and (36) are special cases of (20). Indeed, all the quantities in (20) remain well-defined when s_p approaches a positive integer, and so Theorem 3 is valid even for positive integer values of s_p .

3.3. The case when s_p is equal to a negative integer. When s_p is equal to a negative integer $-t$, then the required continuation formula is [5]

$$(37) \quad \frac{\Gamma(a_1)\Gamma(a_2)\cdots\Gamma(a_{p+1})}{\Gamma(b_1)\cdots\Gamma(b_p)} {}_{p+1}F_p \left(\begin{matrix} a_1, a_2, \dots, a_{p+1} \\ b_1, \dots, b_p \end{matrix} \middle| z \right) \\ = (1-z)^{-t} \sum_{n=0}^{t-1} h_n (1-z)^n + \sum_{n=0}^{\infty} [u_n + v_n \ln(1-z)] (1-z)^n,$$

where

$$(38) \quad h_n = (-1)^n \frac{(a_1-t)_n (a_2-t)_n \Gamma(t-n)}{\Gamma(1+n)} \sum_{k=0}^n \frac{(-n)_k}{(a_1-t)_k (a_2-t)_k} A_k^{(p)},$$

$$(39) \quad v_n = -(-1)^t \frac{(a_1-t)_{t+n} (a_2-t)_{t+n}}{\Gamma(1+n)\Gamma(1+t+n)} \sum_{k=0}^{t+n} \frac{(-t-n)_k}{(a_1-t)_k (a_2-t)_k} A_k^{(p)},$$

$$(40) \quad u_0 = \sum_{k=1}^{\infty} \frac{\Gamma(k)}{(a_1)_k (a_2)_k} A_{t+k}^{(p)} \\ + (-1)^t \frac{(a_1-t)_t (a_2-t)_t}{\Gamma(1+t)} \sum_{k=0}^t \frac{(-t)_k}{(a_1-t)_k (a_2-t)_k} A_k^{(p)} \\ \times [\psi(1+t-k) + \psi(1) - \psi(a_1) - \psi(a_2)],$$

and the other u_n are not needed here. Proceeding as above and keeping the contributions from the logarithmic terms up to and including $n = 2$, we then may get

Theorem 7. *If s_p is equal to a negative integer $-t$, then*

$$(41) \quad \sum_{l=0}^{m-1} \frac{\Gamma(a_1+l)\Gamma(a_2+l)\cdots\Gamma(a_{p+1}+l)}{\Gamma(b_1+l)\cdots\Gamma(b_p+l)\Gamma(1+l)} \\ = \sum_{n=0}^{t-1} (-1)^n \frac{1}{t-n} \frac{(a_1-t)_n (a_2-t)_n \Gamma(t-n)}{\Gamma(1+n)} \frac{\Gamma(t-n+m)}{\Gamma(m)} \\ \times \sum_{k=0}^n \frac{(-n)_k}{(a_1-t)_k (a_2-t)_k} A_k^{(p)} \\ + \sum_{k=1}^{\infty} \frac{\Gamma(k)}{(a_1)_k (a_2)_k} A_{t+k}^{(p)} \\ + (-1)^t \frac{(a_1-t)_t (a_2-t)_t}{\Gamma(1+t)} \sum_{k=0}^t \frac{(-t)_k}{(a_1-t)_k (a_2-t)_k} A_k^{(p)}$$

$$\begin{aligned}
& \times [\psi(1+t-k) - \psi(a_1) - \psi(a_2) + \psi(m)] \\
& + (-1)^t \frac{(a_1-t)_{t+1}(a_2-t)_{t+1}}{\Gamma(2+t)} \sum_{k=0}^{t+1} \frac{(-t-1)_k}{(a_1-t)_k(a_2-t)_k} A_k^{(p)} (m-1)^{-1} \\
& \quad - (-1)^t \frac{(a_1-t)_{t+2}(a_2-t)_{t+2}}{2\Gamma(3+t)} \\
& \times \sum_{k=0}^{t+2} \frac{(-t-2)_k}{(a_1-t)_k(a_2-t)_k} A_k^{(p)} [(m-1)(m-2)]^{-1} + O(m^{-3})
\end{aligned}$$

as $m \rightarrow \infty$, where the infinite sum converges if $\Re(a_j) > 0$ for $j = 3, \dots, p+1$.

4. SPECIAL EXAMPLES

If the parameters $a_1, \dots, a_{p+1}, b_1, \dots, b_p$ satisfy certain additional relations or even have special values, it may well be that the appropriate formula above can be simplified, in particular its constant term, which involves an infinite sum.

As our first example of this kind, let us consider a zero-balanced ${}_5F_4$ with the special values of the parameters $a_1 = a_2 = a_3 = a_4 = 1/2$, $a_5 = 5/4$, $b_1 = b_2 = b_3 = 1$, $b_4 = 1/4$. More convenient than (9) for this purpose is the alternative representation (4.2) of [5]. The constant term CT of (30) can then be reduced to

$$(42) \quad CT = -\psi(1) + 4 \ln(2) - \frac{1}{18} \sum_{l=0}^{\infty} (1+l) \frac{(2)_l(2)_l}{(\frac{5}{2})_l(\frac{5}{2})_l} {}_3F_2 \left(\begin{matrix} 3/2, 3/2, -l \\ 2, 3 \end{matrix} \middle| 1 \right).$$

Here the infinite sum seems to be difficult to evaluate analytically, although it must simply be equal to $18 \ln(2)$. For we know from Berndt [3] that

$$(43) \quad CT = -\psi(1) + 3 \ln(2).$$

Using this information, we obtain from (30)

Corollary 1.

$$(44) \quad \frac{1}{4} \pi^2 \sum_{l=0}^{m-1} \frac{(\frac{1}{2})_l(\frac{1}{2})_l(\frac{1}{2})_l(\frac{1}{2})_l(\frac{5}{4})_l}{(1)_l(1)_l(1)_l(1)_l(\frac{1}{4})_l} = \psi(m) - \psi(1) + 3 \ln(2) \\ + \frac{1}{4}(m-1)^{-1} - \frac{1}{8}[(m-1)(m-2)]^{-2} + O(m^{-3}).$$

Our second example is a zero-balanced ${}_5F_4$ which depends only on three independent parameters a, b, c according to $a_1 = a$, $a_2 = b$, $a_3 = c$, $a_4 = a + b + c - 1$, $a_5 = (a + b + c + 1)/2$, $b_1 = b + c$, $b_2 = c + a$, $b_3 = a + b$, $b_4 = (a + b + c - 1)/2$. Here again, the infinite sum in the constant term of (30) is difficult to evaluate analytically, but fortunately we know the constant term from a different source [8], [3], [11]. Thus we obtain

Corollary 2.

$$(45) \quad \sum_{l=0}^{m-1} \frac{\Gamma(a+l)\Gamma(b+l)\Gamma(c+l)\Gamma(a+b+c-1+l)\Gamma(\frac{1}{2}[a+b+c+1]+l)}{\Gamma(b+c+l)\Gamma(c+a+l)\Gamma(a+b+l)\Gamma(\frac{1}{2}[a+b+c-1]+l)\Gamma(1+l)} \\ = \psi(m) + \frac{1}{2}[\psi(1) - \psi(a) - \psi(b) - \psi(c)] + \frac{1}{2}(a+b+c-1)(m-1)^{-1} \\ + \frac{1}{4}[2abc - (a+b+c)(a+b+c-1)][(m-1)(m-2)]^{-2} + O(m^{-3}).$$

As a special case of this formula, with $a = b = c = 1/2$, we can again get Corollary 1 above.

Other partial sums with interrelated parameters have been obtained by A. K. Srivastava [15].

REFERENCES

1. W. N. Bailey, *Generalized hypergeometric series*, Cambridge Tracts in Mathematics and Mathematical Physics, No. 32, Stechert-Hafner, New York, 1964. MR **32**:2625
2. B. C. Berndt, *Chapter 11 of Ramanujan's second notebook*, Bull. London Math. Soc. **15** (1983), 273–320. MR **85a**:01043
3. B. C. Berndt, *Ramanujan's notebooks*, Part II, Springer-Verlag, New York, 1989. MR **90b**:01039
4. W. Bühring, *The behavior at unit argument of the hypergeometric function ${}_3F_2$* , SIAM J. Math. Anal. **18** (1987), 1227–1234. MR **88j**:33004
5. W. Bühring, *Generalized hypergeometric functions at unit argument*, Proc. Amer. Math. Soc. **114** (1992), 145–153. MR **92c**:33004
6. W. Bühring, *Asymptotic expansions for ratios of products of gamma functions*, Internat. J. Math. Math. Sci. **2003**:**18** (2003), 1167–1171.
7. W. Bühring and H. M. Srivastava, *Analytic continuation of the generalized hypergeometric series near unit argument with emphasis on the zero-balanced series*, Approximation Theory and Applications (Th. M. Rassias, ed.), Hadronic Press, Palm Harbor, 1998, pp. 17–35 (updated version: e-print math.CA/0102032 on <http://arXiv.org>).
8. R. J. Evans, *Ramanujan's second notebook: Asymptotic expansions for hypergeometric series and related functions*, Ramanujan Revisited (G. E. Andrews et al., eds.), Academic Press, Boston, MA, 1988, pp. 537–560. MR **89c**:33003
9. R. J. Evans and D. Stanton, *Asymptotic formulas for zero-balanced hypergeometric series*, SIAM J. Math. Anal. **15** (1984), 1010–1020. MR **85i**:33003
10. Y. L. Luke, *The special functions and their approximations*, vol. 1, Academic Press, New York, 1969. MR **39**:3039
11. S. Ramanujan, *Notebooks*, vol. 2, Tata Institute of Fundamental Research, Bombay, 1957. MR **20**:6340
12. M. Saigo, *On properties of the Appell hypergeometric functions F_2 and F_3 and the generalized Gauss function ${}_3F_2$* , Bull. Central Research Institute Fukuoka University **66** (1983), 27–32. MR **85m**:33006
13. M. Saigo and H. M. Srivastava, *The behavior of the zero-balanced hypergeometric series ${}_pF_{p-1}$ near the boundary of its convergence region*, Proc. Amer. Math. Soc. **110** (1990), 71–76. MR **91d**:33005
14. L. J. Slater, *Generalized hypergeometric functions*, Cambridge Univ. Press, Cambridge, 1966. MR **34**:1570
15. A. K. Srivastava, *Asymptotic behaviour of certain zero-balanced hypergeometric series*, Proc. Indian Acad. Sci. (Math. Sci.) **106** (1996), 39–51. MR **97h**:33013

PHYSIKALISCHES INSTITUT, UNIVERSITÄT HEIDELBERG, PHILOSOPHENWEG 12, 69120 HEIDELBERG, GERMANY

E-mail address: buehring@physi.uni-heidelberg.de