

A PROOF OF W. T. GOWERS' c_0 THEOREM

VASSILIS KANELLOPOULOS

(Communicated by N. Tomczak-Jaegermann)

ABSTRACT. W. T. Gowers' c_0 theorem asserts that for every Lipschitz function $F : S_{c_0} \rightarrow \mathbb{R}$ and $\varepsilon > 0$, there exists an infinite-dimensional subspace Y of c_0 such that the oscillation of F on S_Y is at most ε . The proof of this theorem has been reduced by W. T. Gowers to the proof of a new Ramsey type theorem. Our aim is to present a proof of the last result.

1. INTRODUCTION AND NOTATION

In this paper we give a proof of W. T. Gowers' c_0 theorem (see [G1], [BL] and [G2]). As W. T. Gowers has shown, the theorem is essentially purely combinatorial and reduces to a Ramsey type result on the space of finitely supported functions $f : \mathbb{N} \rightarrow \{-k, \dots, k\}$. The proof presented here is based on an abstract lemma, implicitly contained in Gowers' proof and which may be considered as an extension of the well-known lemma of the existence of an idempotent in a compact semigroup. Both the positive and the general case of the theorem are naturally derived from it. At the end of the paper we briefly describe some Hales-Jewett versions of the above combinatorial results.

Notation. Let us fix some notation. By $\mathbb{N} = \{0, 1, 2, \dots\}$ we denote the set of all non-negative integers. For $k \geq 1$ we define

$$X_k = \{f : \mathbb{N} \rightarrow \{0, \dots, k\} : \text{supp } f \text{ is finite and } \exists n \in \mathbb{N} \text{ with } f(n) = k\} \text{ and} \\ X_{\pm k} = \{f : \mathbb{N} \rightarrow \{-k, \dots, k\} : \text{supp } f \text{ is finite and } \exists n \in \mathbb{N} \text{ with } |f(n)| = k\}.$$

By θ we denote the constant zero map $\theta : \mathbb{N} \rightarrow \{0\}$ and we set

$$X_{[k]} = \bigcup_{i=0}^k X_i \quad \text{and} \quad X_{[\pm k]} = \bigcup_{i=0}^k X_{\pm i}, \quad \text{where } X_0 = \{\theta\}.$$

For $f_1, f_2 \in X_{[\pm k]} \setminus X_0$, we write $f_1 < f_2$ if $\max \text{supp } f_1 < \min \text{supp } f_2$. A sequence $\vec{f} = (f_n)_n$ in $X_{[\pm k]}$ is called a block sequence if $f_n < f_{n+1}$ for every n .

Received by the editors February 26, 2003 and, in revised form, March 23, 2003 and June 13, 2003.

2000 *Mathematics Subject Classification.* Primary 46B45, 46T20.

Key words and phrases. Lipschitz functions, compact semigroups, idempotents, ultrafilters, variable words.

Partially supported by Thales program of NTUA.

Let $T : X_{[\pm k]} \rightarrow X_{[\pm k]}$ be defined by

$$Tf(n) = \begin{cases} f(n) - 1 & \text{if } f(n) > 0, \\ f(n) + 1 & \text{if } f(n) < 0, \\ 0 & \text{if } f(n) = 0. \end{cases}$$

For a block sequence $\vec{f} = (f_n)_n$ in X_k the *span* of \vec{f} in X_k is the set

$$\langle \vec{f} \rangle_k = \{ T^{\epsilon_0} f_{n_0} + \dots + T^{\epsilon_m} f_{n_m} : n_0 < \dots < n_m, \\ \epsilon_0, \dots, \epsilon_m \in \{0, \dots, k-1\} \\ \text{and } \exists i \in \{0, \dots, m\} \text{ with } \epsilon_i = 0 \},$$

and for a block sequence $\vec{f} = (f_n)_n$ in $X_{\pm k}$ the *span* of \vec{f} in $X_{\pm k}$ is the set

$$\langle \vec{f} \rangle_{\pm k} = \{ \pm T^{\epsilon_0} f_{n_0} \pm \dots \pm T^{\epsilon_m} f_{n_m} : n_0 < \dots < n_m, \\ \epsilon_0, \dots, \epsilon_m \in \{0, \dots, k-1\} \\ \text{and } \exists i \in \{0, \dots, m\} \text{ with } \epsilon_i = 0 \}.$$

A *coloring* of a set X is a partition of X into disjoint pieces, and a subset of X is called *monochromatic* if it is contained in one of these pieces. Under the above notation the positive case of W. T. Gowers' theorem is stated as follows.

Theorem 1. *For every finite coloring of X_k , $k \geq 1$, there exists an infinite block sequence \vec{f} in X_k such that $\langle \vec{f} \rangle_k$ is monochromatic.*

For the general case we need one more definition. Two elements f_1, f_2 of $X_{[\pm k]}$ are called *neighbours* if $\|f_1 - f_2\|_\infty \leq 1$. For a set $C \subseteq X_{[\pm k]}$, by \widehat{C} we denote the set of all neighbours of elements of C , i.e. $\widehat{C} = \{f \in X_{[\pm k]} : \exists g \in C \text{ such that } \|f - g\|_\infty \leq 1\}$.

Finally, if $X_{[\pm k]} = C_0 \cup \dots \cup C_p$ is a finite coloring of $X_{[\pm k]}$, then a set $A \subseteq X_{[\pm k]}$ is called *approximately monochromatic* if there exists an i such that $A \subseteq \widehat{C}_i$. The general and most interesting case of W. T. Gowers' theorem is the following.

Theorem 2. *For every finite coloring of $X_{\pm k}$, $k \geq 1$, there exists an infinite block sequence \vec{f} in $X_{\pm k}$ such that $\langle \vec{f} \rangle_{\pm k}$ is approximately monochromatic.*

As has been shown by Gowers the above result yields the c_0 theorem. For the sake of completeness we include the proof here. Let F be a real-valued Lipschitz (or more generally a uniformly continuous) function on the unit sphere of c_0 . Fix $\varepsilon > 0$. Then there exists a $\delta > 0$ such that $|F(x) - F(y)| \leq \varepsilon/5$, for all $x, y \in S_{c_0}$ with $\|x - y\| \leq \delta$. Choose k with $(1 + \delta)^{-(k-1)} < \delta$ and define Δ to be the set of all finitely supported vectors of S_{c_0} all of whose coordinates belong to $\{0\} \cup \{\pm(1 + \delta)^{-(k-i)} : i = 1, \dots, k\}$. Then Δ is in one-to-one correspondence to $X_{\pm k}$ via the map $G : X_{\pm k} \rightarrow \Delta$, defined by $G(f) = f'$ where $f'(n) = (1 + \delta)^{-(k-i)}$ if $f(n) = i > 0$, $f'(n) = -(1 + \delta)^{-(k+i)}$ if $f(n) = i < 0$ and $f'(n) = 0$ if $f(n) = 0$. Since F is bounded, for the given $\varepsilon > 0$, there exist I_0, \dots, I_p disjoint intervals of length $\varepsilon/5$ that cover the range of F . Define a coloring $\{C_r : r = 1, \dots, p\}$ of $X_{\pm k}$ by putting f in C_r iff $F(f') \in I_r$.

By Theorem 2, there exists an infinite block sequence $\vec{f} = (f_n)_n$ in $X_{\pm k}$ and an r_0 such that $\langle \vec{f} \rangle_{\pm k} \subseteq \widehat{C}_{r_0}$. By the choice of k , for f, g neighbours of $X_{\pm k}$, we have that $\|f' - g'\| < \delta$, and this gives that the oscillation of F on $G(\langle \vec{f} \rangle_{\pm k})$ is at most $3\varepsilon/5$. We set $\vec{f}' = (f'_n)_n$, and let Y denote the linear span of \vec{f}' in c_0 . Then

$G(\langle \vec{f} \rangle_{\pm k})$ is easily seen to be δ -dense in S_Y , and so the oscillation of F on S_Y is at most ε .

2. A LEMMA ON COMPACT SEMIGROUPS

A *compact semigroup* is a semigroup $(S, +)$ with a topology with respect to which S is a compact Hausdorff space and the map $t \rightarrow t + s, t \in S$, is continuous for each $s \in S$. An element $\alpha \in S$ is called *idempotent* if $\alpha + \alpha = \alpha$. The following lemma due to R. Ellis is fundamental in the theory of compact semigroups.

Lemma 3. *If $(S, +)$ is a compact semigroup, then S contains an idempotent.*

Proof. Let $\mathcal{C} = \{S' \subseteq S : S' \text{ is a compact subsemigroup of } S\}$ ordered by inclusion. By Zorn's lemma there exists a minimal element M of \mathcal{C} . Pick an $\alpha \in M$ and observe that $M + \alpha$ is a compact subsemigroup of M and so $M + \alpha = M$. Hence the set $M' = \{\alpha' \in M : \alpha' + \alpha = \alpha\}$ is non-empty. Notice that M' is a compact subsemigroup of M , and so again by the minimality of M , we get that $M' = M$. Therefore $\alpha \in M'$ and $\alpha + \alpha = \alpha$. \square

A *two-sided ideal* of a semigroup $(S, +)$, is a subset I of S such that $S + I \subseteq I$ and $I + S \subseteq I$. On the set of all idempotents of a semigroup the following partial order is defined:

$$\alpha \leq \beta \iff \alpha + \beta = \beta + \alpha = \alpha.$$

Under the above notation a sequence $\alpha_0, \alpha_1, \alpha_2, \dots$ of idempotents will be called *increasing* if $\alpha_n \leq \alpha_{n+1}$ for all n . For $k = 0, 1, \dots$ and a map $T : S \rightarrow S$ we say that T is *k-stabilized* if $T^k = T^{k+1}$ or equivalently $T^k = T^{k+m}$, for all $m \geq 1$ (where by convention T^0 is the identity on S). Similarly a sequence $(\alpha_n)_n$ in S is called *k-stabilized* if $\alpha_k = \alpha_{k+m}$ for all $m \geq 1$. Notice that a map $T : S \rightarrow S$ is 0-stabilized if and only if T is the identity and that a sequence $(\alpha_n)_n$ is 0-stabilized if and only if it is constant.

Although the basic arguments below are essentially due to W. T. Gowers, the next lemma is inspired by a presentation of Gowers' theorem due to S. Todorćević [T].

Lemma 4. *Let $(S, +)$ be a compact semigroup, I a closed two-sided ideal of S , $k = 0, 1, \dots$, and $T : S \rightarrow S$ a continuous k -stabilized endomorphism. Then there exists an $\alpha \in I$ such that $\alpha, T\alpha, T^2\alpha, \dots$ is an increasing (k -stabilized) sequence of idempotents of $(S, +)$.*

Proof. We proceed by induction on k . If $k = 0$, then T is the identity on S and since I is a compact subsemigroup of S the result follows trivially by the previous lemma.

Suppose that the result is true for some $k \geq 0$ and let T be a continuous $(k + 1)$ -stabilized endomorphism of S . Let $S' = T(S), I' = T(I)$ and notice that $(S', +)$ is a compact semigroup, I' is a two-sided closed ideal of S' and the restriction of T to $(S', +)$ is a continuous k -stabilized endomorphism of S' . By our inductive assumption there exists an idempotent $\alpha' \in I'$ such that the sequence $(T^n \alpha')_n$ is an increasing k -stabilized sequence of idempotents of $(S, +)$. We set

$$G = \{\beta \in I : T\beta = \alpha'\} + \alpha'$$

and we claim that $(G, +)$ is a semigroup. Indeed, let $\beta_1 + \alpha', \beta_2 + \alpha'$ be in G . We have that $(\beta_1 + \alpha') + (\beta_2 + \alpha') = (\beta_1 + \alpha' + \beta_2) + \alpha'$ and since $(T^n \alpha')_n$ is increasing,

$\alpha' \leq T\alpha'$. Hence, $T(\beta_1 + \alpha' + \beta_2) = T\beta_1 + T\alpha' + T\beta_2 = \alpha' + T\alpha' + \alpha' = \alpha'$, which gives that $(G, +)$ is a semigroup.

By the continuity properties of T and $+$, the semigroup $(G, +)$ is also compact. Hence by Lemma 3, the compact semigroup $(G, +)$ contains an idempotent $\gamma + \alpha'$ with $\gamma \in I$ and $T\gamma = \alpha'$. We put $\alpha = \alpha' + \gamma + \alpha'$, and it is easily checked that α is an idempotent in I , $\alpha \leq \alpha'$ and $T\alpha = \alpha'$. Therefore $T^{n+1}\alpha = T^n\alpha'$ for all n , and it is straightforward that $(T^n\alpha)_n$ is an increasing $(k+1)$ -stabilized sequence of idempotents of $(S, +)$. \square

3. THE PROOF OF W. T. GOWERS' c_0 THEOREM

We will apply Lemma 4 to certain semigroups and two-sided ideals of ultrafilters. Before this let us start by recalling certain basic facts concerning ultrafilters. An *ultrafilter* α on a set X is a maximal filter on X , that is, a family of subsets of X that satisfies the following properties: (i) $\emptyset \notin \alpha$, (ii) if $A \in \alpha$ and $A \subseteq B \subseteq X$, then $B \in \alpha$, (iii) if $A, B \in \alpha$, then $A \cap B \in \alpha$, (iv) for every $A \subseteq X$, either A or $X \setminus A$ belongs to α . The set of all ultrafilters on X will be denoted by βX . An ultrafilter α is called *principal* if there exists $x \in X$ such that $\alpha = \{A \subseteq X : x \in A\}$. The principal ultrafilter determined by the element $x \in X$ is usually denoted by x^* . The set X is naturally identified with the set of principal ultrafilters via the map $X \ni x \rightarrow x^* \in \beta X$.

The set βX becomes a compact Hausdorff topological space, with basis the family of sets $A^* = \{\alpha \in \beta X : A \in \alpha\}$ where A ranges over all non-empty subsets of X . With this topology X is a dense open subset of βX . Of particular importance is the following property of βX . Whenever $\sigma : X \rightarrow X'$ is a map, then σ has a continuous extension $\beta\sigma : \beta X \rightarrow \beta X'$ defined by

$$\beta\sigma(\alpha) = \{A' \subseteq X' : \sigma^{-1}(A') \in \alpha\}.$$

It is easy to see that if $\sigma : X \rightarrow X'$ and $\tau : X' \rightarrow X''$, then $\beta(\tau \circ \sigma) = \beta\tau \circ \beta\sigma$. In particular for a map $\sigma : X \rightarrow X$ we have that $\beta\sigma^j = (\beta\sigma)^j$ for every j . In the sequel we shall denote the extension $\beta\sigma$ also by σ and we will write $\sigma\alpha$ instead of $\beta\sigma(\alpha)$.

Besides the above set-theoretic description of βX there is another one which will be very useful for our purposes. To every ultrafilter α on X , we associate the quantifier (αx) reading as “for α -almost $x \in X$ ”, in the following sense. If $P(x)$ is a statement concerning the elements $x \in X$, we write

$$(\alpha x)P(x) \iff \{x \in X : P(x)\} \in \alpha,$$

that is, “for α -almost $x \in X$, $P(x)$ is true” if and only if the set of all x that satisfy $P(x)$ belongs to the ultrafilter α . Substituting $P(x)$ with $(x \in A)$, we get that

$$(\alpha x)(x \in A) \iff A \in \alpha.$$

Standard properties of ultrafilters yield that the quantifier (αx) commutes with conjunction (\wedge) and negation (\neg) , namely

$$(\alpha x)P(x) \wedge (\alpha x)Q(x) \iff (\alpha x)(P(x) \wedge Q(x))$$

and

$$\neg(\alpha x)P(x) \iff (\alpha x)(\neg P(x)).$$

For example, if $\sigma : X \rightarrow X'$, then for every $\alpha \in \beta X$ the associated to $\sigma\alpha$ quantifier is defined by

$$(\sigma\alpha x')P(x') \iff (\alpha x)P(\sigma(x))$$

or equivalently,

$$A' \in \sigma\alpha \iff (\sigma\alpha x')(x' \in A') \iff (\alpha x)(\sigma(x) \in A').$$

Another important fact concerning βX is that it inherits the algebraic properties of X . Specifically if X is a semigroup, then it can be shown that βX is a compact semigroup. The classical example is the compact semigroup $(\beta\mathbb{N}, +)$. In our case the set $X_{[k]} = \bigcup_{i=0}^k X_i$, $k \geq 1$, with the usual operation $+$, although it is not a semigroup it is a partial semigroup, where $f + g$ is defined if either f or g are equal to $\theta \equiv 0$, or otherwise if $\max \text{supp } f < \min \text{supp } g$. However this is enough to guarantee that $\beta X_{[k]}$ contains a compact semigroup. To be more precise let us set

$$X_{[k]}^n = \{f \in X_{[k]} : f(0) = \dots = f(n) = 0\}$$

for every $n = 0, 1, \dots$. Then the family $\{X_{[k]}^n\}_n$ has the finite intersection property and therefore we can define

$$\delta X_{[k]} = \{\alpha \in \beta X_{[k]} : X_{[k]}^n \in \alpha, \text{ for every } n\}.$$

Observe that $\delta X_{[k]} = \bigcap_n (X_{[k]}^n)^*$, and so $\delta X_{[k]}$ is a non-empty compact subspace of $\beta X_{[k]}$. Also since $\theta \in X_{[k]}^n$ for all n , the principal ultrafilter $\theta^* = \{A \subseteq X_{[k]} : A \text{ contains the function } \theta \equiv 0\}$ belongs to $\delta X_{[k]}$ and moreover it is the only principal ultrafilter in $\delta X_{[k]}$.

On $\delta X_{[k]}$ we define the addition $+$ by

$$\alpha + \beta = \{A \subseteq X_{[k]} : (\alpha f)(\beta g)(f + g \text{ is defined and } f + g \in A)\}$$

or equivalently,

$$A \in \alpha + \beta \iff \{f \in X_{[k]} : A_f \in \beta\} \in \alpha$$

where $A_f = \{g \in X_{[k]} : f + g \text{ is defined and } f + g \in A\}$. The following three propositions are essentially from [BBH] where they are stated and proved in greater generality.

Proposition 5. *The set $\delta X_{[k]}$ with the operation $+$ is a compact semigroup and θ^* is its unit element.*

Proof. We leave to the reader the verification that $\alpha + \beta$ is indeed an ultrafilter, the operation $+$ is associative and the map $\alpha \rightarrow \alpha + \beta$, $\alpha \in \delta X_k$ is continuous for any $\beta \in \delta X_k$ that is obtained in a similar manner as in the case of $\beta\mathbb{N}$.

We show the rest of the conclusions. We begin by showing that $\alpha + \beta \in \delta X_k$ for $\alpha, \beta \in \delta X_k$. By the definition of $\delta X_{[k]}$, it is enough to check that $X_{[k]}^n \in \alpha + \beta$ for all n . Fix n and set for simplicity $A = X_{[k]}^n$. Then for every $f \in A$, we have that $A_f \in \beta$. Indeed, if $f = \theta$, then $A_f = A$; otherwise $A_f = X_{[k]}^{n_0}$ where $n_0 = \max \text{supp } f$. Therefore $A \subseteq \{f \in X_{[k]} : A_f \in \beta\}$, and since $A \in \alpha$, we obtain that $\{f \in X_{[k]} : A_f \in \beta\} \in \alpha$, which by definition means that $A \in \alpha + \beta$.

To show that $\alpha = \alpha + \theta^*$, observe that $A \in \alpha \implies \forall f \in A, \theta \in A_f \iff \forall f \in A, A_f \in \theta^* \iff A \subseteq \{f \in X_{[k]} : A_f \in \theta^*\} \implies \{f \in X_{[k]} : A_f \in \theta^*\} \in \alpha \iff A \in \alpha + \theta^*$. Hence $\alpha \subseteq \alpha + \theta^*$, and since both are ultrafilters $\alpha = \alpha + \theta^*$. The equality $\theta^* + \alpha = \alpha$ is similarly shown. \square

We now describe the properties of the extension of $T : X_{[k]} \rightarrow X_{[k]}$ to $\beta X_{[k]}$. Observe that if $f + g$ is defined, then $Tf + Tg$ is also defined (the converse fails) and $T(f + g) = Tf + Tg$, that is, T is an endomorphism of the partial semigroup $(X_{[k]}, +)$.

Proposition 6. *The extension of $T : X_{[k]} \rightarrow X_{[k]}$ to $T : \beta X_{[k]} \rightarrow \beta X_{[k]}$ maps $\delta X_{[k]}$ into $\delta X_{[k]}$, and its restriction to $(\delta X_{[k]}, +)$ is a continuous k -stabilized endomorphism.*

Proof. First let us verify that $T(\alpha + \beta) = T\alpha + T\beta$ for $\alpha, \beta \in \delta X_{[k]}$. Indeed,

$$\begin{aligned} A \in T(\alpha + \beta) &\iff T^{-1}(A) \in \alpha + \beta \\ &\iff (\alpha f)(\beta g)(f + g \text{ is defined and } f + g \in T^{-1}(A)) \\ &\iff (\alpha f)(\beta g)(f + g \text{ is defined and } T(f + g) \in A) \\ &\iff (\alpha f)(\beta g)(f + g \text{ is defined and } Tf + Tg \in A) \\ &\implies (\alpha f)(\beta g)(Tf + Tg \text{ is defined and } Tf + Tg \in A) \\ &\iff (T\alpha f')(T\beta g')(f' + g' \text{ is defined and } f' + g' \in A) \\ &\iff A \in T\alpha + T\beta. \end{aligned}$$

Hence $T(\alpha + \beta) \subseteq T\alpha + T\beta$, and so $T(\alpha + \beta) = T\alpha + T\beta$. To see that T maps $\delta X_{[k]}$ into itself, observe that $X_{[k]}^n \subseteq T^{-1}(X_{[k]}^n)$ for all n and so $T\alpha \in \delta X_{[k]}$ for every $\alpha \in \delta X_{[k]}$. To show that T is k -stabilized notice that for every $f \in X_{[k]}$, $T^{k+1}f = T^k f = \theta$, which easily gives $T^{k+1}\alpha = T^k\alpha = \theta^*$ for all $\alpha \in \beta X_{[k]}$. Finally the continuity of T follows from the continuity of $T : \beta X_{[k]} \rightarrow \beta X_{[k]}$. \square

We set

$$X_k^n = \{f \in X_k : f(0) = \dots = f(n) = 0\}$$

for every $n = 0, 1, \dots$ and we define

$$\delta X_k = \{\alpha \in \beta X_k : X_k^n \in \alpha \text{ for every } n\}.$$

Again by the finite intersection property of the family $\{X_k^n\}_n$, the set δX_k is a compact non-empty subset of βX_k . As in the case of $\delta X_{[k]}$, it can be shown that $(\delta X_k, +)$ is a compact semigroup with the operation $+$ having an analogous meaning. Notice that for $f, g \in X_{[k]}$, $f + g \in X_k$ whenever $f + g$ is defined and f or $g \in X_k$ i.e. X_k is a two-sided ideal of the partial semigroup $(X_{[k]}, +)$. This property is carried over to δX_k in the following sense.

Proposition 7. *The extension of the inclusion map $I : X_k \hookrightarrow X_{[k]}$ to $I : \beta X_k \rightarrow \beta X_{[k]}$ maps δX_k into $\delta X_{[k]}$, and its restriction to $(\delta X_k, +)$ is a continuous isomorphism onto a (closed) two-sided ideal of $(\delta X_{[k]}, +)$.*

Proof. As in Proposition 6, it can be checked that $I : \beta X_k \rightarrow \beta X_{[k]}$ maps $(\delta X_k, +)$ isomorphically into $(\delta X_{[k]}, +)$. Observe that for an ultrafilter $\beta \in \delta X_{[k]}$, we have that

$$\beta \in I(\delta X_k) \iff X_k \in \beta.$$

Since X_k is a two-sided ideal of $X_{[k]}$, it is easy to see that for $\alpha \in \delta X_{[k]}$ and $\beta \in I(\delta X_k)$, $X_k \in \alpha + \beta$, and so by the above remark $\alpha + \beta \in I(\delta X_k)$. Similarly it is shown that $\beta + \alpha \in I(\delta X_k)$, and so $I(\delta X_k)$ is a two-sided ideal of $\delta X_{[k]}$. \square

In the sequel we will identify δX_k with the closed two-sided ideal $I(\delta X_k)$ of $(\delta X_{[k]}, +)$.

Finally, the semigroups $(\delta X_{[\pm k]}, +)$ and $(\delta X_{\pm k}, +)$ are defined analogously, and it is easy to see that they satisfy what corresponds to the above propositions. Since for every $f \in X_{[k]}$ and every $\beta \in \delta X_{[k]}$, we have that $(\beta g)(f + g)$ is defined, in what follows we simply write $(\alpha f)(\beta g)(f + g \in A)$ to denote that $A \in \alpha + \beta$ for $\alpha, \beta \in \delta X_{[k]}$ (and similarly for $\delta X_{[\pm k]}$).

We are now ready to prove W. T. Gowers' theorem.

Proof of Theorem 1. We set

$$\mathcal{I}(\delta X_k, T) = \{\alpha \in \delta X_k : \alpha + T^j \alpha = T^j \alpha + \alpha = \alpha \text{ for every } j = 0, \dots, k\}.$$

By the above propositions and Lemma 4, for $S = \delta X_{[k]}$ and $I = \delta X_k$, we obtain that

$$\mathcal{I}(\delta X_k, T) \neq \emptyset,$$

and so we can choose an $\alpha \in \mathcal{I}(\delta X_k, T)$.

Lemma 8. *For every $A \in \alpha$ there exists a block sequence \vec{f} in X_k such that $\langle \vec{f} \rangle_k \subseteq A$.*

Proof. Since $\alpha \in \mathcal{I}(\delta X_k, T)$ we have that for every $A \in \alpha$,

$$(\alpha f)(\alpha g)(\{f, g, f + T^j g, T^j f + g : j = 0, 1, \dots, k - 1\} \subseteq A)$$

or equivalently,

$$(1) \quad (\alpha f)(\alpha g)(\langle f, g \rangle_k \subseteq A).$$

By induction pick $A = A_0 \supseteq A_1 \supseteq \dots \supseteq A_n \supseteq \dots$ and $f_0 < f_1 < \dots < f_n < \dots$ such that $\forall n, A_n \in \alpha, f_n \in A_n$ and $A_{n+1} = \{g \in X_k : \langle f_n, g \rangle_k \subseteq A_n\}$.

To start set $A_0 = A$ and by (1) choose an $f_0 \in X_k$ so that $(\alpha g)(\langle f_0, g \rangle_k \subseteq A_0)$ and put $A_1 = \{g \in X_k : \langle f_0, g \rangle_k \subseteq A_0\}$. Then $f_0 \in A_0, A_1 \in \alpha$ and $A_0 \supseteq A_1$. Suppose that A_0, \dots, A_n and f_0, \dots, f_{n-1} have been constructed so as to satisfy the required properties. Since A_n belongs to α , (1) holds for A_n in place of A , and so we can pick an $f_n \in X_k$ so that $f_{n-1} < f_n$ and $(\alpha g)(\langle f_n, g \rangle_k \subseteq A_n)$. We set $A_{n+1} = \{g \in X_k : \langle f_n, g \rangle_k \subseteq A_n\}$ and clearly we have that $A_{n+1} \in \alpha, f_n \in A_n, A_{n+1} \subseteq A_n$, and so the induction can be carried out. By downwards induction it is easily checked that $\langle f_i, \dots, f_n \rangle_k \subseteq A_i$, for all $i \leq n$ and so $\langle f_0, f_1, \dots, f_n \rangle_k \subseteq A_0 = A$, for all n . \square

Since one of the pieces of every finite coloring of X_k belongs to α the proof of Theorem 1 is complete. \square

Proof of Theorem 2. The main idea here is to apply Lemma 4 for $S = \delta X_{[\pm k]}$, $I = \delta X_{\pm k}$ and $-T$ instead of T defined by $-Tf = -(Tf) = T(-f)$ for all $f \in X_{[\pm k]}$.

We set

$$\mathcal{I}(\delta X_{\pm k}, -T) = \{\alpha \in \delta X_{\pm k} : \alpha + (-T)^j \alpha = (-T)^j \alpha + \alpha = \alpha \text{ for every } j = 0, \dots, k\}.$$

It is easily shown that $-T$ satisfies the obvious analogue of Proposition 6, and so Lemma 4 gives that for every $k \geq 1$,

$$\mathcal{I}(\delta X_{\pm k}, -T) \neq \emptyset.$$

For a block sequence $\vec{f} = (f_n)_n$ in $X_{\pm k}$ we define

$$\langle \vec{f} \rangle_{(-T)} = \{(-T)^{\epsilon_0} f_{n_0} + \dots + (-T)^{\epsilon_m} f_{n_m} : 0 \leq \epsilon_i \leq k-1, n_0 < \dots < n_m \\ \text{and } \exists i \text{ with } \epsilon_i = 0\}.$$

Lemma 9. For a symmetric set $A \subseteq X_{\pm k}$ (i.e. $A = -A$) and a block sequence \vec{f} in $X_{\pm k}$, if $\langle \vec{f} \rangle_{(-T)} \subseteq A$, then $\langle \vec{f} \rangle_{\pm k} \subseteq \widehat{A}$.

Proof. Let $f \in \langle \vec{f} \rangle_{\pm k}$. Then there exist $m \in \mathbb{N}$, $n_0 < \dots < n_m$, $\epsilon_0, \dots, \epsilon_m \in \{0, \dots, k-1\}$, with $\epsilon_i = 0$ for some i and $\delta_0, \dots, \delta_m \in \{-1, +1\}$, such that

$$f = \delta_0 T^{\epsilon_0} f_{n_0} + \dots + \delta_m T^{\epsilon_m} f_{n_m}.$$

We distinguish two cases:

Case 1. For some $i \in \{0, \dots, m\}$, $\epsilon_i = 0$ and $\delta_i = +1$. Then for every $i \in \{0, \dots, m\}$ we set $g_i = \delta_i T^{\epsilon_i} f_{n_i}$ and

(a) If $\delta_i = -1$ and ϵ_i is odd, or $\delta_i = +1$ and ϵ_i is even, we set $g'_i = g_i$. Observe that in these subcases $g'_i = g_i = (-T)^{\epsilon_i} f_{n_i}$.

(b) If $\delta_i = -1$ and ϵ_i is even, or $\delta_i = +1$ and ϵ_i is odd, we set $g'_i = Tg_i$. Then it is easy to see that $g'_i = (-T)^{\epsilon_i+1} f_{n_i}$.

In both subcases, by the choice of g'_i , we have that g'_i, g_i are neighbours for every $i = 0, \dots, m$. We set $f' = g'_0 + \dots + g'_m$. Then $f' \in \langle \vec{f} \rangle_{(-T)} \subseteq A$ and $\|f - f'\|_\infty \leq 1$. Hence $f \in \widehat{A}$.

Case 2. For every $i \in \{0, \dots, m\}$ if $\epsilon_i = 0$, then $\delta_i = -1$. Let $g = -f$. Then the representation of g satisfies the condition of Case 1, and so there exists $g' \in \langle \vec{f} \rangle_{(-T)}$ such that $\|g - g'\|_\infty \leq 1$. Since $\langle \vec{f} \rangle_{(-T)} \subseteq A$ and A is symmetric we have that $h = -g' \in A$ and $\|f - h\|_\infty = \|g - g'\|_\infty \leq 1$.

Therefore for every $f \in \langle \vec{f} \rangle_{\pm k}$, we have that $f \in \widehat{A}$, that is, $\langle \vec{f} \rangle_{\pm k} \subseteq \widehat{A}$. \square

In the following two lemmas, we assume that $\alpha \in \mathcal{I}(\delta X_{\pm k}, -T)$.

Lemma 10. For every $A \in \alpha$ there exists a block sequence \vec{f} in $X_{\pm k}$ such that $\langle \vec{f} \rangle_{(-T)} \subseteq A$.

Proof. This is identical to that of Lemma 8. \square

Lemma 11. For every $A \in \alpha$, $-\widehat{A} \in \alpha$.

Proof. Since $\alpha \leq -T\alpha$, we have that

$$\begin{aligned} A \in \alpha = -T\alpha + \alpha &\iff (\alpha f)(\alpha g)(-Tf + g \in A) \\ &\implies (\alpha f)(\alpha g)(-f + Tg \in \widehat{A}) \text{ (since } f < g) \\ &\iff (\alpha f)(\alpha g)(f - Tg \in -\widehat{A}) \\ &\iff -\widehat{A} \in \alpha - T\alpha = \alpha. \end{aligned}$$

\square

For $A \subseteq X_{\pm k}$, by $\widehat{\widehat{A}}$ we denote the set of all neighbours of the elements of \widehat{A} . It is straightforward that $\widehat{\widehat{A}} = \{f \in X_{[\pm k]} : \exists g \in A \text{ such that } \|f - g\|_\infty \leq 2\}$. The following proposition is a weak version of Theorem 2.

Proposition 12. For every $\alpha \in \mathcal{I}(\delta X_{\pm k}, -T)$ and every $A \in \alpha$, there exists a block sequence \vec{f} in $X_{\pm k}$ such that $\langle \vec{f} \rangle_{\pm k} \subseteq \widehat{\widehat{A}}$.

Proof. Applying Lemma 11 we get that the set $A' = \widehat{A} \cap -\widehat{A}$ belongs to α . By Lemma 10 there exists a block sequence \vec{f} in $X_{\pm k}$ such that $\langle \vec{f} \rangle_{(-T)} \subseteq A'$, and since A' is symmetric Lemma 9 gives that $\langle \vec{f} \rangle_{\pm k} \subseteq \widehat{A}'$. Since $A' \subseteq \widehat{A}$ we have that $\langle \vec{f} \rangle_{\pm k} \subseteq \widehat{A}$. \square

Remark 1. Notice that the above weak version of Theorem 2 suffices to prove that every real-valued Lipschitz function on S_{c_0} is oscillation stable.

To complete the proof we define the map $\Phi : X_{\pm 2k} \rightarrow X_{\pm k}$, where for every $f \in X_{\pm 2k}$ and $n \in \mathbb{N}$, $\Phi f(n)$ is defined as follows:

$$\Phi f(n) = \begin{cases} \frac{f(n)}{2} & \text{if } f(n) \text{ is even,} \\ \frac{f(n)-1}{2} & \text{if } f(n) > 0 \text{ and } f(n) \text{ is odd,} \\ \frac{f(n)+1}{2} & \text{if } f(n) < 0 \text{ and } f(n) \text{ is odd.} \end{cases}$$

The following properties of Φ are straightforward:

- (i) It is onto $X_{\pm k}$.
- (ii) For every $f_1 < f_2$ in $X_{\pm 2k}$, $\Phi(f_1 + f_2) = \Phi f_1 + \Phi f_2$.
- (iii) For every $f \in X_{\pm 2k}$, $\Phi(-f) = -(\Phi f)$.
- (iv) For every $f_1 < f_2$ in $X_{\pm 2k}$ and every $\epsilon_1, \epsilon_2 \in \{0, \dots, k\}$ with $\epsilon_i = 0$ for at least one $i \in \{1, 2\}$, $T^{\epsilon_1}(\Phi f_1) + T^{\epsilon_2}(\Phi f_2) = \Phi(T^{2\epsilon_1} f_1 + T^{2\epsilon_2} f_2)$.
- (v) For every $f_1, f_2 \in X_{\pm 2k}$, if $\|f_1 - f_2\|_\infty \leq 2$, then $\|\Phi f_1 - \Phi f_2\|_\infty \leq 1$.

We recall that Φ is extended to $\Phi : \beta X_{\pm 2k} \rightarrow \beta X_{\pm k}$, where for every $\alpha \in \beta X_{\pm 2k}$, $\Phi\alpha = \{A' \subseteq X_{\pm k} : \Phi^{-1}(A') \in \alpha\}$.

Lemma 13. For every $\alpha \in \mathcal{I}(\delta X_{\pm 2k}, -T)$ and every $A' \in \Phi\alpha$ there exists a block sequence \vec{f}' in $X_{\pm k}$ such that $\langle \vec{f}' \rangle_{\pm k} \subseteq \widehat{A}'$.

Proof. By the definition of $\Phi\alpha$ we have that $A = \Phi^{-1}(A') \in \alpha$. By Proposition 12 there exists a block sequence \vec{f} in $X_{\pm 2k}$ such that $\langle \vec{f} \rangle_{\pm 2k} \subseteq \widehat{A}$. We set $f'_n = \Phi f_n$ and $\vec{f}' = (f'_n)_n$. It is clear that \vec{f}' is a block sequence in $X_{\pm k}$. We will show that $\langle \vec{f}' \rangle_{\pm k} \subseteq \widehat{A}'$.

So let $f' \in \langle \vec{f}' \rangle_{\pm k}$. Then there exist $m \in \mathbb{N}$, $n_0 < \dots < n_m$, $\epsilon_0, \dots, \epsilon_m \in \{0, \dots, k-1\}$, with $\epsilon_i = 0$ for some i and $\delta_0, \dots, \delta_m \in \{-1, +1\}$, such that

$$\begin{aligned} f' &= \delta_0 T^{\epsilon_0} f'_{n_0} + \dots + \delta_m T^{\epsilon_m} f'_{n_m} \\ &= \delta_0 T^{\epsilon_0} \Phi f_{n_0} + \dots + \delta_m T^{\epsilon_m} \Phi f_{n_m} = \Phi(\delta_0 T^{2\epsilon_0} f_{n_0} + \dots + \delta_m T^{2\epsilon_m} f_{n_m}) \end{aligned}$$

by the properties (ii)-(iv) of Φ . Setting $f = \delta_0 T^{2\epsilon_0} f_{n_0} + \dots + \delta_m T^{2\epsilon_m} f_{n_m}$ we have that $f' = \Phi f$ and $f \in \langle \vec{f} \rangle_{\pm 2k}$. Since $\langle \vec{f} \rangle_{\pm 2k} \subseteq \widehat{A}$ where $A = \Phi^{-1}(A')$, there exists $g \in \Phi^{-1}(A')$, such that $\|f - g\|_\infty \leq 2$. We set $g' = \Phi g$ and we have that $g' \in A'$ and by the property (v) of Φ , $\|f' - g'\|_\infty = \|\Phi f - \Phi g\|_\infty \leq 1$. Hence for every $f' \in \langle \vec{f}' \rangle_{\pm k}$, we have that $f' \in \widehat{A}'$, that is, $\langle \vec{f}' \rangle_{\pm k} \subseteq \widehat{A}'$. \square

Since for every $\alpha \in \mathcal{I}(\delta X_{\pm 2k}, -T)$ and every finite coloring of $X_{\pm k}$ one of the pieces of the partition belongs to $\Phi\alpha$, the proof of Theorem 2 is complete. \square

4. NOTES

1. Theorems 1 and 2 have interesting corollaries that are actually equivalent to them and they illustrate the property $\alpha \leq T\alpha \leq \dots \leq T^k\alpha$ of the sequence $(T^n\alpha)_n$ obtained in Lemma 4. To state them we need to extend slightly the definition of the span of a block sequence as follows. For an infinite block sequence $\vec{f} = (f_n)_n$ in $X_{[k]}$ define its span in $X_{[k]}$ to be the set

$$\langle \vec{f} \rangle_{[k]} = \left\{ T^{\epsilon_0} f_{n_0} + T^{\epsilon_1} f_{n_1} + \dots + T^{\epsilon_m} f_{n_m} : m \in \mathbb{N}, \epsilon_i = 0, \dots, k-1 \text{ and } n_0 < n_1 < \dots < n_m \right\},$$

and for all $i = 1, \dots, k$ define the *relative span* in X_i by $\langle \vec{f} \rangle_i = \langle \vec{f} \rangle_{[k]} \cap X_i$. We will also use the notation $T^j \vec{f} = (T^j f_n)_n$ for a block sequence $\vec{f} = (f_n)_n$ of X_k and $j = 1, \dots, k-1$. Then $\langle \vec{f} \rangle_i = \langle T^{k-i} \vec{f} \rangle_i = T^{k-i}(\langle \vec{f} \rangle_k)$ for all $i = 1, \dots, k$.

Given two block sequences \vec{f}, \vec{g} in X_k , we say that $\vec{f} = (f_n)$ is a block subsequence of \vec{g} , denoted by $\vec{f} \leq \vec{g}$, if $f_n \in \langle \vec{g} \rangle_k$ for all n . Under the above notation we have the following corollary of Theorem 1.

Corollary 14. *Let $k \geq 1$ and suppose that for each $i = 1, \dots, k$, the set X_i is finitely colored. Then for every block sequence $\vec{g} = (g_n)_n$ in X_k there exists an infinite block subsequence $\vec{f} \leq \vec{g}$ such that for each $i = 1, \dots, k$, the span $\langle \vec{f} \rangle_i$ is monochromatic.*

Proof. Let us denote by $\vec{e} = (e_n)_n$ the block sequence in X_k consisting of the functions e_n where $e_n(m) = k$ if $n = m$ and $e_n(m) = 0$ otherwise. Notice that $X_k = \langle \vec{e} \rangle_k$ and every $f \in X_k$ has a unique representation $f = T^{\epsilon_0} e_{n_0} + T^{\epsilon_1} e_{n_1} + \dots + T^{\epsilon_m} e_{n_m}$, where $\epsilon_0, \dots, \epsilon_m \in \{0, \dots, k-1\}$. We define the map $P : X_k \rightarrow \langle \vec{g} \rangle_k$ by $P(T^{\epsilon_0} e_{n_0} + T^{\epsilon_1} e_{n_1} + \dots + T^{\epsilon_m} e_{n_m}) = T^{\epsilon_0} g_{n_0} + T^{\epsilon_1} g_{n_1} + \dots + T^{\epsilon_m} g_{n_m}$.

It is easily verified that P is bijective and carries block sequences in X_k to block subsequences of \vec{g} . Since the coloring of $\langle \vec{g} \rangle_k$ induces via P^{-1} a corresponding coloring of X_k , Theorem 1 and the above property of P yield that there exists a block sequence \vec{f}_1 of X_k with $\vec{f}_1 \leq \vec{g}$ such that $\langle \vec{f}_1 \rangle_k$ is monochromatic. Using the same arguments for the case $k-1$ instead of k and the block sequence $T\vec{f}_1$ in place of \vec{g} , we obtain a block sequence $\vec{f}_2 \leq T\vec{f}_1$ such that $\langle \vec{f}_2 \rangle_{k-1}$ is monochromatic. It is now easily shown that there exists a block sequence \vec{f}_2 of X_k such that $\vec{f}_2 \leq \vec{f}_1$ and $T\vec{f}_2 = \vec{f}_2$. Then $\langle \vec{f}_2 \rangle_{k-1} = \langle \vec{f}_2 \rangle_{k-1}$, $\langle \vec{f}_2 \rangle_k \subseteq \langle \vec{f}_1 \rangle_k$ and so the relative spans of \vec{f}_2 in X_k and X_{k-1} are monochromatic. Repeating the same procedure, after k steps we obtain an $\vec{f} = \vec{f}_k \leq \vec{g}$ with monochromatic relative span in each X_i , $i = 1, \dots, k$. \square

Stating the analogous definitions concerning the general case $X_{[\pm k]}$ we get the next corollary of Theorem 2.

Corollary 15. *Let $k \geq 1$ and suppose that for each $i = 1, \dots, k$, the set $X_{\pm i}$ is finitely colored. Then for every block sequence $\vec{g} = (g_n)_n$ in $X_{\pm k}$ there exists an infinite block subsequence $\vec{f} \leq \vec{g}$ such that for each $i = 1, \dots, k$, the span $\langle \vec{f} \rangle_{\pm i}$ is approximately monochromatic.*

2. The sequence of idempotents obtained by Lemma 4 can be chosen so that all its terms $\alpha, T\alpha, \dots$ will, in addition, be minimal. This can be done by using in the inductive step of the proof the fact that for every idempotent α and every

two-sided ideal I there exists a minimal idempotent α' with $\alpha' \leq \alpha$ and $\alpha' \in I$ (see for example [FK] for a proof of this statement). However we do not see any further property resulting from the minimality of the idempotent element.

3. It is also possible to derive some Hales-Jewett versions of W. T. Gowers' combinatorial theorems in the context of variable words. Given a non-empty set Γ , a *word* over Γ is a map from a proper initial segment of \mathbb{N} into Γ (i.e. a finite sequence in Γ). The set of all words over Γ , denoted by $W(\Gamma)$, with the operation of concatenation $*$ is a semigroup.

To state the first abstract version take for an alphabet a singleton set $A = \{u_0\}$ and for $k \geq 1$ let u_1, \dots, u_k be distinct variables. For $i = 0, \dots, k$ we set $W_{[i]} = W(\{u_0, \dots, u_i\})$ and $V_i = W_{[i]} \setminus W_{[i-1]}$, for all $i = 0, \dots, k$ (where by convention $W_{[-1]} = \emptyset$). Let T be the homomorphic extension to $W_{[k]}$ of the map $u_i \rightarrow u_{i-1}$ for $i = 1, \dots, k$ and $u_0 \rightarrow u_0$. For an infinite sequence $\vec{w} = (w_n)_n$ of words of V_k define its span in V_k to be the set

$$\langle \vec{w} \rangle_k = \{T^{\epsilon_0} w_{n_0} * T^{\epsilon_1} w_{n_1} * \dots * T^{\epsilon_m} w_{n_m} : \epsilon_i = 0, \dots, k, \quad n_0 < n_1 < \dots < n_m, \quad \text{and } \exists i \text{ with } \epsilon_i = 0\}.$$

Under the above definitions, we have the following analogue of Theorem 1, which is related to Carlson's results [C].

Theorem 16. *For $k \geq 0$ and every finite coloring of V_k there exists an infinite sequence \vec{w} in V_k such that the span $\langle \vec{w} \rangle_k$ is monochromatic.*

Identifying $n \in \{1, 2, \dots\}$ with the word $w = u_0 * \dots * u_0$ (n times), the case $k = 0$ of the above theorem is Hindman's theorem for monochromatic finite sums of a sequence of positive integers. The proof of Theorem 16 is similar to that of Theorem 1 and actually it is simpler, since Lemma 4 is applied directly to $S = \beta W_k$ and $I = \beta V_k$.

We have that Theorem 1 can be derived from the above theorem. To show this we define the map $\Psi : V_k \rightarrow X_k$ for $k \geq 1$ where for every $w \in V_k$, $\Psi w(n) = i$ if $w(n) = u_i$ and $\Psi w(n) = 0$ if $w(n)$ is undefined. The coloring of X_k induces a coloring of V_k by lifting via Ψ . Let $\vec{w} = (w_n)_n$ be the resulting (by Theorem 16) sequence in V_k , and define $f_0 = \Psi w_0$ and $f_n = \Psi(T^k(w_0 * \dots * w_{n-1}) * w_n)$, for all $n \geq 1$. It is easy to show that $\langle \vec{f} \rangle_k \subseteq \Psi(\langle \vec{w} \rangle_k)$ and so $\langle \vec{f} \rangle_k$ is monochromatic.

Other abstract versions can be derived using the concept of located words, which was introduced by V. Bergelson, A. Blass and N. Hindman in [BBH], in order to extend the results of T. Carlson in [C]. For a set Γ , a *located* word over Γ is a map from a nonempty finite subset of \mathbb{N} into Γ . Note that this notion is actually a generalization of the common notion of a word over Γ since its domain is not necessarily an initial segment of \mathbb{N} . The set of located words over Γ , denoted by $L(\Gamma)$ becomes a partial semigroup under the operation $+$, defined by $w + w' = w \cup w'$ whenever $\text{dom}(w) < \text{dom}(w')$. As above let $A = \{u_0\}$ and for $k \geq 1$ let u_1, \dots, u_k be distinct variables. Set $L_{[i]} = L(\{u_0, \dots, u_i\})$ and $L_i = L_{[i]} \setminus L_{[i-1]}$, for all $i = 0, \dots, k$ (where $L_{[-1]} = \emptyset$). An infinite sequence of located words $\vec{w} = (w_n)_n$ is said to be a block sequence if $\text{dom}(w_n) < \text{dom}(w_{n+1})$ for every n . The span of a block sequence $\langle \vec{w} \rangle_k$ is defined as in the case of ordinary words.

Theorem 17. *For $k \geq 0$ and every finite coloring of L_k there exists an infinite block sequence \vec{w} in L_k such that the span $\langle \vec{w} \rangle_k$ is monochromatic.*

One also could consider abstract versions of Theorem 2 by introducing “negative” variables u_{-1}, \dots, u_{-k} and defining the span of a sequence of words (or of a block sequence of located words) analogously.

ACKNOWLEDGEMENT

The author would like to thank Prof. S.A. Argyros and Dr. P. Dodos for their help in the preparation of the paper.

REFERENCES

- [BL] Y. Benyamini, J. Lindenstrauss, *Geometric Nonlinear Functional Analysis*, Vol. 1, American Mathematical Society Colloquium Publications 48, American Mathematical Society, Providence, RI, (2000). MR 1727673 (2001b:46001)
- [BBH] V. Bergelson, A. Blass, N. Hindman, *Partition theorems for spaces of variable words*, Proc. London Math. Soc. 3, 68, 449-476, (1994). MR 1262304 (95i:05107)
- [C] T. Carlson, *Some unified principles in Ramsey theory*, Discrete Math. 68, 117-169, (1988). MR 0926120 (89b:04006)
- [FK] H. Furstenberg and Y. Katznelson, *Idempotents in compact semigroups and Ramsey theory*, Israel Jour. Math. 68, 257-270, (1989). MR 1039473 (92d:05170)
- [G1] W. T. Gowers, *Lipshitz functions on classical spaces*, Europ. Jour. Combinatorics, 13, 141-151, (1992). MR 1164759 (93g:05142)
- [G2] W. T. Gowers, *Ramsey methods in Banach spaces*, Handbook of the Geometry of Banach Spaces, Vol. 2, pp. 1071–1097, W. B. Johnson and J. Lindenstrauss, eds., Elsevier, Amsterdam (2003). MR 1999191
- [T] S. Todorcevic, *Lecture notes in infinite-dimensional Ramsey theory*, (manuscript, Univ. of Toronto (1998)).

DEPARTMENT OF MATHEMATICS, NATIONAL TECHNICAL UNIVERSITY OF ATHENS, ATHENS 15780, GREECE

E-mail address: `bkanel@math.ntua.gr`