

LATTICE POLYGONS AND GREEN'S THEOREM

HAL SCHENCK

(Communicated by Michael Stillman)

ABSTRACT. Associated to an n -dimensional integral convex polytope P is a toric variety X and divisor D , such that the integral points of P represent $H^0(\mathcal{O}_X(D))$. We study the free resolution of the homogeneous coordinate ring $\bigoplus_{m \in \mathbb{Z}} H^0(mD)$ as a module over $Sym(H^0(\mathcal{O}_X(D)))$. It turns out that a simple application of Green's theorem yields good bounds for the linear syzygies of a projective toric surface. In particular, for a planar polytope $P = H^0(\mathcal{O}_X(D))$, D satisfies Green's condition N_p if ∂P contains at least $p + 3$ lattice points.

1. GREEN'S THEOREM AND HYPERPLANE SECTIONS

For a curve C of genus g , a divisor D of degree $d \geq 2g + 1$ is very ample, so gives an embedding of C into projective space. In fact, when $d \geq 2g + 1$, work of Castelnuovo, Mattuck and Mumford shows that the embedding is *projectively normal*, which means that $S = Sym(H^0(\mathcal{O}_X(D)))$ surjects onto $\bigoplus_{m \in \mathbb{Z}} H^0(mD) = R$. When $d \geq 2g + 2$, results of Fujita and St. Donat show that the homogeneous ideal of I_C is generated by quadrics. Let F_\bullet be a minimal free resolution of R over S . A very ample divisor is said to satisfy property N_p if $F_0 = S$ and $F_q \simeq \bigoplus S(-q - 1)$ for all $q \in \{1, \dots, p\}$. Thus, N_0 means projectively normal, N_1 means that the homogeneous ideal is generated by quadrics, N_2 means that the minimal syzygies on the quadrics are linear, and so on. In [7], Green used Koszul cohomology to give a beautiful generalization of the classical results above: if $\deg(D) \geq 2g + p + 1$, then D satisfies N_p .

In this brief note, we investigate the N_p property for toric varieties. For any divisor D and variety X such that R is arithmetically Cohen-Macaulay, it is natural to slice with hyperplanes until X has been reduced to a curve, and then apply Green's theorem. Results of Hochster [8] show that projectively normal toric varieties are always arithmetically Cohen-Macaulay. So it makes sense to apply the technique in this setting. In [4], Ewald and Wessels prove that if D is an ample divisor on a toric variety of dimension n , then $(n - 1)D$ is very ample and satisfies N_0 . Bruns, Gubeladze and Trung [2] give another proof and also show that nD satisfies property N_1 . While it is often difficult to determine if a given divisor satisfies N_0 , for a lattice polygon P and corresponding divisor on a toric surface, the property N_0 holds "for free".

Received by the editors April 10, 2002 and, in revised form, August 26, 2003.

2000 *Mathematics Subject Classification*. Primary 14M25; Secondary 14J30, 52B35.

Key words and phrases. Toric variety, Green's theorem, free resolution, syzygy.

The author was supported in part by NSA Grant #MDA904-03-1-0006.

In [6] Gallego and Purnaprajna give criteria for the N_p property for smooth rational surfaces. Toric varieties are rational, and in the case of smooth surfaces the result we obtain is a toric restatement of the result in [6]. However, the proof is simpler in the toric case, applies to singular surfaces, and extends several results in the toric literature. For example, in [10] Koelman proves that a toric surface defined by P satisfies N_1 iff ∂P contains at least four lattice points, and Ewald and Schmeink [3] prove that certain polytopes associated to smooth toric varieties with $\text{Pic}(X) = 2$ satisfy N_1 .

Theorem 1.1. *Let P be an n -dimensional lattice polytope, and X, D the associated projective toric variety and ample divisor; so $P = H^0(\mathcal{O}_X(D))$. If D satisfies N_0 , then D satisfies N_p if P satisfies*

$$\sum_{\text{facets } F_i} \text{vol}(F_i) \geq n(n-2)\text{vol}(P) + \frac{p+3}{(n-1)!}.$$

Proof. Hochster's results mentioned earlier show that R is arithmetically Cohen-Macaulay. In [9], Khovanskii shows that a toric variety X defined by a lattice polytope P is *normal* iff the Hilbert polynomial of X and the Ehrhart polynomial of P agree. Projective normality implies normality, and so X is normal. Hence, the singular locus of X is of codimension at least two. So a general member of $|D|$ is smooth. Slicing with $n-1$ general hyperplanes, we obtain a smooth curve C with the same minimal free resolution as X . By Khovanskii's result,

$$\chi(\mathcal{O}_X(mD)) = |mP \cap \mathbb{Z}^n| = am^n + bm^{n-1} + \dots.$$

After slicing with $n-1$ general hyperplanes, the resulting curve C has

$$\chi(\mathcal{O}_C(m)) = n!am + (n-1)!b - (n-1)! \binom{n}{2} a.$$

The first two coefficients of the Ehrhart polynomial are

$$\begin{aligned} a &= \text{vol}(P), \\ b &= \frac{1}{2} \sum_{\text{facets } F_i} \text{vol}(F_i). \end{aligned}$$

Thus, applying Green's theorem, the divisor D associated to P satisfies N_p if

$$\sum_{\text{facets } F_i} \text{vol}(F_i) \geq n(n-2)\text{vol}(P) + \frac{p+3}{(n-1)!}.$$

□

2. APPLICATIONS

In [12], Wills shows that an n -dimensional lattice polytope P that contains an interior point satisfies $n \cdot \text{vol}(P) \geq \sum_{\text{facets}} \text{vol}(F_i)$. So at first glance the bound above seems useless. However, when $n=2$ the term $n(n-2)\text{vol}(P)$ vanishes, and by [4] the divisor associated to a lattice polygon P satisfies N_0 . So we obtain:

Corollary 2.1. *The divisor D associated to a lattice polygon P satisfies N_p if*
 $\# \text{ integral points in } \partial P \geq p+3.$

Example 2.2. If P is the unit lattice two-simplex, then dP defines the d -uple Veronese embedding of \mathbb{P}^2 . By Corollary 2.1, dP satisfies N_p if $p \leq 3d-3$, recovering a result of [1]. In fact, Ottaviani and Paoletti [11] show that this bound is tight.

Example 2.3. The ideal sheaf of a projective toric surface X is two-regular iff N_p holds for all $p \leq \text{codim}(X)$. By Corollary 2.1, this is true if P has no interior points. In this case R is level with a -invariant -2 , which gives half of Theorem 1.27 of [2]. If P has no interior points, then the corresponding divisor has arithmetic genus zero ([5], p. 91). Thus X is a surface of minimal degree. So if X is smooth, then it must be a rational normal scroll or the Veronese surface in \mathbb{P}^5 .

If P is three-dimensional, then P satisfies N_p if $2 \sum \text{vol}(F_i) - 6 \text{vol}(P) - 3 \geq p$ and N_0 holds. In order to obtain a useful bound, we require that P have no interior points, so that the Ehrhart polynomial evaluated at -1 is zero. For such a polytope, this implies that $\sum \text{vol}(F_i) = \#$ integral points in $P - 2$, which yields:

Corollary 2.4. *A lattice three-polytope P with no interior points satisfies N_p if D is projectively normal and $\#$ integral points in $P \geq 3 \text{vol}(P) + \frac{p+7}{2}$.*

Example 2.5. Polytopes corresponding to smooth torics with $\text{Pic}(X) = 2$ are studied in [3]; for threefolds there are only two families. Ewald and Schmei\nc show that the polytopes below satisfy N_1 :

$$\begin{aligned} P_1(a) &= \text{conv}\{\mathbf{0}, \mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3, \mathbf{e}_1 + (a+1)\mathbf{e}_3, \mathbf{e}_1 + (a+1)\mathbf{e}_2\}, \\ P_2(a, b) &= \text{conv}\{\mathbf{0}, \mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3, \mathbf{e}_1 + (a+1)\mathbf{e}_3, \mathbf{e}_2 + (b+1)\mathbf{e}_3\}. \end{aligned}$$

A calculation shows that

$$\begin{aligned} \text{vol}(P_1(a)) &= \frac{a^2+3a+3}{6}, \quad \# \text{ integral points in } P = \frac{a^2+5a+12}{2}, \\ \text{vol}(P_2(a, b)) &= \frac{a+b+3}{6}, \quad \# \text{ integral points in } P = a + b + 6. \end{aligned}$$

Thus, $P_1(a)$ satisfies N_p if $p \leq 2a + 2$, and $P_2(a, b)$ satisfies N_p if $p \leq a + b + 2$.

ACKNOWLEDGEMENT

I thank Greg Smith for useful comments.

REFERENCES

1. C. Birkenhake, *Linear systems on projective spaces*, Manuscripta Math. **88** (1995), 177–184. MR 96h:14003
2. W. Bruns, J. Gubeladze, N. Trung, *Normal polytopes, triangulations, and Koszul algebras*, J. Reine Angew. Math. **485** (1997), 123–160. MR 99c:52016
3. G. Ewald, A. Schmei\nc, *Representation of the Hirzebruch-Kleinschmidt varieties by quadrics*, Beitr\ncge Algebra Geom. **34** (1993), 151–156. MR 95b:14036
4. G. Ewald, U. Wessels, *On the ampleness of invertible sheaves in complete projective toric varieties*, Results Math. **19** (1991), 275–278. MR 92b:14028
5. W. Fulton, *Introduction to Toric Varieties*, Princeton University Press, Princeton N.J., 1993. MR 94g:14028
6. F. Gallego, B. Purnaprajna, *Some results on rational surfaces and Fano varieties*, J. Reine Angew. Math. **538** (2001), 25–55. MR 2002f:14024
7. M. Green, *Koszul cohomology and the geometry of projective varieties*, J. Differential Geometry **19** (1984), 125–171. MR 85e:14022
8. M. Hochster, *Rings of invariants of tori, Cohen-Macaulay rings generated by monomials, and polytopes*, Ann. of Math. **96** (1972), 318–337. MR 46:3511
9. A. Khovanskii, *The Newton polytope, the Hilbert polynomial and sums of finite sets*, Funct. Anal. Appl. **26** (1992), 276–281. MR 94e:14068
10. R. Koelman, *A criterion for the ideal of a projectively embedded toric surface to be generated by quadrics*, Beitr\ncge Algebra Geom. **34** (1993), 57–62. MR 94h:14051

11. G. Ottaviani, R. Paoletti, *Syzygies of Veronese embeddings*, *Compositio Math.* **125** (2001), 31–37. MR 2002g:13023
12. J. Wills, *On an analog to Minkowski's lattice point theorem. The geometric vein*, Springer, New York-Berlin (1981), 285–288. MR 84b:52014

DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TEXAS 77843
E-mail address: `schenck@math.tamu.edu`