

## CLOSED SIMILARITY LORENTZIAN AFFINE MANIFOLDS

TSEMO ARISTIDE

(Communicated by Ronald A. Fintushel)

ABSTRACT. A  $Sim(n-1, 1)$  affine manifold is an  $n$ -dimensional affine manifold whose linear holonomy lies in the similarity Lorentzian group but not in the Lorentzian group. In this paper, we show that a compact  $Sim(n-1, 1)$  affine manifold is incomplete. Let  $\langle \cdot, \cdot \rangle_L$  be the Lorentz form, and  $q$  the map on  $\mathbb{R}^n$  defined by  $q(x) = \langle x, x \rangle_L$ . We show that for a compact radiant  $Sim(n-1, 1)$  affine manifold  $M$ , if a connected component  $C$  of  $\mathbb{R}^n - q^{-1}(0)$  intersects the image of the universal cover of  $M$  by the developing map, then either  $C$  or a connected component of  $C - H$ , where  $H$  is a hyperplane, is contained in this image.

### INTRODUCTION

An  $n$ -dimensional affine manifold  $M$  is an  $n$ -dimensional differentiable manifold endowed with an atlas whose coordinate changes are locally affine maps. The affine structure of  $M$  pulls back to its universal cover  $\hat{M}$  and defines on it an affine structure determined by a local diffeomorphism  $D : \hat{M} \rightarrow \mathbb{R}^n$ , called the developing map. The developing map gives rise to a representation  $h : \pi_1(M) \rightarrow Aff(\mathbb{R}^n)$ , called the holonomy of the affine manifold. Its linear part  $L(h)$ , is called the linear holonomy of the affine manifold. We will say that the affine manifold is complete if and only if the developing map is a diffeomorphism. An  $n$ -affine manifold is said to be radiant if its holonomy fixes an element of  $\mathbb{R}^n$ .

We denote by  $O(p, q)$  the subgroup of linear automorphisms of  $\mathbb{R}^n$  that preserve a bilinear symmetric form of type  $p, q$ , and by  $Sim(p, q)$  the group generated by  $O(p, q)$  and the homotheties. An  $O(p, q)$  affine manifold  $M$  is an affine manifold  $M$  such that the image of its linear holonomy  $L(h)$  is a subgroup of  $O(p, q)$ . A  $Sim(p, q)$  affine manifold  $M$  is an affine manifold  $M$  such that the image of its linear holonomy  $L(h)$  is a subgroup of  $Sim(p, q)$  and contains an element that is not in  $O(p, q)$ .

Let us consider the flat riemannian torus  $T^n$ . Bieberbach has shown that closed  $O(n, 0)$  affine manifolds are finitely covered by  $T^n$ . Using the notion of discreteness, Yves Carrière has shown that closed  $O(n-1, 1)$  affine manifolds are complete. It is obvious that a  $Sim(n, 0)$  affine manifold is incomplete, since an element of its holonomy that does not lie in  $O(n, 0)$  fixes an element of  $\mathbb{R}^n$ . There exist examples of complete  $Sim(n-1, 1)$  affine manifolds. Let us give one: Endow  $\mathbb{R}^n$  with its

---

Received by the editors April 28, 2001.

2000 *Mathematics Subject Classification.* Primary 53C30, 53C50.

©2004 American Mathematical Society

basis  $(e_1, \dots, e_n)$  and with the Lorentzian product defined by

$$\langle e_i, e_i \rangle_L = 1; 0 < i < n; \langle e_i, e_j \rangle_L = 0; i \neq j; \langle e_n, e_n \rangle_L = -1.$$

We restrict this product to  $\mathbb{R}^2$ . The affine map whose linear part is

$$\begin{pmatrix} 2 & 0 \\ 0 & 1 \end{pmatrix}$$

in the basis  $(e_1 + e_2, e_1 - e_2)$ , and whose translation part is  $e_1 - e_2$  generates a group that acts properly and freely on  $\mathbb{R}^2$ .

The goal of this paper is to study closed  $Sim(n - 1, 1)$  affine manifolds. First we show:

**Theorem 1.** *A compact  $Sim(n - 1, 1)$  affine manifold is incomplete.*

After, using the notion of discompactity, we show

**Theorem 2.** *Let  $M$  be a closed radiant  $Sim(n - 1, 1)$  affine manifold. If a connected component  $C$  of  $\mathbb{R}^n - q^{-1}(0)$  intersects  $D(\hat{M})$ , then either  $C$  is contained in  $D(\hat{M})$  or a connected component of  $C - H$ , where  $H$  is a hyperplane.*

Interesting structures of  $Sim(n - 1, 1)$  affine manifolds can be constructed using the work of Goldman on projective structures on surfaces; see [Gol]. For instance, a  $Sim(2, 1)$  structure whose linear holonomy is Zariski dense in  $Gl(3, \mathbb{R})$  is given in this paper.

1. CLOSED  $Sim(n - 1, 1)$  AFFINE MANIFOLDS ARE INCOMPLETE

The main goal of this part is to show that a closed  $Sim(n - 1, 1)$  affine manifold cannot be complete.

Let us suppose that there exists a complete closed  $Sim(n - 1, 1)$  affine manifold  $M$ ;  $M$  is the quotient of  $\mathbb{R}^n$  by a subgroup of affine transformations  $\Gamma$ , whose linear part is contained in  $Sim(n - 1, 1)$ .

**Lemma 1.1.** *Let  $\gamma$  be an element of  $\Gamma$  whose linear part has a determinant  $< 1$ . Then there exists a basis  $(e_1, \dots, e_n)$  of  $\mathbb{R}^n$  such that the linear part of  $\gamma$  in this basis has the following form:*

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & \frac{1}{\lambda^2} & 0 \\ 0 & 0 & \frac{1}{\lambda} B'' \end{pmatrix}$$

where  $\lambda$  is a real number strictly superior to 1 in absolute value, and  $B''$  is a matrix that preserves the restriction of a euclidean product to the vector subspace generated by  $e_3, \dots, e_n$ .

*Proof.* We have supposed that the determinant of the linear part  $L(\gamma)$  of  $\gamma$  is strictly inferior to 1 in absolute value. This implies that there exists a real number  $\lambda > 1$  such that  $\lambda L(\gamma) = L(\gamma)'$ , where  $L(\gamma)'$  is an element of  $O(n - 1, 1)$ . The linear map  $L(\gamma)$  has 1 as an eigenvalue, since  $\gamma$  acts freely. We deduce that  $\lambda$  is an eigenvalue of  $L(\gamma)'$ . We remark that  $L(\gamma)'$  has another eigenvalue  $\alpha$  whose module is different from 1 and the module of  $\lambda$  since the absolute value of its determinant is 1. If  $\alpha$  is not a real number, then  $\alpha$  and its complex conjugate  $\bar{\alpha}$  are eigenvalues associated to the complex eigenvectors  $u_1$  and  $u_2$ . In this case the restriction of  $L(\gamma)'$  to the plane generated by  $u_1 + u_2$  and  $i(u_1 - u_2)$  is a euclidean similitude whose ratio is different from 1. This is impossible since  $L(\gamma)'$  lies in  $O(n - 1, 1)$ . Let  $v_1$  and  $v_2$

be the eigenvectors associated to  $\lambda$  and  $\alpha$ , and let  $\langle \cdot, \cdot \rangle_L$  be the Lorentzian product preserved by the linear holonomy. We have:

$$\langle v_1, v_1 \rangle_L = \langle v_2, v_2 \rangle_L = 0.$$

We deduce that the restriction of  $\langle \cdot, \cdot \rangle_L$  to the plane  $P$  generated by  $v_1$  and  $v_2$  is nondegenerate and has signature  $(1, 1)$ . This implies that the restriction of  $\langle \cdot, \cdot \rangle_L$  to the orthogonal  $W$  of  $P$  with respect to itself is a scalar product. The restriction  $B''$  of  $L(\gamma)'$  to  $W$  is an orthogonal linear map. We can suppose that its determinant is 1. We deduce that  $\alpha = \frac{1}{\lambda}$ .  $\square$

Up to a change of origin, we can suppose that  $\gamma(0) = (a_1, 0, \dots, 0)$  where  $a_1$  is a real number. The restriction  $B$  of  $L(\gamma)$  to the linear subspace generated by  $(e_2, \dots, e_n)$  is strictly contracting. It is easy to show that the group generated by  $\gamma$  is not cocompact. So  $\Gamma$  contains another element  $\gamma_1$  different from  $\gamma$ .

**Lemma 1.2.** *Let  $C$  be the linear part of  $\gamma_1$ . Then  $C(e_1) = e_1 + b$  where  $b$  lies in the linear subspace generated by  $e_2, \dots, e_n$ .*

*Proof.* Let  $k$  be an element of  $\mathbb{N}$ . Consider the element  $\gamma^k \gamma_1$ . Its linear part has 1 as an eigenvalue. The matrix of this linear part in the basis  $(e_1, \dots, e_n)$  is  $A^k C$ , where  $A$  is the matrix of the linear part of  $\gamma$ . Let  $u_k$  be an eigenvector of  $A^k C$  associated to 1. We assume that the norm of  $u_k$  with respect to the euclidean scalar product defined by  $\langle e_i, e_j \rangle = \delta_{ij}$  is 1. Let  $u_k = (u_k^1, u_k^2)$ . We have  $C(u_k) = (v_k^1, v_k^2)$ , where  $u_k^1$  and  $v_k^1$  are elements of  $\mathbb{R}$ , and  $u_k^2$  and  $v_k^2$  are elements of the vector space  $F$  generated by  $e_2, \dots, e_n$ . We have  $v_k^1 = u_k^1$  since  $A^k(e_1) = e_1$ , and  $A^k$  preserves  $F$ . Since  $B$  is strictly contracting, the norm of  $A^k(0, v_k^2)$  goes to 0 with respect to the euclidean norm. So  $u_k$  goes to  $e_1$ , and  $C(e_1)$ , which is the limit of  $C(u_k) = A^{-k}(u_k)$ , is  $e_1 + b$  where  $b$  is an element of  $F$ .  $\square$

*Proof of the Theorem 1.* Let  $c$  be the translational part of  $\gamma_1$  in the basis  $(e_1, \dots, e_n)$ . Put  $c = (c_1, \dots, c_n)$ . We have  $\gamma^k \circ \gamma_1 \circ \gamma^{-k}(0) = (c_1, B^k(-ka_1 b + (c_2, \dots, c_n)))$ . Since  $B$  is contracting and the action of  $\Gamma$  is proper, we deduce that  $(c_2, \dots, c_n) = b = 0$ . This will imply that the action of  $\Gamma$  preserves proper affine subspaces, which is impossible. See [FGH], Theorem 2.2.  $\square$

*Remark.* In contrast to the  $Sim(n, 0)$  affine manifolds (see [Fr], Theorem 1), there exist compact  $Sim(n - 1, 1)$  affine manifolds that are not radiant. Here is an example.

Endow  $\mathbb{R}^2$  with the Lorentzian product  $(\cdot, \cdot)$  such that  $(e_1, e_1) = (e_2, e_2) = 0$  and  $(e_1, e_2) = 1$ .

Consider the subgroup  $\Gamma$  of  $Aff(\mathbb{R}^2)$  generated by the following transformations:

$$\begin{aligned} \gamma_1(x, y) &= (x + 1, y), \\ \gamma_2(x, y) &= (x, 2y). \end{aligned}$$

The quotient of  $\mathbb{R} \times (\mathbb{R} - \{0\})$  by  $\Gamma$  is a compact  $Sim(n - 1, 1)$  affine manifold.

2. ON THE UNIVERSAL COVER OF COMPACT  $Sim(n - 1, 1)$  AFFINE MANIFOLDS

In this part we are going to find properties of the universal cover of a closed radiant  $Sim(n - 1, 1)$  affine manifold. We use the notion of discompactness defined by Carrière ([Car], 2.2.1). Let us recall it.

We consider in  $\mathbb{R}^n$  the unit ball  $B_n$ . The euclidean metric induces on closed subsets of  $\mathbb{R}^n$  the Hausdorff distance. Let  $G$  be a subgroup of  $Gl(n, \mathbb{R})$ , and let  $(g_p)_{p \in \mathbb{N}}$  be a sequence of elements of  $G$ . The limit of the family  $(g_p(B_n) \cap B_n)_{p \in \mathbb{N}}$  converges in  $B_n$ . It is a degenerated ellipsoid (see [Car]). The codimension of this ellipsoid is the discompacity  $d$ , of the family  $(g_p)_{p \in \mathbb{N}}$ . The discompacity of the group with respect to the euclidean metric is the smallest  $d$ .

Obviously we cannot use the notion of discompacity in this form since the linear holonomy of our manifold may contain homotheties. Denote  $q : \mathbb{R}^n \rightarrow \mathbb{R}$ ,  $x \rightarrow \langle x, x \rangle_L$ . We can define in  $\mathbb{R}^n - q^{-1}(0)$  the metric

$$(u, v) \longrightarrow \langle u, v \rangle' = \frac{\langle u, v \rangle_{euc}}{q(x)}$$

where  $u$  and  $v$  are vectors of the tangent space at  $x$ , and  $\langle \cdot, \cdot \rangle_{euc}$  is the euclidean scalar product.

**Theorem 2.1.** *Let  $\hat{x}$  be an element of  $\hat{M}$ ,  $u$  and  $v$  elements of  $T_{\hat{x}}\hat{M}$ , such that the geodesics  $c_1 : [0, 1] \rightarrow \hat{M}$ ,  $t \rightarrow \exp_{\hat{x}}(tu)$ , and the one  $c_2 : [0, 1] \rightarrow \hat{M}$ ,  $t \rightarrow \exp_{\hat{x}}(tv)$  are defined. Suppose that the elements  $\exp_{\hat{x}}(u)$  and  $\exp_{\hat{x}}(v)$  cannot be joined by a geodesic, but for every  $t, t' < 1$ , there is a geodesic between  $\exp_{\hat{x}}(tu)$  and  $\exp_{\hat{x}}(t'v)$ . Let  $c : [0, 1] \rightarrow \mathbb{R}^n$ ,  $t \rightarrow \exp_{D(\exp_{\hat{x}}(u))}(tw)$  be the geodesic between  $D(\exp_{\hat{x}}(u))$  and  $D(\exp_{\hat{x}}(v))$ , and let  $U_{\hat{x}}$  be the domain of definition of  $\exp_{\hat{x}}$ . Consider the element  $t_0 \in [0, 1]$  such that for every  $t < t_0$ ,  $\exp_{D(\exp_{\hat{x}}(u))}(tw)$  is an element of  $D(\exp_{\hat{x}}(U_{\hat{x}}))$ , but  $\exp_{D(\exp_{\hat{x}}(u))}(t_0w) = y$  is not an element of  $D(\exp_{\hat{x}}(U_{\hat{x}}))$ . Then  $y$  is an element of  $q^{-1}(0)$ .*

*Proof.* There is a geodesic  $\hat{c}_3 : [0, 1[ \rightarrow \hat{M}$ ,  $t \rightarrow \exp_{\hat{x}}(tb)$  such that  $y$  is an element of the adherence of  $D(\hat{c}_3([0, 1[))$  and such that  $D(\hat{c}_3)$  is contained in the convex hull of  $D(\hat{c}_1)$  and  $D(\hat{c}_2)$ , where  $\hat{c}_1$ , and  $\hat{c}_2$  are geodesics of  $\hat{M}$  respectively above  $c_1$  and  $c_2$ . Set  $p(\hat{x}) = x$ . The image  $c_3$  of  $p(\hat{c}_3)$  is a maximal incomplete geodesic of  $M$ . Since  $M$  is compact, there exists an element  $z$  of  $M$  such that the geodesic  $c_3$  is recurrent in an affine chart  $U$  that contains  $z$ . We deduce as in Carrière the existence of a family of ellipsoids  $s_p$  of  $\mathbb{R}^n$  whose centers are elements of  $D(\hat{c}_3)$ , such that for each  $p, p'$ , there is an element  $\gamma_{p,p'}$  of the holonomy such that  $\gamma_{p,p'}(s_p) = s_{p'}$  and the centers  $x_p$  of  $s_p$  go to  $y$ .

Suppose that  $y$  is not an element of  $q^{-1}(0)$ .

Let  $z_p$  be an element of an ellipsoid  $s_p$ , and let  $u_p, v_p$  be two vectors in its tangent space. Put  $\gamma_{p,p'} = \lambda_{p,p'} g_{p,p'}$  where  $g_{p,p'}$  is an element of  $O(n - 1, 1)$ . We have

$$\frac{\langle \gamma_{p,p'}(u_p), \gamma_{p,p'}(v_p) \rangle_{euc}}{q(\gamma_{p,p'}(x))} = \frac{\langle g_{p,p'}(u_p), g_{p,p'}(v_p) \rangle_{euc}}{q(x)},$$

since the holonomy of  $M$  is supposed to be radiant.

The metrics  $\langle \cdot, \cdot \rangle_{euc}$  and  $\langle \cdot, \cdot \rangle'$  are equivalent in a neighborhood of  $y$  since  $q(y)$  is different from 0. We know that the discompacity of the family of  $g_p$  with respect to the riemannian metric  $\langle \cdot, \cdot \rangle_{euc}$  is 1. The family of ellipsoids  $s_p$  goes to an ellipsoid, or a codimension 1 degenerated ellipsoid centered in  $y$ . We conclude as in Carrière that  $y$  must be an element of  $D(\exp_{\hat{x}}(U_{\hat{x}}))$ . This is not possible; so  $q(y) = 0$ .  $\square$

A similar result is given in [Gol].

**Corollary 2.2.** *Let  $M$  be a compact radiant  $Sim(n - 1, 1)$  affine manifold, and let  $\hat{x}$ ,  $u$ , and  $v$  be respectively elements of  $\hat{M}$  and  $T_{\hat{x}}\hat{M}$ , such that  $\exp_{\hat{x}}(u)$  and  $\exp_{\hat{x}}(v)$*

are defined. If the convex hull  $E$  of  $(D(\hat{x}), D(\exp_{\hat{x}}(u)), D(\exp_{\hat{x}}(v)))$  is contained in a connected component of  $\mathbb{R}^n - q^{-1}(0)$ , then it is contained in  $D(\exp_{\hat{x}}(U_{\hat{x}}))$ .

*Proof.* Suppose that  $E$  is not contained in  $D(\exp_{\hat{x}}(U_{\hat{x}}))$ . Let  $y$  and  $z$  be two elements of  $E \cap D(\exp_{\hat{x}}(U_{\hat{x}}))$  such that  $y = D(\exp_{\hat{x}}(u_1))$ ,  $z = D(\exp_{\hat{x}}(u_2))$ , and for every  $t_1, t_2 < 1$ ,  $\exp_{\hat{x}}$  is defined on the convex hull of  $0, tu_1, tu_2$ , but the elements  $\exp_{\hat{x}}(u_1)$  and  $\exp_{\hat{x}}(u_2)$  cannot be joined by a geodesic. Consider the geodesic  $c : [0, 1] \rightarrow \mathbb{R}^n$ ,  $t \rightarrow \exp_y(tw)$  between  $y$  and  $z$ . There exists a real number  $0 < t_0 < 1$ , such that for  $0 < t < t_0$ ,  $\exp_y(tw)$  lies in  $D(\exp_{\hat{x}}(U_{\hat{x}}))$ , but  $\exp_y(t_0w)$  does not lie in  $D(\exp_{\hat{x}}(U_{\hat{x}}))$ . We deduce from Theorem 2.1 that  $\exp_y(t_0w)$  must lie in  $q^{-1}(0)$ . This is contrary to the hypothesis.  $\square$

More generally, we can determine the boundary of the image of the developing map of a compact radiant  $Sim(n - 1, 1)$  affine manifold. More precisely, we have the following proposition, which implies Theorem 2.

**Proposition 2.3.** *Let  $M$  be a compact radiant  $Sim(n - 1, 1)$  affine manifold whose developing map is injective. Then the boundary of  $D(\hat{M})$  is contained in the union of  $q^{-1}(0)$  and a hyperplane.*

*Proof.* As in [Car], p. 625, one can remark that elements of the boundary of  $D(\hat{M})$  that are not elements of  $q^{-1}(0)$  are limits of  $(\gamma_n e)_{n \in \mathbb{N}}$ , where  $\gamma_n$  is an element of the holonomy and  $e$  is an ellipsoid. We conclude that those elements are contained in at most two hyperplanes  $H_1, H_2$ . The case of two hyperplanes is impossible, since those hyperplanes are stable by the holonomy. The affine function  $\alpha$  such that  $\alpha(H_1) = 0$  and  $\alpha(H_2) = 1$  will be invariant by the holonomy and so will define a differentiable function on  $M$  without a maximum. (It is the same argument used in [Car]).  $\square$

**Proposition 2.4.** *Let  $M$  be a compact radiant affine manifold. If the image of the developing map is a convex set contained in an open set of  $\mathbb{R}^n - q^{-1}(0)$ , then the developing map is injective.*

*Proof.* Let  $\hat{x}$  be an element of  $\hat{M}$ . For all elements  $u$  and  $v$  of  $U_{\hat{x}}$ , the convex hull of  $D(\hat{x})$  is a subset of  $D(\hat{M}) \cap (\mathbb{R}^n - q^{-1}(0))$ , where  $y = D(\exp_{\hat{x}}(u))$  and  $z = D(\exp_{\hat{x}}(v))$ . We deduce from Corollary 2.2 that  $y$  and  $z$  are elements of  $D(U_{\hat{x}})$ . This implies that  $U_{\hat{x}}$  is a convex set. We can conclude the proof by using [Kos].  $\square$

A particular case of the situation of Corollary 2.4 is the following: endow a compact oriented surface  $S$  of genus  $> 2$  with a hyperbolic structure, and consider  $q$ , the Lorentzian form defined on  $\mathbb{R}^3$ , by  $q(x_1, x_2, x_3) = x_1^2 + x_2^2 - x_3^2$ . The hyperbolic structure can be defined by a representation of the fundamental group of  $S$ ,  $\pi_1(S) \rightarrow O(2, 1)$  such that the quotient of  $H = q^{-1}(-1)$  by  $\pi_1(S)$  is  $S$ . The quotient of  $W = \{x : q(x) \langle 0, x_3 \rangle 0\}$  by the group generated by  $\pi_1(S)$ , and a homothety of ratio  $0 < \lambda < 1$ , is a compact  $Sim(n - 1, 1)$  affine manifold whose universal cover is  $W$ .

More generally we have

**Corollary 2.5.** *Let  $M$  be a radiant compact affine manifold such that the image of its developing map is contained in  $W = \{x : q(x) \langle 0, z \rangle 0\}$ .  $M$  is the quotient of a connected component of  $W - H$  by a discrete group of  $Sim(n - 1, 1)$ , where  $H$  is a hyperplane of  $\mathbb{R}^n$ .*

*Proof.* We remark that the interior of a connected component of  $W - H$  is convex. This implies that the image of the developing map is a convex set. The result follows by using Propositions 2.3 and 2.4.  $\square$

Let  $M$  be a compact radiant  $Sim(n - 1, 1)$  affine manifold. The foliation  $D(\hat{\mathcal{F}}_q)$  of  $\mathbb{R}^n - \{0\}$  whose leaves are the submanifolds defined by  $q = \text{constant}$  is invariant by the holonomy of  $M$ . Its pullback on  $\hat{M}$  defines a foliation  $\hat{\mathcal{F}}_q$  of  $\hat{M}$ , which gives rise to a foliation  $\mathcal{F}_q$  of  $M$ . If  $D(\hat{N}) = D(\hat{M}) \cap q^{-1}(0)$  is not empty, then  $N = p(D^{-1}(D(\hat{N})))$  is a compact submanifold of  $M$ . Note that the 1-parameter group  $\phi_t$ , generated by the radiant vector field, preserves the foliation  $\mathcal{F}_q$ , and is transverse to all the leaves but not to the connected components of  $N$ .

**Proposition 2.6.** *If  $N$  is empty, then  $M$  is the total space of a bundle over  $S^1$ .*

*Proof.* If  $N$  is empty, then  $\phi_t$  is transverse to the foliation  $\mathcal{F}_q$ . This implies that this foliation is a Lie foliation. We conclude the proof by using [God], Corollary 2.6, p. 154.  $\square$

*Remark.* If the image of the developing map is contained in  $\{x/q(x) < 0\}$ , one may expect that the fibers of the previous fibration are hyperbolic manifolds.

#### REFERENCES

- [Car] Carrière, Y. Autour de la conjecture de L. Markus sur les variétés affines. *Invent. Math.* 95, (1989) 615-628. MR 89m:53116
- [Fr] Fried, D. Closed similarity manifolds. *Comment. Math. Helvetici* 55, (1980) 576-582. MR 83e:53049
- [FGH] Fried, D. Goldman, W. Hirsch, M. Affine manifolds with nilpotent holonomy. *Comment. Math. Helvetici* 56, (1981) 487-523. MR 83h:53062
- [God] Godbillon, C. Feuilletages, études géométriques, I. *Progress. in Math.*, vol. 98 (1991). MR 93i:57038
- [Gol] Goldman, W. Projective structures with Fuchsian holonomy. *J. Differential Geometry* 25, (1987) 297-326. MR 88i:57006
- [Kos] Koszul, J-L. Variétés localement plates et convexité. *Osaka J. Math.* 2, (1965) 285-290. MR 33:4849
- [T] Tsemo, A. Thèse Université de Montpellier II, (1999).
- [Wo] Wolf, J. Spaces of constant curvature. McGraw-Hill, 1967. MR 36:829

THE INTERNATIONAL CENTER FOR THEORETICAL PHYSICS, STRADA COSTIERA, 11, TRIESTE, ITALY

*Current address:* 3738, Avenue de Laval, Appt. 106, Montreal, Canada H2X 3C9

*E-mail address:* tsemoaristide@hotmail.com