

FRAMES AND THE FEICHTINGER CONJECTURE

PETER G. CASAZZA, OLE CHRISTENSEN, ALEXANDER M. LINDNER,
AND ROMAN VERSHYNIN

(Communicated by David R. Larson)

ABSTRACT. We show that the conjectured generalization of the Bourgain-Tzafriri *restricted-invertibility theorem* is equivalent to the conjecture of Feichtinger, stating that every bounded frame can be written as a finite union of Riesz basic sequences. We prove that any bounded frame can at least be written as a finite union of linearly independent sequences. We further show that the two conjectures are implied by the *paving conjecture*. Finally, we show that Weyl-Heisenberg frames over rational lattices are finite unions of Riesz basic sequences.

1. INTRODUCTION

The purpose of this paper is to relate a large number of conjectures appearing in different branches of analysis. We state the exact conjectures later in this section, but in order to proceed we need some definitions.

A *frame* for a Hilbert space \mathcal{H} is a family of vectors $\{f_i\}_{i \in I}$ in \mathcal{H} so that there are constants $A, B > 0$ satisfying:

$$A\|f\|^2 \leq \sum_{i \in I} |\langle f, f_i \rangle|^2 \leq B\|f\|^2, \quad \text{for all } f \in \mathcal{H}.$$

The constants A and B are called *lower* and *upper frame bounds*, respectively. If we can choose $A = B$ we say that $\{f_i\}_{i \in I}$ is a *B-tight frame*. If at least the upper frame condition is satisfied we call $\{f_i\}_{i \in I}$ a *Bessel sequence*, with *Bessel constant* B . A sequence $\{f_i\}_{i \in I}$ in \mathcal{H} is *bounded* if $0 < \inf_{i \in I} \|f_i\| \leq \sup_{i \in I} \|f_i\| < \infty$. A bounded unconditional basis for \mathcal{H} is called a *Riesz basis* for \mathcal{H} . A sequence $\{f_i\}_{i \in I}$ which is a Riesz basis for its closed linear span in H is called a *Riesz basic sequence* in H . It is known that $\{f_i\}_{i \in I}$ is a Riesz basis for \mathcal{H} if and only if $\{f_i\}_{i \in I}$ is complete in \mathcal{H} and there are constants $A, B > 0$ so that for all finite families of scalars $\{a_i\}_{i \in I' \subset I}$ we have

$$A \sum_{i \in I'} |a_i|^2 \leq \left\| \sum_{i \in I'} a_i f_i \right\|^2 \leq B \sum_{i \in I'} |a_i|^2.$$

Received by the editors February 18, 2003 and, in revised form, July 3, 2003.

2000 *Mathematics Subject Classification*. Primary 46C05, 46L05; Secondary 42C40.

Key words and phrases. Kadison-Singer problem, paving conjecture, Feichtinger conjecture, frames.

The first author was supported by NSF DMS 0102686.

The last author thanks PIMS for support.

In this case we call A a *lower Riesz basis bound* of $\{f_i\}_{i \in I}$ and B an *upper Riesz basis bound*. For the basic properties of frames, Bessel sequences, Riesz sequences and Riesz basic sequences we refer the reader to [Ca, Ch, Y].

We now formulate the conjectures we will deal with in this paper. We begin with the original conjecture by Feichtinger:

Conjecture 1.1 (Feichtinger). *Every bounded frame can be written as a finite union of Riesz basic sequences.*

Given $N \in \mathbb{N}$, let ℓ_2^N denote \mathbb{C}^N equipped with the ℓ_2 -norm. We now state a conjecture concerning frames for ℓ_2^N , which we refer to as the *finite Feichtinger conjecture*.

Conjecture 1.2 (Finite Feichtinger Conjecture). *For every $B, C > 0$ there is a natural number $M = M(B, C)$ and an $A = A(B, C) > 0$ so that whenever $\{f_i\}_{i \in I}$ is a frame for ℓ_2^N ($N \in \mathbb{N}$) with upper frame bound B and $\|f_i\| \geq C$ for all $i \in I$, then I can be partitioned into $\{I_j\}_{j=1}^M$ so that for each $1 \leq j \leq M$, $\{f_i\}_{i \in I_j}$ is a Riesz basic sequence with lower Riesz basis bound A and upper Riesz basis bound B .*

The corresponding conjectures for Bessel sequences are:

Conjecture 1.3. *Every bounded Bessel sequence can be written as a finite union of Riesz basic sequences.*

Conjecture 1.4. *For every $B > 0$ there exists a natural number $M = M(B)$ and an $A = A(B)$ so that every Bessel sequence $\{f_i\}_{i=1}^n$ with Bessel constant $B > 0$ and $\|f_i\| = 1$, for all $1 \leq i \leq n$, can be written as a union of M Riesz basic sequences each with lower Riesz basis bound A .*

Throughout this paper, $\{e_i\}$ will denote an orthonormal basis for whatever Hilbert space we are working in. In 1987, Bourgain and Tzafriri [BT1] proved the following fundamental result known as the *Restricted-Invertibility Theorem*:

Theorem 1.5 (Bourgain-Tzafriri). *There is a universal constant $c > 0$ so that whenever $T : \ell_2^n \rightarrow \ell_2^n$ is a linear operator for which $\|Te_i\| = 1$ for $1 \leq i \leq n$, then there exists a subset $\sigma \subset \{1, 2, \dots, n\}$ of cardinality $|\sigma| \geq \frac{cn}{\|T\|^2}$ so that*

$$\left\| \sum_{j \in \sigma} a_j Te_j \right\|^2 \geq c \sum_{j \in \sigma} |a_j|^2,$$

for all choices of scalars $\{a_j\}_{j \in \sigma}$.

Theorem 1.5 gave rise to the following conjecture, which is still open today:

Conjecture 1.6. *For every $B > 0$ there is a natural number $M = M(B)$ and an $A = A(B) > 0$ so that if $T : \ell_2^n \rightarrow \ell_2^n$ is a linear operator for which $\|Te_i\| = 1$ for all $1 \leq i \leq n$, and $\|T\| \leq \sqrt{B}$, then there is a partition $\{I_j\}_{j=1}^M$ of $\{1, 2, \dots, n\}$ so that for each $1 \leq j \leq M$ and all choices of scalars $\{a_i\}_{i \in I_j}$ we have*

$$\left\| \sum_{i \in I_j} a_i Te_i \right\|^2 \geq A \sum_{i \in I_j} |a_i|^2.$$

We will show that Conjectures 1.1, 1.2 and 1.6 are equivalent in the sense that all three have positive answers or all three have negative answers. We will also show that these conjectures are equivalent to the corresponding conjectures about

Bessel sequences and that all of these are true if the well-known *Paving Conjecture* holds. Given a subset I of the integers, we denote by P_I the orthogonal projection in ℓ_2 onto the subspace spanned by $\{e_i\}_{i \in I}$.

Conjecture 1.7 (The Paving Conjecture [KS]). *For any $\varepsilon > 0$, there is a constant $M = M(\varepsilon)$ such that for every integer n and every linear operator S on ℓ_2^n whose matrix with respect to $\{e_i\}_{i=1}^n$ has zero diagonal, one can find a partition $\{\sigma_j\}_{j=1}^M$ of $\{1, \dots, n\}$ such that*

$$\|P_{\sigma_j} S P_{\sigma_j}\| \leq \varepsilon \|S\| \quad \text{for all } j = 1, 2, \dots, M.$$

The Paving Conjecture is known to be equivalent to the Kadison-Singer conjecture [KS] (see also [BT2] for a deep analysis of the paving conjecture). In an interesting paper [W], Weaver gives several reformulations of the Kadison-Singer conjecture and thus of the Paving Conjecture in terms of frames.

It is possible that all these conjectures have negative answers in general. So in Section 4 we will give conditions under which related results hold. We also prove that any Bessel sequence can at least be decomposed into a finite union of linearly independent sets. Finally we consider frames with a special structure, namely, Weyl-Heisenberg frames, and give a sufficient condition for the decomposition into a finite number of Riesz basic sequences.

2. EQUIVALENCE OF THE CONJECTURES

To simplify the proof of the main result of this section, we first prove an elementary proposition.

Proposition 2.1. *Fix a natural number M and assume for every natural number n we have a partition $\{I_i^n\}_{i=1}^M$ of $\{1, 2, \dots, n\}$. Then there are natural numbers $\{n_1 < n_2 < \dots\}$ so that if $j \in I_i^{n_j}$ for some $i \in \{1, \dots, M\}$, then $j \in I_i^{n_k}$, for all $k \geq j$. Hence, if $I_i = \{j | j \in I_i^{n_j}\}$, then*

- (1) $\{I_i\}_{i=1}^M$ is a partition of \mathbb{N} .
- (2) If $I_i = \{j_1 < j_2 < \dots\}$, then for every natural number k we have $\{j_1, j_2, \dots, j_k\} \subset I_i^{n_{j_k}}$.

Proof. For each natural number n , 1 is in one of the sets $\{I_i^n\}_{i=1}^M$. Hence, there are natural numbers $n_1^1 < n_2^1 < n_3^1 < \dots$ and an $1 \leq i \leq M$ so that $1 \in I_i^{n_j^1}$, for all $j \in \mathbb{N}$. Now, for every natural number n_j^1 , 2 is in one of the sets $\{I_i^{n_j^1}\}_{i=1}^M$. Hence, there is a subsequence $\{n_j^2\}$ of $\{n_j^1\}$ and an $1 \leq i \leq M$ so that $2 \in I_i^{n_j^2}$, for all $j \in \mathbb{N}$. Continuing by induction, we get a subsequence $\{n_j^{\ell+1}\}_{j=1}^\infty$ of $\{n_j^\ell\}_{j=1}^\infty$ and an $1 \leq i \leq M$ so that $\ell + 1 \in I_i^{n_j^{\ell+1}}$, for all $j \in \mathbb{N}$. Letting $\{n_j\}_{j=1}^\infty$ be $\{n_j^\ell\}_{j=1}^\infty$ gives the conclusion of the proposition. \square

We can now state the main result of this section.

Theorem 2.2. *Conjectures 1.1, 1.2, 1.3, 1.4 and 1.6 are all equivalent in the sense that either all four of these conjectures have positive answers or all four have negative answers.*

Proof. Conjecture 1.3 \Rightarrow Conjecture 1.1: This is obvious.

Conjecture 1.1 \Rightarrow Conjecture 1.4: We will prove the contrapositive. So we assume that Conjecture 1.4 fails. Then there is a constant $B > 0$ so that for every $M \in \mathbb{N}$ and for every $A > 0$ there is an $n = n(M, A) \in \mathbb{N}$, a finite-dimensional Hilbert space H and a Bessel sequence $\{f_i\}_{i=1}^n$ in H with Bessel constant B and $\|f_i\| = 1$, for all $1 \leq i \leq n$, and whenever we partition $\{1, 2, \dots, n\}$ into sets $\{I_j\}_{j=1}^M$, then there exists some $1 \leq \ell \leq M$ and a set of scalars $\{a_i\}_{i \in I_\ell}$ with

$$\left\| \sum_{i \in I_\ell} a_i f_i \right\|^2 \leq A \sum_{i \in I_\ell} |a_i|^2.$$

Now, for each $k \in \mathbb{N}$, we can choose a finite-dimensional Hilbert space H_k of dimension, say m_k , and letting $M = k$ and $A = 1/k$ above we can choose $n_k = n(k, 1/k)$ and $\{f_i^k\}_{i=1}^{n_k}$ satisfying the above conditions. Let

$$H = \left(\sum \oplus H_k \right)_{\ell_2}$$

and consider $\{f_i^k\}_{i=1, k=1}^{n_k, \infty}$ as elements of H . For each $k \in \mathbb{N}$, let $\{e_i^k\}_{i=1}^{m_k}$ be an orthonormal basis for H_k and consider $\{e_i^k\}_{i=1, k=1}^{m_k, \infty}$ as elements of H . Since $\{e_i^k\}_{i=1, k=1}^{m_k, \infty}$ is an orthonormal basis for H , the family $\{f_i^k\}_{i=1, k=1}^{n_k, \infty} \cup \{e_i^k\}_{i=1, k=1}^{m_k, \infty}$ is a family of norm one vectors in H with Bessel bound $B + 1$ and lower frame bound ≥ 1 and hence forms a frame for H . Fix $M, A > 0$ and assume we can partition this frame into M sets of Riesz basic sequences each with lower Riesz basis bound A . In particular, we can partition $\{f_i^k\}_{i=1, k=1}^{n_k, \infty}$ into M sets of Riesz basic sequences each with lower Riesz basis bound A . But, for all k with $k \geq M$ and $1/k \leq A$, $\{f_i^k\}_{i=1}^{n_k}$ cannot be partitioned into M sets each with lower Riesz basis bound $\geq A$, and hence $\{f_i^k\}_{i=1, k=1}^{n_k, \infty}$ cannot be partitioned this way. This shows that Conjecture 1.1 fails.

Conjecture 1.4 \Rightarrow Conjecture 1.2: Given $\{f_i\}$ as in Conjecture 1.2, the sequence $\left\{ \frac{f_i}{\|f_i\|} \right\}$ is a Bessel sequence in $\ell_2^{\mathbb{N}}$ with Bessel constant $\frac{B}{C^2}$. So Conjecture 1.2 follows from Conjecture 1.4, since every frame is automatically bounded from above.

Conjecture 1.2 \Rightarrow Conjecture 1.6: This is obvious.

Conjecture 1.6 \Rightarrow Conjecture 1.3: Let $\{f_i\}_{i=1}^\infty$ be a bounded Bessel sequence for an infinite-dimensional Hilbert space H with Bessel constant B . Without loss of generality we may assume that $\|f_i\| = 1$, for all $1 \leq i < \infty$. For each n , choose an n -dimensional Hilbert space H_n containing the span of $\{f_i\}_{i=1}^n$ and let $\{e_i^n\}_{i=1}^n$ be an orthonormal basis for H_n . Define $T_n : H_n \rightarrow H_n$ by $T_n e_i^n = f_i$. Then $\|T_n\| \leq \sqrt{B}$ and so, by assuming that Conjecture 1.6 has a positive answer, we can find a partition $\{I_j^n\}_{j=1}^M$, $M = M(B)$, of $\{1, 2, \dots, n\}$ so that for every $1 \leq j \leq k$, $\{f_i\}_{i \in I_j^n}$ is a Riesz basic sequence with lower Riesz basis bound $A = A(B)$. By Proposition 2.1, we can partition \mathbb{N} into sets $\{I_i\}_{i=1}^M$ so that if $I_i = \{j_1 < j_2 < \dots\}$, then for every natural number k we have that $\{j_1, j_2, \dots, j_k\} \subset I_i^{n_{j_k}}$. It follows that $\{f_{j_\ell}\}_{\ell=1}^k$ is a Riesz basic sequence with the same lower Riesz basis bound A for all $k \in \mathbb{N}$. Hence, $\{f_j\}_{j \in I_i}$ has lower Riesz basis bound A , for all $1 \leq i \leq M$. Also, B is an upper Riesz basis bound for all these sets. This shows that Conjecture 1.3 has a positive answer. \square

3. THE PAVING CONJECTURE

Kadison and Singer raised the problem, which is still open, whether every pure state on \mathbb{D} , the C^* -algebra of the diagonal operators on ℓ_2 , admits a unique extension to a (pure) state on $\mathcal{L}(\ell_2)$, the C^* -algebra of all bounded linear operators on ℓ_2 . The problem of Kadison and Singer reduces to (and is equivalent to) the Paving Conjecture [KS] (see also [DS]).

Proposition 3.1. *The Paving Conjecture implies Conjecture 1.4.*

Proof. Let $\{f_i\}_{i=1}^n$ be a unit norm Bessel sequence in ℓ_2^n with Bessel constant B . Define the linear operator T on ℓ_2^n by setting $Te_i = f_i$ for all i . Then $\|T\| \leq \sqrt{B}$. Consider the operator $S = T^*T - I$. Then the (i, j) -entry of the matrix of S is

$$\langle Se_i, e_j \rangle = \begin{cases} \langle f_i, f_j \rangle, & i \neq j, \\ 0, & i = j. \end{cases}$$

By the Paving Conjecture, for any $\varepsilon > 0$ there exists a number $M = M(\varepsilon)$ and a partition $\{\sigma_k\}_{k=1}^M$ of the set $\{1, \dots, n\}$ such that

$$\|P_{\sigma_k}SP_{\sigma_k}\| \leq \varepsilon\|S\| \quad \text{for all } x \in \ell_2^n \text{ and all } k.$$

Applying this with $\varepsilon = \frac{1}{2(B+1)}$, and noting that $\|S\| \leq B + 1$, we obtain

$$\|P_{\sigma_k}SP_{\sigma_k}x\| \leq \frac{1}{2}\|x\| \quad \text{for all } k.$$

Now,

$$\begin{aligned} \langle P_{\sigma_k}SP_{\sigma_k}x, x \rangle &= \langle P_{\sigma_k}(T^*T - I)P_{\sigma_k}x, x \rangle \\ &= \langle (T^*T - I)P_{\sigma_k}x, P_{\sigma_k}x \rangle \\ &= \langle T^*TP_{\sigma_k}x, P_{\sigma_k}x \rangle - \langle P_{\sigma_k}x, P_{\sigma_k}x \rangle \\ &= \langle TP_{\sigma_k}x, TP_{\sigma_k}x \rangle - \|P_{\sigma_k}x\|^2 \\ &= \|TP_{\sigma_k}x\|^2 - \|P_{\sigma_k}x\|^2. \end{aligned}$$

Hence

$$\frac{|\|TP_{\sigma_k}x\|^2 - \|P_{\sigma_k}x\|^2|}{\|x\|} = \frac{|\langle P_{\sigma_k}SP_{\sigma_k}x, x \rangle|}{\|x\|} \leq \|P_{\sigma_k}SP_{\sigma_k}x\| \leq \frac{1}{2}\|x\|.$$

In particular,

$$|\|TP_{\sigma_k}x\|^2 - \|P_{\sigma_k}x\|^2| \leq \frac{1}{2}\|P_{\sigma_k}x\|^2 \quad \text{for all } x \in \ell_2^n.$$

Thus,

$$\frac{1}{2}\|P_{\sigma_k}x\|^2 \leq \|TP_{\sigma_k}x\|^2 \leq \frac{3}{2}\|P_{\sigma_k}x\|^2 \quad \text{for all } x \in \ell_2^n.$$

By the definition of T , this implies that $\{f_i\}_{i \in \sigma_k}$ is a Riesz basic sequence with lower Riesz basis bound $1/2$ and upper Riesz basis bound $3/2$. The proposition is proved. \square

The Paving Conjecture is known to be true for various classes of operators T on ℓ_2^n ; see [BT2] for references. In particular, the Paving Conjecture is proved for the operators whose matrices have small entries, $O(1/\log^{1+\gamma} n)$ for some $\gamma > 0$.

Theorem 3.2 (Bourgain-Tzafriri). *Let $\varepsilon > 0$ and S be a linear operator on ℓ_2^n whose matrix has zero diagonal and all entries are bounded by $1/\log^{1+\gamma} n$ for some $\gamma > 0$. Then S satisfies the conclusion of the Paving Conjecture: there exists a partition $\{\sigma_k\}_{k \leq M}$ of the set $\{1, \dots, n\}$, where $M = M(\gamma, \varepsilon)$, and such that*

$$\|P_{\sigma_k} S P_{\sigma_k}\| \leq \varepsilon \|S\| \quad \text{for all } k.$$

Actually, the partition $\{\sigma_k\}$ constructed by Bourgain and Tzafriri is random, i.e. σ_k is the image of the interval $\{1, \dots, n/M\}$ under a random permutation π of the interval $\{1, \dots, n\}$; for such a partition, the conclusion holds with probability close to one.

Theorem 3.2 implies the positive answer to Conjecture 1.4 for sequences which are in a certain sense “well separated”. It is clear that similar statements hold for conjectures 1.2 and 1.6 as well.

Corollary 3.3. *Let $\{f_i\}_{i=1}^n$ be a Bessel sequence with Bessel constant $B > 0$ and with $\|f_i\| = 1$ for all i . Assume that*

$$|\langle f_i, f_j \rangle| \leq \frac{1}{\log^{1+\gamma} n} \quad \text{for all } i \neq j.$$

Then the sequence $\{f_i\}_{i=1}^n$ can be written as a union of $M = M(B, \gamma)$ Riesz basic sequences each with lower Riesz basis bound $1/2$ and upper Riesz basis bound $3/2$.

Proof. This follows from Theorem 3.2 with an argument analogous to that of Proposition 3.1. □

4. POSITIVE RESULTS

First we will show that bounded Bessel sequences can be decomposed into a finite union of linearly independent sets. For this, we need a result of Christensen and Lindner [CL].

Proposition 4.1. *Let $M \in \mathbb{N}$, I a finite subset of \mathbb{N} and let $\{f_i\}_{i \in I}$ be a sequence of nonzero elements in a Hilbert space. The following are equivalent:*

- (1) *I can be partitioned into M disjoint sets I_1, I_2, \dots, I_M so that each family $\{f_i\}_{i \in I_j}$ ($j = 1, 2, \dots, M$) is linearly independent.*
- (2) *For any nonempty subset $J \subset I$ we have*

$$\frac{|J|}{\dim \text{span}\{f_j\}_{j \in J}} \leq M.$$

We now can show:

Theorem 4.2. *Every Bessel sequence $\{f_i\}_{i \in I}$ with Bessel bound B and $\|f_i\| \geq C > 0$, for every $i \in I$, can be decomposed into $\lceil B/C^2 \rceil$ linearly independent sets.*

Proof. We proceed by way of contradiction. Assume that $\{f_i\}_{i \in I}$ is a sequence, with Bessel bound B and $\|f_i\| \geq C > 0$, which cannot be decomposed into $\lceil B/C^2 \rceil$ linearly independent sets. By Proposition 2.1, with the same reasoning as used for the implication “Conjecture 1.6 \implies Conjecture 1.3”, we can assume that I is finite. By Proposition 4.1 there is a finite subset $J \subset I$ so that

$$\frac{|J|}{\dim \text{span}\{f_j\}_{j \in J}} > \left\lceil \frac{B}{C^2} \right\rceil.$$

Now, $\{f_j/\|f_j\|\}_{j \in J}$ is a frame for its span with upper frame bound $B_J \leq \lceil \frac{B}{C^2} \rceil$. Denote the corresponding frame operator by S , i.e.

$$S : \text{span} \{f_j\}_{j \in J} \rightarrow \text{span} \{f_j\}_{j \in J}, \quad f \mapsto \sum_{j \in J} \langle f, f_j \rangle f_j.$$

It is known that S has exactly $\dim \text{span}\{f_j\}_{j \in J}$ eigenvalues (counted with multiplicity), that all these eigenvalues are positive and less than or equal to B_J , and that their sum equals $|J|$; see [Ch, Th. 1.2.1]. Thus it follows that the largest eigenvalue λ_{max} must satisfy

$$\lambda_{max} \geq \frac{|J|}{\dim \text{span}\{f_j\}_{j \in J}} > \left\lceil \frac{B}{C^2} \right\rceil.$$

But, $\lambda_{max} \leq B_J$ and so $B_J > \lceil \frac{B}{C^2} \rceil$, which is a contradiction. □

Next, we will show that, up to a logarithmic factor, the generalized Bourgain-Tzafriri invertibility theorem (and hence the finite Feichtinger conjecture) is true. Namely, we can iterate Theorem 1.5 to obtain:

Proposition 4.3. *There is a universal constant $c > 0$ and a function $d = d(\|T\|)$ so that whenever $T : \ell_2^n \rightarrow \ell_2^n$ ($n \geq 2$) is a linear operator for which $\|Te_i\| = 1$, for $1 \leq i \leq n$, then there is a partition $\{I_j\}_{j=1}^{\lfloor d \log n \rfloor}$ of $\{1, 2, \dots, n\}$ so that for each $1 \leq j \leq \lfloor d \log n \rfloor$ and all choices of scalars $\{a_i\}_{i \in I_j}$ we have*

$$(4.1) \quad \left\| \sum_{i \in I_j} a_i Te_i \right\|^2 \geq c \sum_{i \in I_j} |a_i|^2.$$

Proof. Let $0 < c < 1$ be as in Theorem 1.5, $0 < b = \frac{c}{\|T\|^2} < 1$ and let $d > 0$ be such that $\lfloor d \log n \rfloor > \frac{-\log n}{\log(1-b)}$ for all $n \geq 2$. By Theorem 1.5 we can find a set $I_1 \subset \{1, 2, \dots, n\}$ with $|I_1| \geq bn$ and satisfying inequality (4.1). Let $J_1 = \{1, 2, \dots, n\} \setminus I_1$. Choose a (possibly into) isometry $U_1 : \text{Rng } T|_{\ell_2^{J_1}} \rightarrow \text{span} \{e_i\}_{i \in J_1}$. Then $U_1 T : \ell_2^{J_1} \rightarrow \ell_2^{J_1}$ satisfies Theorem 1.5, so there is a set $I_2 \subset J_1$ with $|I_2| \geq b(n - |I_1|)$ and satisfying inequality (4.1). Continuing, there are disjoint sets $\{I_j\}_{j=1}^{\lfloor d \log n \rfloor}$ with

$$|I_j| \geq b(n - |I_1| - |I_2| - \dots - |I_{j-1}|)$$

and each I_j satisfies inequality (4.1). Denoting $a_k := \sum_{j=1}^k |I_j|$, we have $a_k \geq bn + (1-b)a_{k-1}$ for any $k \geq 2$, and with $a_1 \geq bn$ this shows that

$$\sum_{j=1}^{\lfloor d \log n \rfloor} |I_j| = a_{\lfloor d \log n \rfloor} \geq bn \sum_{j=0}^{\lfloor d \log n \rfloor - 1} (1-b)^j = n(1 - (1-b)^{\lfloor d \log n \rfloor}) > n - 1$$

by the definition of d . Hence $\sum_{j=1}^{\lfloor d \log n \rfloor} |I_j| = n$, completing the proof. □

We can obtain a slightly more general result, namely

Theorem 4.4. *There is a universal constant $c > 0$ and a $D = D(B)$ so that whenever $\{f_i\}_{i=1}^k$ is a Bessel sequence in an n -dimensional Hilbert space \mathcal{H} with $\|f_i\| = 1$ for all $1 \leq i \leq k$ and Bessel constant B , then there is a partition $\{I_j\}_{j=1}^{\lfloor D \log n \rfloor}$ of $\{1, 2, \dots, k\}$ so that for each $1 \leq j \leq \lfloor D \log n \rfloor$, $\{f_i\}_{i \in I_j}$ is a Riesz basic sequence with lower Riesz basis bound c .*

Proof. By Theorem 4.2, $\{f_i\}_{i=1}^k$ can be decomposed into $\lceil B \rceil$ linearly independent sets, each of which has dimension at most n . In particular, $k \leq \lceil B \rceil n$. Then if d denotes the constant appearing in Proposition 4.3, it suffices to choose D such that

$$\lfloor d \log(\lceil B \rceil n) \rfloor = \lfloor d \log \lceil B \rceil + d \log n \rfloor \leq \lfloor D \log n \rfloor \quad \text{for all } n \geq 2.$$

□

5. WEYL-HEISENBERG FRAMES

In this section we show that the Feichtinger conjecture is true for certain Weyl-Heisenberg frames. If $g \in L^2(\mathbb{R})$, $a, b > 0$ we define for all $m, n \in \mathbb{Z}$:

$$E_{mb}g(t) = e^{2\pi i m b t} g(t) \quad \text{and} \quad T_{na}g(t) = g(t - na).$$

If $\{E_{mb}T_{na}g\}_{m,n \in \mathbb{Z}}$ is a frame for $L^2(\mathbb{R})$, we call it a *Weyl-Heisenberg* or *Gabor frame*. Our purpose is to show that whenever ab is rational, a Weyl-Heisenberg frame can be written as a finite union of Riesz basic sequences. In [G], Gröchenig shows that frames with a certain “localization property” can always be written as finite unions of Riesz basic sequences. This includes the case of Weyl-Heisenberg frames when g lies in a certain *modulation space*. The latter assumption is not required in our approach at a cost of having to work with rational lattices.

Theorem 5.1. *Let $g \in L^2(\mathbb{R})$ and $0 < ab < 1$ with ab rational. If $\{E_{mb}T_{na}g\}_{m,n \in \mathbb{Z}}$ is a Weyl-Heisenberg frame for $L^2(\mathbb{R})$, then it can be written as a finite union of Riesz basic sequences.*

Proof. After a change of variables we may assume that $b = 1$ and $a = \frac{p}{q}$ with $p, q \in \mathbb{N}$. We first reduce the problem to the case of integer oversampling. Notice that

$$\begin{aligned} \{E_m T_{\frac{1}{q}n} g\}_{m,n \in \mathbb{Z}} &= \bigcup_{k=0}^{p-1} \{E_m T_{\frac{1}{q}(np+k)} g\}_{m,n \in \mathbb{Z}} \\ &= \bigcup_{k=0}^{p-1} \{E_m T_{\frac{p}{q}n + \frac{k}{q}} g\}_{m,n \in \mathbb{Z}}. \end{aligned}$$

Since each of the families $\{E_m T_{\frac{p}{q}n + \frac{k}{q}} g\}_{m,n \in \mathbb{Z}}$ is a Bessel sequence (and a frame for $k = 0$) we conclude that $\{E_m T_{\frac{1}{q}n} g\}_{m,n \in \mathbb{Z}}$ is a frame. In the rest of the proof we show that $\{E_m T_{\frac{1}{q}n} g\}_{m,n \in \mathbb{Z}}$ is a finite union of Riesz basic sequences. Since $\{E_m T_{\frac{p}{q}n} g\}_{m,n \in \mathbb{Z}} \subset \{E_m T_{\frac{1}{q}n} g\}_{m,n \in \mathbb{Z}}$ the conclusion of the theorem follows from here.

We use a result by Ron and Shen [RS] (see also [G, Th. 7.4.3]), stating that $\{E_{qm}T_n g\}_{m,n \in \mathbb{Z}}$ is a Riesz basic sequence. Now,

$$\begin{aligned} \{E_m T_{\frac{n}{q}} g\}_{m,n \in \mathbb{Z}} &= \bigcup_{k=0}^{q-1} \{E_m T_{\frac{nq+k}{q}} g\}_{m,n \in \mathbb{Z}} \\ &= \bigcup_{k=0}^{q-1} \{E_m T_{n + \frac{k}{q}} g\}_{m,n \in \mathbb{Z}} \\ &= \bigcup_{k=0}^{q-1} \bigcup_{j=0}^{q-1} \{E_{qm+j} T_{n + \frac{k}{q}} g\}_{m,n \in \mathbb{Z}}. \end{aligned}$$

By the commutator relations between the translation and modulation operators, $\{E_{qm+j}T_{n+\frac{k}{q}}g\}_{m,n\in\mathbb{Z}}$ is a Riesz basic sequence (with the same Riesz basis bounds as $\{E_{qm}T_n g\}_{m,n\in\mathbb{Z}}$) for all j, k , from which the result follows. \square

Remark 5.2. The proof of Theorem 5.1 shows that a frame $\{E_m T_{\frac{k}{q}} g\}_{m,n\in\mathbb{Z}}$ is a union of q^2 Riesz basic sequences.

REFERENCES

- [BT1] J. Bourgain and L. Tzafriri: *Invertibility of “large” submatrices with applications to the geometry of Banach spaces and harmonic analysis*. Israel J. Math., **57** (1987) 137–223. MR0890420 (89a:46035)
- [BT2] J. Bourgain, L. Tzafriri, *On a problem of Kadison and Singer*, J. Reine Angew. Math. **420** (1991), 1–43. MR1124564 (92j:46104)
- [Ca] P.G. Casazza, *The art of frame theory*. Taiwanese Journal of Math., **4** (2) (2000) 129–202. MR1757401 (2001f:42046)
- [Ch] O. Christensen, *An introduction to frames and Riesz bases*, Birkhäuser, 2002. MR1946982 (2003k:42001)
- [CL] O. Christensen and A.M. Lindner, *Decomposition of Riesz frames and wavelets into a finite union of linearly independent sets*, Lin. Alg. Appl. **355** no. 1 (2002), 147–160. MR1930142 (2003g:42057)
- [DS] K. Davidson, S. Szarek, *Local operator theory, random matrices and Banach spaces*, Handbook of the geometry of Banach spaces, Vol. I, 317–366, North-Holland, Amsterdam, 2001. MR1863696 (2004f:47002a)
- [G] K. Gröchenig, *Localized frames are finite unions of Riesz sequences*. Adv. Comp. Math. **18** no. 2-4 (2003), 149–157. MR1968117 (2004a:42044)
- [KS] R. Kadison and I. Singer, *Extensions of pure states*, American Jour. Math. **81** (1959), 547–564. MR0123922 (23:A1243)
- [RS] A. Ron and Z. Shen, *Weyl-Heisenberg systems and Riesz bases in $L^2(\mathbb{R}^d)$* , Duke Math. Jour. **89** (1997), 237–282. MR1460623 (98i:42013)
- [W] N. Weaver, *The Kadison-Singer conjecture in discrepancy theory*, Preprint.
- [Y] R. Young, *An introduction to nonharmonic Fourier series - revised first edition*, Academic Press, 2001. MR1836633 (2002b:42001)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MISSOURI-COLUMBIA, COLUMBIA, MISSOURI 65211

E-mail address: pete@math.missouri.edu

DEPARTMENT OF MATHEMATICS, TECHNICAL UNIVERSITY OF DENMARK, BUILDING 303, 2800 LYNGBY, DENMARK

E-mail address: Ole.Christensen@mat.dtu.dk

CENTER OF MATHEMATICAL SCIENCES, MUNICH UNIVERSITY OF TECHNOLOGY, BOLTZMANNSTR. 3, D-85747 GARCHING, GERMANY

E-mail address: lindner@mathematik.tu-muenchen.de

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA AT DAVIS, ONE SHIELDS AVENUE, DAVIS, CALIFORNIA 95016

E-mail address: vershynin@math.ucdavis.edu