

FOURIER TRANSFORMS HAVING ONLY REAL ZEROS

DAVID A. CARDON

(Communicated by Joseph A. Ball)

ABSTRACT. Let $G(z)$ be a real entire function of order less than 2 with only real zeros. Then we classify certain distribution functions F such that the Fourier transform $H(z) = \int_{-\infty}^{\infty} G(it)e^{izt} dF(t)$ has only real zeros.

1. INTRODUCTION

Pólya [13] suggested that determining the class of functions whose Fourier transforms have only real zeros would be a ‘rather artificial question’ if it were not for the Riemann Hypothesis. For $\Re(s) > 1$, the Riemann zeta function is defined by $\zeta(s) = \sum_{n=1}^{\infty} n^{-s}$. It has an analytic continuation, and the function

$$\xi(s) = \frac{1}{2}s(s-1)\pi^{-s/2}\Gamma(s/2)\zeta(s)$$

is entire. The Riemann Hypothesis states that all the zeros of $\xi(s)$ satisfy $\Re(s) = 1/2$. A proof of the Riemann Hypothesis would be a major advance for analytic number theory. Let $\Xi(z) = \xi(\frac{1}{2} + iz)$. It is well known (see Titchmarsh [18], chapter 10) that

$$\Xi(z) = \int_{-\infty}^{\infty} \Phi(x)e^{izx} dx$$

where

$$\Phi(x) = \sum_{n=1}^{\infty} \left(4n^4\pi^2 e^{9x/2} - 6n^2\pi e^{5x/2} \right) \exp(-n^2\pi e^{2x}).$$

In other words, the Riemann Hypothesis is true if and only if the Fourier transform $\Xi(z)$ has only real zeros.

Pólya wrote several papers (such as [11, 12, 13, 14], which can all be found in [15]) in which he studied the reality of the zeros of various Fourier transforms. A particularly interesting result is the following:

Proposition 1 (Pólya [14]). *Let f be an integrable function of a real variable t such that $f(t) = \overline{f(-t)}$ and $f(t) = O(e^{-|t|^b})$ for $t \rightarrow \pm\infty$ and $b > 2$. Assume that*

$$\int_{-\infty}^{\infty} f(t)e^{izt} dt$$

Received by the editors September 23, 2003 and, in revised form, December 23, 2003.
2000 *Mathematics Subject Classification*. Primary 42A38, 30C15.

Key words and phrases. Fourier transform, zeros of entire functions, Laguerre-Pólya class.

©2004 American Mathematical Society
Reverts to public domain 28 years from publication

has only real zeros. Let ϕ be a real entire function having only real zeros, and assume that ϕ has a Weierstrass product of the form

$$\phi(z) = cz^m e^{\alpha z - \beta z^2} \prod_k \left(1 - \frac{z}{\alpha_k}\right) e^{z/\alpha_k}$$

where c, α, α_k are real, $\beta \geq 0$, and m is a nonnegative integer. (In other words, ϕ belongs to the Laguerre-Pólya class.) Then

$$\int_{-\infty}^{\infty} \phi(it) f(t) e^{izt} dt$$

has only real zeros.

In this paper we are concerned with constructing Fourier transforms with only real zeros in which the measure is *not* assumed to be the ordinary Lebesgue measure dt . The main result is the following theorem:

Theorem 1. *Suppose G is an entire function of order < 2 that is real on the real axis and has only real zeros. Let $\{a_k\}$ be a nonincreasing sequence of positive real numbers, let $\{X_k\}$ be the sequence of independent random variables such that X_k takes values ± 1 with equal probability, and let F_n be the distribution function of the normalized sum $Y_n = (a_1 X_1 + \dots + a_n X_n)/s_n$ where $s_n^2 = a_1^2 + \dots + a_n^2$. The functions F_n converge pointwise to a continuous distribution $F = \lim_{n \rightarrow \infty} F_n$. Let H be the Fourier transform of $G(it)$ with respect to the measure dF . That is,*

$$H(z) = \int_{-\infty}^{\infty} G(it) e^{izt} dF(t).$$

Then H is an entire function of order ≤ 2 that is real on the real axis. If H is not identically zero, then H has only real zeros.

Theorem 1 includes cases not covered in Proposition 1 because the distribution function $F(t)$, although continuous, need not be differentiable. If $F(t)$ is differentiable, we may write

$$\int_{-\infty}^{\infty} G(it) e^{izt} dF(t) = \int_{-\infty}^{\infty} G(it) e^{izt} F'(t) dt.$$

However, since not all functions $f(t)$ in Proposition 1 are of the form $f(t) = F'(t)$ for the types of distributions in Theorem 1, Proposition 1 covers cases not included in Theorem 1. So, while there is some overlap between Proposition 1 and Theorem 1, neither implies the other.

The proof of Theorem 1 is given in §2. Before proceeding with the proof we mention that the proof relies on a result about sums of exponential functions. Let $h_n(z)$ be the function of a complex variable z defined by

$$h_n(z) = \sum G(\pm ia_1 \pm \dots \pm ia_n) e^{iz(\pm b_1 \pm \dots \pm b_n)}$$

where the summation is over all 2^n possible sign combinations, the same sign combination being used in both the argument of G and in the exponent. The numbers a_1, a_2, a_3, \dots and b_1, b_2, b_3, \dots are positive, and G is as in Theorem 1. The author shows in [3] that all the zeros of the exponential sum $h_n(z)$ are real. Interestingly, the proof of this fact is similar to the proof of the Lee-Yang Circle Theorem from statistical mechanics (cf. [8] Appendix II or [17] Chapter 5). It should be pointed

out that this result of the author is related to de Branges' Hilbert spaces of entire functions [5]. Let $a_k = b_k$, $s_n^2 = a_1^2 + \cdots + a_n^2$, and

$$H_n(z) = 2^{-n} \sum G((\pm ia_1 \pm \cdots \pm ia_n)/s_n) e^{iz(\pm a_1 \pm \cdots \pm a_n)/s_n}.$$

All of the zeros of $H_n(z)$ are real. Theorem 1 is established by showing that the limit

$$H(z) = \lim_{n \rightarrow \infty} H_n(z)$$

is uniform on compact sets.

2. PROOF OF THEOREM 1

The proof of Theorem 1 is presented in this section as a sequence of lemmas. We begin with some notation.

The Laguerre-Pólya class \mathcal{LP} of functions consists of the entire functions having only real zeros with a Weierstrass factorization of the form

$$az^q e^{\alpha z - \beta z^2} \prod (1 - z/\alpha_n) e^{z/\alpha_n}$$

where a, α, β are real, $\beta \geq 0$, q is a nonnegative integer, and the α_n are nonzero real numbers such that $\sum \alpha_n^{-2} < \infty$. We shall be most interested in the subset \mathcal{LP}^* of the Laguerre-Pólya class consisting of all elements of \mathcal{LP} of order < 2 . Thus, β is necessarily 0 for functions in \mathcal{LP}^* .

The distribution function T for a random variable Y is $T(x) = \Pr(Y \leq x)$. We will consider the following types of random variables and their distribution functions: Let $\{a_k\}$ be a nonincreasing sequence of positive real numbers. Let $\{X_k\}$ be a sequence of independent random variables such that X_k takes values ± 1 with equal probability. Let Y_n be the sum

$$Y_n = \frac{a_1 X_1 + \cdots + a_n X_n}{s_n}$$

where $s_n^2 = a_1^2 + \cdots + a_n^2$. F_n will denote the distribution function of Y_n , and F will denote the limit $F = \lim_{n \rightarrow \infty} F_n$. In Theorem 1 the distribution F has variance 1. However, F could be rescaled to have any other positive value for its variance. The following lemma describes this F .

Lemma 1. *The sequence F_n converges pointwise to a continuous distribution F . If the sequence s_n is unbounded, F is the normal distribution. If the sequence s_n is bounded, F is not the normal distribution.*

Proof. This is proved in Lemma 1 of [2]. □

Lemma 2 (Pólya [12], Hilfssatz II). *Let a be a positive constant, let b be real, and let $G(z)$ be an entire function of genus 0 or 1 that for real z takes real values, has at least one real zero, and has only real zeros. Then the function*

$$e^{ib}G(z + ia) + e^{-ib}G(z - ia)$$

has only real zeros.

Proof. Pólya's original statement is Hilfssatz II in [12]. Pólya's argument is reiterated as Proposition 2 in [4]. □

We need the following important fact about $H_n(z)$.

Lemma 3. *Suppose $G \in \mathcal{LP}^*$. Let $a_1, \dots, a_n, b_1, \dots, b_n$ be positive real numbers. The exponential sum*

$$h_n(z) = \sum G(\pm ia_1 \pm \dots \pm ia_n) e^{iz(\pm b_1 \pm \dots \pm b_n)}$$

obtained by summing over all sign combinations, the same combination being used in the argument of G as in the exponent, is in \mathcal{LP}^ .*

Proof. It is clear that $h_n(z)$ is real for real z and has order 1. The fact that $h_n(z)$ has real zeros is proved in [3] by a method similar to that of the Lee-Yang Circle Theorem (found in Appendix II in [8]). □

If $s_n^2 = a_1^2 + \dots + a_n^2$ and if we use the notation of Riemann-Steiltjes integration, an immediate corollary to Lemma 3 is the following:

Corollary 4. *The function*

$$H_n(z) = 2^{-n} \sum G((\pm ia_1 \pm \dots \pm ia_n)/s_n) e^{iz(\pm a_1 \pm \dots \pm a_n)/s_n} = \int_{-\infty}^{\infty} G(it) e^{it} dF_n(t)$$

is in \mathcal{LP}^ .*

In Lemmas 6 and 7 we will show that the integrals $\int_{-\infty}^{\infty} G(it) e^{it} dF_n(t)$ converge uniformly to $\int_{-\infty}^{\infty} G(it) e^{it} dF_n(t)$ for z in compact sets. The proof of Lemma 6 will require the following 1994 result of Pinelis, which is an improvement of a conjecture by Eaton [6]:

Lemma 5 (Pinelis [10], Corollary 2.6). *Let X_k be independent random variables taking values ± 1 with equal probability. Let $s_n^2 = a_1^2 + \dots + a_n^2$, and let*

$$Y_n = \frac{a_1 X_1 + \dots + a_n X_n}{s_n}.$$

Then

$$\Pr(|Y_n| > u) < 2c(1 - \Phi(u))$$

where $c = 2e^3/9$, $\Phi(u) = \int_{-\infty}^u \phi(t) dt$, and $\phi(t) = (2\pi)^{-1/2} e^{-t^2/2}$.

Lemma 6. *Let $\epsilon > 0$ be given, suppose $G \in \mathcal{LP}^*$, and let K be a compact subset of \mathbb{C} . Then there is a positive number A (depending on ϵ and K) such that*

$$\int_{|t|>A} |G(it) e^{izt}| dF_n(t) < \epsilon$$

for all n and all $z \in K$.

Proof. Let λ denote the order of G . By hypothesis, $\lambda < 2$. Choose δ with $\max(1, \lambda) < \delta < 2$. Then choose $A > 0$ large enough so that $|G(it) e^{izt}| < e^{|t|^\delta}$ for all $z \in K$ and $|t| > A$. Such an A exists as follows: Choose δ' with $\lambda < \delta' < \delta$. Then for sufficiently large A , $|G(it)| < e^{|t|^{\delta'}}$ whenever $|t| > A$. Since K is compact, there is an R so that $|z| < R$ for all $z \in K$. For sufficiently large A and $|t| > A$,

$$|G(it) e^{izt}| \leq |G(it)| e^{|z||t|} < e^{|t|^{\delta'} + R|t|} < e^{|t|^\delta}.$$

Thus, A exists as claimed. Then

$$\int_{A < |t| < B} |G(it)e^{izt}| dF_n(t) < 2 \int_A^B e^{t^\delta} dF_n(t).$$

After integration by parts the right-hand side becomes

$$\begin{aligned} & 2e^{B^\delta} (F_n(B) - 1) - 2e^{A^\delta} (F_n(A) - 1) - 2 \int_A^B (F_n(t) - 1) d(e^{t^\delta}) \\ & < 2e^{A^\delta} (1 - F_n(A)) + 2 \int_A^B (1 - F_n(t)) \delta t^{\delta-1} e^{t^\delta} dt. \end{aligned}$$

According to Lemma 5 for $t \geq A$ and if A is sufficiently large, then

$$1 - F_n(t) \leq \beta \int_t^\infty e^{-u^2/2} du < \frac{e^{-t^2/2}}{2}$$

where $\beta = 4e^3/(9\sqrt{2\pi})$. This gives

$$\int_{A < |t| < B} |G(it)e^{izt}| dF_n(t) < e^{A^\delta - A^2/2} + \int_A^B \delta t^{\delta-1} e^{t^\delta - t^2/2} dt$$

and

$$\int_{A < |t|} |G(it)e^{izt}| dF_n(t) < e^{A^\delta - A^2/2} + \int_A^\infty \delta t^{\delta-1} e^{t^\delta - t^2/2} dt.$$

For sufficiently large A the last integral is bounded above by $e^{A^\delta - A^2/2}$. So,

$$\int_{A < |t| < B} |G(it)e^{izt}| dF_n(t) < 2e^{A^\delta - A^2/2}.$$

The right-hand side of the last inequality can be made arbitrarily small for sufficiently large A . Therefore, we obtain $\int_{|t| > A} |G(it)e^{izt}| dF_n(t) < \epsilon$ as desired. \square

Lemma 7. *Let K be a compact subset of \mathbb{C} . Then*

$$\int_{-A}^A G(it)e^{izt} dF_n(t) \rightarrow \int_{-A}^A G(it)e^{izt} dF(t)$$

uniformly as $n \rightarrow \infty$ for $z \in K$.

Proof. By the Helly-Bray Theorem (see [9, p. 182] or [7, p. 298]) it is immediate that convergence occurs pointwise. We must, however, verify uniform convergence for $z \in K$.

Let $\epsilon > 0$ be given, and write $g_z(t) = G(it)e^{izt}$. Choose κ such that $\kappa > |g_z(t)|$ and $\kappa > |g'_z(t)|$ for all $z \in K$ and $t \in [-A, A]$. Integration by parts yields

$$\int_{-A}^A G(it)e^{itz} dF(t) = g_z(A)F(A) - g_z(-A)F(-A) - \int_{-A}^A F(t)g'_z(t) dt.$$

Then

$$\begin{aligned} & \left| \int_{-A}^A G(it)e^{itz} dF(t) - \int_{-A}^A G(it)e^{itz} dF_n(t) \right| \\ & \leq |g_z(A)| |F(A) - F_n(A)| + |g_z(-A)| |F(-A) - F_n(-A)| \\ & \quad + \int_{-A}^A |F(t) - F_n(t)| |g'_z(t)| dt \\ & \leq \kappa |F(A) - F_n(A)| + \kappa |F(-A) - F_n(-A)| + \kappa 2A \max_{t \in [-A, A]} |F(t) - F_n(t)|. \end{aligned}$$

Since the functions F_n and F are distributions functions such that F_n converges to the continuous distribution F pointwise on $[-A, A]$, F_n converges to F uniformly on $[-A, A]$. Thus for sufficiently large n and all $t \in [-A, A]$,

$$|F_n(t) - F(t)| < \min\left(\frac{\epsilon}{6A\kappa}, \frac{\epsilon}{3\kappa}\right).$$

Therefore, for all sufficiently large n and all $z \in K$,

$$\left| \int_{-A}^A G(it)e^{itz} dF(t) - \int_{-A}^A G(it)e^{itz} dF_n(t) \right| < \frac{\epsilon}{3} + \frac{\epsilon}{3} + \frac{\epsilon}{3} = \epsilon.$$

This shows that the convergence $\int_{-A}^A G(it)e^{itz} dF_n(t) \rightarrow \int_{-A}^A G(it)e^{itz} dF(t)$ is uniform as claimed. □

Lemma 8. *Suppose $G \in \mathcal{LP}^*$. Then $H(z) = \int_{-\infty}^{\infty} G(it)e^{izt} dF(t)$ is an entire function that is real for real z , and if it does not vanish identically, then it has only real zeros.*

Proof. Let $H_n(z) = \int_{-\infty}^{\infty} G(it)e^{izt} dF_n(t)$. By Lemmas 6 and 7, $H_n(z)$ converges uniformly to $H(z)$ on compact sets. Since $H_n(z)$ is real for real z , its limit $H(z)$ is real for real z . By Hurwitz's Theorem (see [1, Thm. 2, p. 178]), if $H(z)$ is not identically zero, its zeros are limit points of the zeros of the $H_n(z)$. Since, for each n , $H_n(z)$ has only real zeros, $H(z)$ also has only real zeros. □

Lemma 9. *Suppose $G \in \mathcal{LP}^*$. The order of $H(z) = \int_{-\infty}^{\infty} G(it)e^{izt} dF(t)$ is ≤ 2 .*

Proof. Choose δ with $\lambda < \delta < 2$ where λ is the order of G . Let M be large enough so that $|G(z)| < Me^{|z|^\delta}$ for all z . By applying Hölder's Inequality (see [16, p. 63]) we obtain

$$\begin{aligned} \left| \int_{-\infty}^{\infty} G(it)e^{-izt} dF(t) \right| & \leq \int_{-\infty}^{\infty} Me^{|t|^\delta + |z||t|} dF(t) \\ & \leq M \left(\int_{-\infty}^{\infty} e^{2|t|^\delta} dF(t) \right)^{1/2} \left(\int_{-\infty}^{\infty} e^{2|z||t|} dF(t) \right)^{1/2}. \end{aligned}$$

By Lemma 5 both integrals in the product converge. The first integral is independent of z . We will determine a bound for the second integral. Integration by parts

and an application of Lemma 5 give

$$\begin{aligned} \int_{-\infty}^{\infty} e^{2|z||t|} dF(t) &= 2 \int_0^{\infty} e^{2|z|t} dF(t) \\ &= 2e^{2|z|t} (F(t) - 1) \Big|_0^{\infty} + 2 \int_0^{\infty} 2|z|e^{2|z|t} (1 - F(t)) dt \\ &= 1 + 4|z| \int_0^{\infty} |z|e^{2|z|t} (1 - F(t)) dt \\ &\leq 1 + K|z| \int_0^{\infty} e^{2|z|t-t^2/2} dt \quad \text{where } K = \frac{16e^3}{9\sqrt{2\pi}}. \end{aligned}$$

Since

$$\int_0^{\infty} e^{2|z|t-t^2/2} dt \leq e^{2|z|^2} \int_{-\infty}^{\infty} e^{-(t-2|z|)^2/2} dt = \sqrt{2\pi} e^{2|z|^2},$$

we see that $\left| \int_{-\infty}^{\infty} G(it)e^{-izt} dF(t) \right|$ is bounded by a constant times $|z|e^{2|z|^2}$. Thus, the order of $\int_{-\infty}^{\infty} G(it)e^{-izt} dF(t)$ is ≤ 2 .

This completes the proof of Theorem 1. \square

REFERENCES

1. Lars V. Ahlfors, *Complex analysis: An introduction to the theory of analytic functions of one complex variable*, third ed., International Series in Pure and Applied Mathematics, McGraw-Hill Book Co., New York, 1978. MR0510197 (80c:30001)
2. David A. Cardon, *Convolution operators and zeros of entire functions*, Proc. Amer. Math. Soc. **130** (2002), no. 6, 1725–1734. MR1887020 (2002m:30006)
3. David A. Cardon, *Sums of exponential functions having only real zeros*, Manuscripta Math. (to appear).
4. David A. Cardon and Pace P. Nielsen, *Convolution operators and entire functions with simple zeros*, Number theory for the millennium, I (Urbana, IL, 2000), A K Peters, Natick, MA, 2002, pp. 183–196. MR1956225 (2003m:30012)
5. Louis de Branges, *Hilbert Spaces of Entire Functions*, Prentice-Hall Inc., Englewood Cliffs, N.J., 1968. MR0229011 (37:4590)
6. M. L. Eaton, *A probability inequality for linear combinations of bounded random variables*, Ann. Stat. **2** (1974), 609–614.
7. Martin Eisen, *Introduction to mathematical probability theory*, Prentice-Hall Inc., Englewood Cliffs, N.J., 1969. MR0258078 (41:2725)
8. T. D. Lee and C. N. Yang, *Statistical theory of equations of state and phase transitions. II. Lattice gas and Ising model*, Physical Rev. (2) **87** (1952), 410–419. MR0053029 (14:711c)
9. Michel Loève, *Probability theory*, third ed., D. Van Nostrand Co., Inc., Princeton, N.J.-Toronto, Ont.-London, 1963. MR0203748 (34:3596)
10. Iosif Pinelis, *Extremal probabilistic problems and Hotelling's T^2 test under a symmetry condition*, Ann. Statist. **22** (1994), no. 1, 357–368. MR1272088 (95m:62115)
11. George Pólya, *On the zeros of an integral function represented by Fourier's integral*, Messenger of Math. **52** (1923), 185–188.
12. George Pólya, *Bemerkung über die Integraldarstellung der Riemannschen ξ -Funktion*, Acta Math. **48** (1926), 305–317.
13. George Pólya, *On the zeros of certain trigonometric integrals*, J. London Math. Soc. **1** (1926), 98–99.
14. George Pólya, *Über trigonometrische Integrale mit nur reellen Nullstellen*, J. Reine Angew. Math. **158** (1927), 6–18.
15. George Pólya, *Collected papers*, The MIT Press, Cambridge, Mass.-London, 1974, Vol. II: Location of zeros, Edited by R. P. Boas, Mathematicians of Our Time, Vol. 8. MR0505094 (58:21342)
16. Walter Rudin, *Real and complex analysis*, third ed., McGraw-Hill Series in Higher Mathematics, McGraw-Hill Book Co., New York, 1987. MR0924157 (88k:00002)

17. David Ruelle, *Statistical mechanics: Rigorous results*, W. A. Benjamin, Inc., New York-Amsterdam, 1969. MR0289084 (44:6279)
18. E. C. Titchmarsh, *The theory of the Riemann zeta-function*, second ed., Clarendon Press Oxford University Press, Oxford, 1986, Revised by D. R. Heath-Brown. MR0882550 (88c:11049)

DEPARTMENT OF MATHEMATICS, BRIGHAM YOUNG UNIVERSITY, PROVO, UTAH 84602
E-mail address: `cardon@math.byu.edu`