

A HEREDITARILY ℓ_1 SUBSPACE OF L_1 WITHOUT THE SCHUR PROPERTY

M. M. POPOV

(Communicated by N. Tomczak-Jaegermann)

ABSTRACT. Let $\infty > p_1 > p_2 > \cdots > 1$. We construct an easily determined 1-symmetric basic sequence in $\left(\sum_{n=1}^{\infty} \oplus \ell_{p_n}\right)_1$, which spans a hereditarily ℓ_1 subspace without the Schur property. An immediate consequence is the existence of hereditarily ℓ_1 subspaces of L_1 without the Schur property.

In our notation and terminology we follow mainly [4]–[6]. Recall that an infinite dimensional Banach space X is said to be hereditarily Y if each infinite dimensional subspace X_0 of X contains a further subspace $Y_0 \subseteq X_0$ which is isomorphic to Y . A Banach space X is said to have the Schur property provided weak convergence of sequences in X implies their norm convergence. It is well known that ℓ_1 has the Schur property. As was shown by J. Bourgain and H. P. Rosenthal in [3], there exists a subspace of L_1 having the Schur property, but which does not embed in ℓ_1 . The first example of a hereditarily ℓ_1 Banach space without the Schur property was constructed by J. Bourgain in [2]. Then P. Azimi and J. N. Hagler in [1] constructed a class of such spaces and investigated their further properties. We construct a class of subspaces of L_1 with the same properties. Actually, we make our construction in the sequence space

$$X = \left(\sum_{n=1}^{\infty} \oplus \ell_{p_n}\right)_1,$$

where $\infty > p_1 > p_2 > \cdots > 1$. Since X embeds isometrically in L_1 for $p_1 \leq 2$ (this can be easily deduced from [6, p. 212]), the construction gives examples of hereditarily ℓ_1 subspaces of L_1 without the Schur property. Of course, each of them must be uncomplemented since every complemented subspace of L_1 without the Schur property contains an isomorphic copy of ℓ_2 [7].

I would like to thank the referee for clarification of some arguments (the proof of Proposition 2 and especially for the idea of considering our example in the setting of the sequence space X , which makes the construction more general, clear and natural) and a number of corrections.

Received by the editors August 24, 2003 and, in revised form, February 26, 2004.
2000 *Mathematics Subject Classification*. Primary 46B20; Secondary 46E30.
Key words and phrases. Schur property, the space L_1 .

©2005 American Mathematical Society

THE CONSTRUCTION

Fix a sequence of reals $p_1 > p_2 > \dots > 1$, and consider the sequence space X defined above. For each $n \geq 1$, by $\{\bar{e}_{i,n}\}_{i=1}^\infty$ denote the unit vector basis of ℓ_{p_n} and by $\{e_{i,n}\}_{i=1}^\infty$ its natural copy in X :

$$e_{i,n} = (\underbrace{0, \dots, 0}_{n-1}, \bar{e}_{i,n}, 0, \dots) \in X.$$

Let $\delta_n > 0$ be such that $\sum_{n=1}^\infty \delta_n = 1$. For $i \geq 1$ put $z_i = \sum_{n=1}^\infty \delta_n e_{i,n}$. Evidently, $\|z_i\| = 1$ for each i .

Denote by Z the closed linear span of $\{z_i\}_{i=1}^\infty$. We show that Z possesses the desired properties. Since X embeds isometrically in L_1 for $p_1 \leq 2$, then so does Z , and our construction can be applied to subspaces of L_1 .

THE SKETCH

For each $I \subseteq \mathbb{N}$, by P_I we denote the natural projection of X onto $[e_{i,n} : i \in \mathbb{N}, n \in I]$ (i.e. with the kernel $[e_{i,n} : i \in \mathbb{N}, n \notin I]$). Of course, $\|P_I\| = \|Id - P_I\| = 1$. The main step here is to prove that Z is hereditarily ℓ_1 . Given an infinite dimensional subspace Z_0 of Z , we find a sequence $\{x_s\}_{s=1}^\infty$ in Z_0 and a block basic subsequence $\{u_s\}_{s=1}^\infty$ of $\{z_i\}_{i=1}^\infty$ having ‘‘almost disjoint supports’’ and which is close enough to $\{x_s\}_{s=1}^\infty$. (Here by ‘‘almost disjoint supports’’ we mean that for each $\varepsilon > 0$ there are disjoint subsets I_s of \mathbb{N} with $\|P_{I_s} u_s\| \geq (1 - \varepsilon)\|u_s\|$.) Hence $\{x_s\}_{s=1}^\infty$ contains a subsequence equivalent to the unit vector basis of ℓ_1 .

THE PROOF

Proposition 1. *For all scalars $\{a_i\}_{i=1}^m$ and each permutation of integers $\tau : \mathbb{N} \rightarrow \mathbb{N}$ one has*

$$\left\| \sum_{i=1}^m a_i z_{\tau(i)} \right\| = \sum_{n=1}^\infty \delta_n \left(\sum_{i=1}^m |a_i|^{p_n} \right)^{\frac{1}{p_n}}.$$

Hence, $\{z_i\}_{i=1}^\infty$ is a 1-symmetric basic sequence.

Proof. The proof is straightforward:

$$\left\| \sum_{i=1}^m a_i z_{\tau(i)} \right\| = \sum_{n=1}^\infty \delta_n \left\| \sum_{i=1}^m a_i e_{\tau(i),n} \right\| = \sum_{n=1}^\infty \delta_n \left(\sum_{i=1}^m |a_i|^{p_n} \right)^{\frac{1}{p_n}}.$$

□

Thus, if a series $\sum_{i=1}^\infty a_i z_i$ converges, then $\sum_{i=1}^\infty |a_i|^{p_n} < \infty$ for each n .

Proposition 2. $\{z_i\}_{i=1}^\infty$ tends weakly to zero.

Proof. Let $z^* \in Z^*$, $\|z^*\| = 1$, and $\varepsilon > 0$. Pick an integer N such that

$$\sum_{n=N+1}^\infty \delta_n < \frac{\varepsilon}{2}.$$

Since $e_{i,n} \rightarrow 0$ weakly as $i \rightarrow \infty$ for each $n \geq 1$, we may pick $K \in \mathbb{N}$ so that if $n = 1, \dots, N$ and $k \geq K$, then

$$|z^*(e_{k,n})| < \frac{\varepsilon}{2}.$$

Since $\sum_{n=1}^N \delta_n < 1$, we have that for $k \geq K$,

$$|z^*(z_k)| = \left| z^* \left(\sum_{n=1}^{\infty} \delta_n e_{k,n} \right) \right| < \sum_{n=1}^N \delta_n |z^*(e_{k,n})| + \sum_{n=N+1}^{\infty} \delta_n < \varepsilon.$$

□

Lemma 3. *Let Z_0 be an infinite dimensional subspace of Z , $n, m, j \in \mathbb{N}$ ($n > 1$) and $\varepsilon > 0$. Then there are $\{x_i\}_{i=1}^m \subset Z_0$ and $\{u_i\}_{i=1}^m \subset Z$ of the form*

$$u_i = \sum_{s=j_i+1}^{j_{i+1}} a_{i,s} z_s \text{ where } j = j_1 < j_2 < \dots < j_{m+1}$$

such that

$$\sum_{s=j_i+1}^{j_{i+1}} |a_{i,s}|^{p_{n-1}} = 1 \text{ and } \|u_i - x_i\| < \frac{\varepsilon}{m} \|u_i\|$$

for each $i = 1, \dots, m$.

Proof. Put $Z_1 = Z_0 \cap [z_i]_{i=j+1}^{\infty}$. Since Z_0 is infinite dimensional and $[z_i]_{i=j+1}^{\infty}$ has finite codimension in Z , Z_1 is infinite dimensional as well. Put $j_1 = j$ and choose any

$$\bar{x}_1 = \sum_{s=j_1+1}^{\infty} \bar{a}_{1,s} z_s \in Z_1 \setminus \{0\}.$$

Without loss of generality we may assume that

$$\sum_{s=j_1+1}^{\infty} |\bar{a}_{1,s}|^{p_{n-1}} = 1$$

(otherwise we multiply \bar{x}_1 by a suitable number). Then choose $j_2 > j_1$ so that for

$$\bar{u}_1 = \sum_{s=j_1+1}^{j_2} \bar{a}_{1,s} z_s$$

we have

$$\|\bar{u}_1 - \bar{x}_1\| < \frac{\varepsilon \|\bar{x}_1\|}{4m}, \quad \lambda_1 = \left(\sum_{s=j_1+1}^{j_2} |\bar{a}_{1,s}|^{p_{n-1}} \right)^{\frac{1}{p_{n-1}}} \geq \frac{1}{2} \quad \text{and} \quad \|\bar{u}_1\| \geq \frac{\|\bar{x}_1\|}{2}.$$

Hence,

$$\|\bar{u}_1 - \bar{x}_1\| < \frac{\varepsilon \|\bar{u}_1\|}{2m}.$$

Now put $a_{1,s} = \lambda_1^{-1} \bar{a}_{1,s}$, $x_1 = \lambda_1^{-1} \bar{x}_1$ and $u_1 = \lambda_1^{-1} \bar{u}_1$. Then

$$\sum_{s=j_1+1}^{j_2} |a_{1,s}|^{p_{n-1}} = \frac{1}{\lambda_1^{p_{n-1}}} \sum_{s=j_1+1}^{j_2} |\bar{a}_{1,s}|^{p_{n-1}} = 1$$

and

$$\|u_1 - x_1\| = \frac{1}{\lambda_1} \|\bar{u}_1 - \bar{x}_1\| < \frac{\varepsilon \|\bar{u}_1\|}{2\lambda_1 m} \leq \frac{\varepsilon \|\bar{u}_1\|}{m} \leq \frac{\varepsilon \|u_1\|}{m}.$$

Continuing the procedure in the obvious manner, we construct the desired sequences. □

For $n \in \mathbb{N}$ denote $Q_n = P_{\{n, n+1, \dots\}}$.

Lemma 4. *Let Z_0 be an infinite dimensional subspace of Z , $j, n \in \mathbb{N}$ and $\varepsilon > 0$. There exist an $x \in Z_0$, $x \neq 0$ and a $u \in Z$ of the form*

$$u = \sum_{i=j+1}^l a_i z_i, \quad \text{where } l > j,$$

such that

- (i) $\|Q_n u\| \geq (1 - \varepsilon) \|u\|$;
- (ii) $\|x - u\| < \varepsilon \|u\|$.

Proof. Choose m so that

$$\frac{1}{\delta_n} m^{\frac{1}{p_{n-1}} - \frac{1}{p_n}} < \varepsilon.$$

Using Lemma 3 choose $\{x_i\}_{i=1}^m \subset Z_0$ and $\{u_i\}_{i=1}^m \subset Z$ to satisfy the claims of the lemma and put

$$x = \sum_{i=1}^m x_i \quad \text{and} \quad u = \sum_{i=1}^m u_i.$$

First, we prove (ii). Since $\{z_s\}_{s=1}^\infty$ is 1-symmetric, then $\|u_i\| \leq \|u\|$ for $i = 1, \dots, m$ and

$$\|x - u\| \leq \sum_{i=1}^m \|x_i - u_i\| < \sum_{i=1}^m \frac{\varepsilon \|u_i\|}{m} \leq \sum_{i=1}^m \frac{\varepsilon \|u\|}{m} = \varepsilon \|u\|.$$

To prove (i), we first show that $\|u\| - \|Q_n u\| < m^{\frac{1}{p_{n-1}}}$. To see this,

$$\begin{aligned} \|u\| - \|Q_n u\| &= \sum_{k=1}^{n-1} \delta_k \left\| \sum_{i=1}^m P_{\{k\}} u_i \right\| = \sum_{k=1}^{n-1} \delta_k \left\| \sum_{i=1}^m \sum_{s=j_i+1}^{j_{i+1}} a_{i,s} e_{s,k} \right\| \\ &= \sum_{k=1}^{n-1} \delta_k \left(\sum_{i=1}^m \sum_{s=j_i+1}^{j_{i+1}} |a_{i,s}|^{p_k} \right)^{\frac{1}{p_k}} \leq \sum_{k=1}^{n-1} \delta_k \left(\sum_{i=1}^m \sum_{s=j_i+1}^{j_{i+1}} |a_{i,s}|^{p_{n-1}} \right)^{\frac{1}{p_{n-1}}} \\ &= \sum_{k=1}^{n-1} \delta_k \left(\sum_{i=1}^m 1 \right)^{\frac{1}{p_{n-1}}} = m^{\frac{1}{p_{n-1}}} \sum_{k=1}^{n-1} \delta_k < m^{\frac{1}{p_{n-1}}}. \end{aligned}$$

On the other hand,

$$\begin{aligned} \|u\| &= \sum_{k=1}^\infty \delta_k \left\| \sum_{i=1}^m \sum_{s=j_i+1}^{j_{i+1}} a_{i,s} e_{s,k} \right\| \geq \delta_n \left\| \sum_{i=1}^m \sum_{s=j_i+1}^{j_{i+1}} a_{i,s} e_{s,n} \right\| \\ &= \delta_n \left(\sum_{i=1}^m \sum_{s=j_i+1}^{j_{i+1}} |a_{i,s}|^{p_n} \right)^{\frac{1}{p_n}} \geq \delta_n \left(\sum_{i=1}^m \left(\sum_{s=j_i+1}^{j_{i+1}} |a_{i,s}|^{p_{n-1}} \right)^{\frac{p_n}{p_{n-1}}} \right)^{\frac{1}{p_n}} \\ &= \delta_n \left(\sum_{i=1}^m 1 \right)^{\frac{1}{p_n}} = \delta_n m^{\frac{1}{p_n}}. \end{aligned}$$

Hence

$$1 - \frac{\|Q_n u\|}{\|u\|} \leq \frac{1}{\delta_n} m^{\frac{1}{p_{n-1}} - \frac{1}{p_n}} < \varepsilon,$$

and $\|Q_n u\| \geq (1 - \varepsilon) \|u\|$. □

Theorem 5. *Z is hereditarily ℓ_1 and does not have the Schur property.*

Proof. By Proposition 2, Z does not have the Schur property. To prove that Z is hereditarily ℓ_1 , let Z_0 be an infinite dimensional subspace of Z . Using Lemma 4, construct inductively sequences $\{x_s\}_{s=1}^\infty \subset Z_0$, $\{u_s\}_{s=1}^\infty \subset Z$ of the form

$$u_s = \sum_{i=j_s+1}^{j_{s+1}} a_i z_i,$$

where $j_1 < j_2 < \dots$ and $\|u_s\| = 1$ and a sequence $1 \leq n_1 < n_2 < \dots$ so that

- (i) $\|Q_{n_s} u_s\| \geq \frac{7}{8},$
- (ii) $\|u_s - x_s\| \leq 2^{-s-1},$
- (iii) $\|Q_{n_{s+1}} u_s\| < \frac{1}{8}.$

To see that this can be done, let $j_1 = n_1 = 1$ and $\varepsilon_1 = \frac{1}{8}$. Choose by Lemma 4 an $x_1 \in Z \setminus \{0\}$ and

$$u_1 = \sum_{i=j_1+1}^{j_2} a_i z_i$$

such that $\|u_1\| = 1$, $\|Q_{n_1} u_1\| \geq 1 - \varepsilon_1 = \frac{7}{8}$ and $\|x_1 - u_1\| < \varepsilon_1 < \frac{1}{4}$. Then choose $n_2 > n_1$ so that $\|Q_{n_2} u_1\| < \frac{1}{8}$. Continuing the procedure in the obvious manner, we construct the desired sequences.

Evidently, (i) yields

$$(i') \quad \|u_s - Q_{n_s} u_s\| \leq \frac{1}{8}.$$

Conditions (i') and (iii) imply the equivalence of $\{u_s\}_{s=1}^\infty$ to the unit vector basis of ℓ_1 . Indeed, for $s \geq 1$ put $v_s = Q_{n_s} u_s - Q_{n_{s+1}} u_s$. Then $\|v_s\| \geq \frac{3}{4}$, and for each sequence of scalars $\{a_s\}_{s=1}^m$ one has

$$\left\| \sum_{s=1}^m a_s v_s \right\| = \sum_{s=1}^m |a_s| \|v_s\| \geq \frac{3}{4} \sum_{s=1}^m |a_s|,$$

and hence

$$\begin{aligned} \left\| \sum_{s=1}^m a_s u_s \right\| &\geq \left\| \sum_{s=1}^m a_s v_s \right\| - \left\| \sum_{s=1}^m a_s (u_s - v_s) \right\| \geq \frac{3}{4} \sum_{s=1}^m |a_s| - \left\| \sum_{s=1}^m a_s (u_s - Q_{n_s} u_s) \right\| \\ &+ \sum_{s=1}^m a_s Q_{n_{s+1}} u_s \geq \frac{3}{4} \sum_{s=1}^m |a_s| - \frac{1}{8} \sum_{s=1}^m |a_s| - \frac{1}{8} \sum_{s=1}^m |a_s| = \frac{1}{2} \sum_{s=1}^m |a_s|. \end{aligned}$$

Thus, $\{u_s\}_{s=1}^\infty$ is equivalent to the unit vector basis of ℓ_1 and so is $\{x_s\}_{s=1}^\infty$ by (ii) and the stability of basic sequences property [5, p. 5]. \square

Since for $p_1 \leq 2$ the space X embeds isometrically in L_1 (this can be easily deduced from [6, p. 212]), we obtain

Corollary 6. *L_1 contains hereditarily ℓ_1 subspaces without the Schur property.*

Finally, we remark that it does not matter whether p_n tends to 1 or not. Furthermore, if we repeat the construction with the space X replaced by $X_p = \left(\sum_{n=1}^{\infty} \oplus \ell_{p_n} \right)_p$, with $1 < p < \infty$ and an arbitrary decreasing sequence p_n (not necessarily satisfying $p \leq \inf_n p_n$), then Z becomes hereditarily ℓ_p , but in this case it is clear that Z does not have the Schur property.

REFERENCES

1. P. Azimi and J. N. Hagler, *Examples of hereditarily ℓ_1 Banach spaces failing the Schur property*, Pacif. J. Math. **122** (2)(1986), 287–297. MR0831114 (87f:46030)
2. J. Bourgain, *ℓ_1 -subspaces of Banach spaces*. Lecture notes. Free University of Brussels.
3. J. Bourgain and H. P. Rosenthal, *Martingales valued in certain subspaces of L^1* , Isr. J. Math. **37** (1-2) (1980), 54–75. MR0599302 (82g:46044)
4. W. B. Johnson and J. Lindenstrauss, *Basic concepts in the geometry of Banach spaces*. Handbook of the geometry of Banach spaces. Vol.I. W. B. Johnson and J. Lindenstrauss eds. Elsevier. Amsterdam. (2001), 1–84. MR1863689 (2003f:46013)
5. J. Lindenstrauss and L. Tzafriri, *Classical Banach spaces. I*, Springer-Verlag, Berlin-Heidelberg-New York, 1977. MR0500056 (58:17766)
6. J. Lindenstrauss and L. Tzafriri, *Classical Banach spaces. II*, Springer-Verlag, Berlin-Heidelberg-New York, 1979. MR0540367 (81c:46001)
7. H. P. Rosenthal, *Convolution by a biased coin*, Altgeld Book (Univ. of Illinois),II, 1975/76.

DEPARTMENT OF MATHEMATICS, CHERNIVTSI NATIONAL UNIVERSITY, STR. KOTSYBYN'SKOGO
 2, CHERNIVTSI, 58012 UKRAINE
E-mail address: `popov@chv.ukrpack.net`