

A TOPOLOGICAL PALEY-WIENER PROPERTY FOR LOCALLY COMPACT GROUPS

EBERHARD KANIUTH, ANTHONY T. LAU, AND GÜNTER SCHLICHTING

(Communicated by Andreas Seeger)

ABSTRACT. We investigate a certain topological Paley-Wiener property and show, for instance, that compact-free nilpotent groups and simply connected solvable groups share this property.

INTRODUCTION

Let f be a bounded and compactly supported measurable function on \mathbb{R}^n ($f \in L_c^\infty(\mathbb{R}^n)$). By the classical Paley-Wiener theorem, the Fourier transform \widehat{f} of f extends to an entire function on \mathbb{C}^n . It follows that $f = 0$ whenever \widehat{f} vanishes on a set of positive Lebesgue measure. If, more generally, G is a (second countable) unimodular locally compact group of type I and \widehat{G} denotes the dual space of G , then G is said to satisfy the *weak Paley-Wiener property* if the operator-valued Fourier transform $\pi \rightarrow \pi(f)$ of a non-zero function $f \in L_c^\infty(G)$ cannot vanish on a set of positive Plancherel measure on \widehat{G} . This weak Paley-Wiener property has been established by several authors for simply connected nilpotent Lie groups [1], [7], [14], [15] and later for completely solvable Lie groups [8].

In this paper we introduce and study another Paley-Wiener type property which can be defined for arbitrary locally compact groups G . We say that G has the *topological Paley-Wiener property* if for every non-zero $f \in L_c^\infty(G)$, the map $\pi \rightarrow \pi(f)$ cannot vanish on any non-empty open subset of the reduced dual \widehat{G}_r of G . Our main result is an extension theorem which asserts that if G contains a closed normal subgroup N that has the topological Paley-Wiener property, then so does G , provided that G/N is abelian and compact-free (Theorem 2.2). As a consequence, compact-free nilpotent locally compact groups and simply connected solvable Lie groups share this topological Paley-Wiener property (Corollaries 2.4 and 2.5).

1. PRELIMINARIES AND SOME BASIC RESULTS

Let G be a locally compact group with fixed left Haar measure, and let $C^*(G)$ be the completion of the convolution algebra $L^1(G)$ with respect to the norm $\|f\|_{C^*(G)} = \sup\{\|\pi(f)\|\}$, where the supremum is taken over all $*$ -representations π of $L^1(G)$ as an algebra of bounded linear operators in a Hilbert space. Let $P(G)$ be the set of all continuous positive definite functions on G , and let $B(G)$ denote the

Received by the editors December 2, 2003.

2000 *Mathematics Subject Classification*. Primary 22D10, 22E25, 43A30, 43A40.

The second author was supported by NSERC grant A7679.

linear span of $P(G)$. Then $B(G)$ can be identified with the dual space $C^*(G)^*$ of $C^*(G)$ by the pairing $\langle f, u \rangle = \int_G f(x)u(x)dx$ for $f \in L^1(G)$ and $u \in B(G)$. With pointwise multiplication and the dual norm, $B(G)$ is a commutative Banach algebra, called the *Fourier-Stieltjes algebra* of G . The *Fourier algebra* $A(G)$ is the norm closure in $B(G)$ of $A_c(G) = B(G) \cap C_c(G)$, the compactly supported functions in $B(G)$. For details regarding $B(G)$ and $A(G)$, see the fundamental paper [5].

As is customary, we shall use the same letter, for example π , for a unitary representation of G and the corresponding $*$ -representations of $L^1(G)$ and $C^*(G)$. For any representation π of G , let $A_\pi(G)$ denote the closed linear subspace of $B(G)$ generated by all coefficient functions of π , and $B_\pi(G)$ the w^* -closure of $A_\pi(G)$ in $B(G)$. Then $B_\pi(G)$ can be identified with the dual of the C^* -algebra $\pi(C^*(G))$. Note that, when $\rho = \rho_G$ denotes the regular representation of G on $L^2(G)$, then $A_\rho(G) = A(G)$, and $B_\rho(G)$ equals $B(G)$ if and only if G is amenable.

Moreover, we need to recall some definitions from representation theory. If S and T are sets of unitary representations, then S is weakly contained in T ($S \prec T$) if $\bigcap\{\ker \sigma : \sigma \in S\} \supseteq \bigcap\{\ker \tau : \tau \in T\}$, where $\ker \pi$ denotes the C^* -kernel of a representation π . The sets S and T are said to be weakly equivalent ($S \sim T$) if $S \prec T$ and $T \prec S$. Then, for any two representations σ and τ , $\sigma \prec \tau$ if and only if $B_\sigma(G) \subseteq B_\tau(G)$. The support of a representation π , $\text{supp } \pi$, is the closed subset of \widehat{G} consisting of all $\tau \in \widehat{G}$ such that $\tau \prec \pi$. Thus $\pi \sim \widehat{G}$ if and only if $\text{supp } \pi = \widehat{G}$. As general references to dual spaces of locally compact groups and weak containment properties, we mention [4] and [6].

Lemma 1.1. *If G contains a non-trivial compact normal subgroup K , then G cannot have the topological Paley-Wiener property.*

Proof. Let μ_K denote the (normalized) Haar measure of K . Let π be an irreducible representation of G such that $\pi \notin \widehat{G/K}$. If σ is an irreducible subrepresentation of $\pi|_K$, then, since $\sigma \neq 1_K$, by the orthogonality relations $\int_K \langle \sigma(k)\xi, \eta \rangle dk = 0$ for all $\xi, \eta \in \mathcal{H}_\sigma$. It follows that $\pi(\mu_K) = 0$. Now, consider $f = g * \mu_K, g \in C_c(G)$. Then $f \in L_c^\infty(G)$ and $\pi(f) = 0$ for all $\pi \in \widehat{G_r} \setminus \widehat{G/K}$, a non-empty open subset of $\widehat{G_r}$. \square

For any $f \in L^1(G)$, let

$$Z_f = \{\pi \in \widehat{G_r} : \pi(f) = 0\},$$

the zero set of f in $\widehat{G_r}$. Note that when studying the topological Paley-Wiener property, we can always assume that the function $f \in L_c^\infty(G)$ actually belongs to $A_c(G)$ since $f * f^*$ is positive definite and $Z_{f*f^*} = Z_f$. We say that $f \in A_c(G)$ generates $B_\rho(G)$ in the w^* -topology if the linear span of all two-sided translates of f is w^* -dense in $B_\rho(G)$.

Lemma 1.2. *Let G be a locally compact group and let $f \in A_c(G)$. Then Z_f has empty interior if and only if f generates $B_\rho(G)$ in the w^* -topology.*

Proof. Let I_f denote the closed ideal of $C_r^*(G)$ generated by $\rho(f)$. If $\overset{\circ}{Z}_f \neq \emptyset$, then the non-zero ideal J of $C_r^*(G)$ defined by $\widehat{J} = \overset{\circ}{Z}_f$ satisfies $J \cap I_f = \{0\}$ since $\widehat{I}_f = \widehat{G_r} \setminus Z_f$. Conversely, if there exists a non-zero ideal J of $C_r^*(G)$ such that $J \cap I_f = \{0\}$, then $\widehat{J} \cap (\widehat{G_r} \setminus Z_f) = \widehat{J} \cap \widehat{I}_f = (J \cap I_f)^\wedge = \emptyset$ and hence $\emptyset \neq \widehat{J} \subseteq Z_f$, so

$\overset{\circ}{Z}_f \neq \emptyset$. Therefore, to prove the lemma it suffices to show that there exists a non-zero ideal J of $C_r^*(G)$ with $J \cap I_f = \{0\}$ if and only if E_f , the w^* -closed subspace of $B_\rho(G)$ generated by all two-sided translates of f , does not coincide with $B_\rho(G)$.

Let Δ denote the modular function of G and for any function h on G , define \tilde{h} by $\tilde{h}(y) = \Delta(y^{-1})h(y^{-1})$. Then note that $\langle \rho(h)\rho(g), u \rangle = \langle \rho(g), \tilde{h} * u \rangle$ for all $g \in C_c(G)$ and $h, u \in A_c(G)$, and hence $\langle \rho(h)T, u \rangle = \langle T, \tilde{h} * u \rangle$ for all $T \in C_r^*(G)$ and $h, u \in A_c(G)$.

Now let J be an ideal as above. Then $\langle T, \tilde{f} * u \rangle = 0$ for all $T \in J$ and $u \in A_c(G)$ since $\rho(f)T = 0$. Writing f as $f(\cdot) = \langle \pi(\cdot)\xi, \eta \rangle$ for some representation π of G and $\xi, \eta \in \mathcal{H}_\pi$ and choosing u such that $u \geq 0$ and $\|u\|_1 = 1$, we have

$$\begin{aligned} (f * u)(x) - f(x) &= \int_G u(y^{-1})\langle \pi(x)\pi(y)\xi, \eta \rangle dy - \langle \pi(x)\xi, \eta \rangle \\ &= \left\langle \pi(x) \left(\int_G u(y^{-1})(\pi(y)\xi - \xi) dy \right), \eta \right\rangle. \end{aligned}$$

This implies that

$$\|f * u - f\|_{B_\rho(G)} \leq \|\eta\| \sup\{\|\pi(y)\xi - \xi\| : y^{-1} \in \text{supp } u\}.$$

Since $\langle T, f * u \rangle = 0$, taking for $u \in A_c(G)$ the usual approximate identity for $L^1(G)$, it follows that

$$|\langle T, f \rangle| \leq \|T\| \cdot \|f * u - f\|_{B_\rho(G)} \rightarrow 0$$

for each $T \in J$. Since J^\perp is a w^* -closed two-sided translation invariant subspace of $B_\rho(G)$, we conclude that $E_f \subseteq J^\perp$, a proper subspace of $B_\rho(G)$.

Conversely, suppose that $E_f \neq B_\rho(G)$ and let

$$J = \{T \in C_r^*(G) : \langle T, u \rangle = 0 \text{ for all } u \in E_f\}.$$

Then J is a non-zero closed ideal $C_r^*(G)$. We show that $J \cap I_f = \{0\}$. For $T \in C_r^*(G)$, $s, t \in G$ and $u \in A_c(G)$, we have

$$\langle \rho(L_{t^{-1}}R_{s^{-1}}\tilde{f})T, u \rangle = \Delta(ts)\langle T, (L_sR_t f) * u \rangle.$$

Therefore, it will follow that $J \cap I_f = \{0\}$ once we have verified that if E is a w^* -closed translation invariant subspace of $B_\rho(G)$, then $E * u \subseteq E$ for every $u \in A_c(G)$. This can be seen as follows. If $v(\cdot) = \langle \pi(\cdot)\xi, \eta \rangle \in E$, then

$$(v * u)(\cdot) = \left\langle \pi(\cdot) \left(\int_G u(y^{-1})\pi(y)\xi dy \right), \eta \right\rangle.$$

Given $\epsilon > 0$, there exist $y_1, \dots, y_n \in G$ and $c_1, \dots, c_n \geq 0$ such that

$$\left\| \int_G u(y^{-1})\pi(y)\xi dy - \sum_{j=1}^n c_j \pi(y_j)\xi \right\| \leq \epsilon.$$

Then $w(\cdot) = \sum_{j=1}^n c_j \langle \pi(\cdot)\pi(y_j)\xi, \eta \rangle \in E$ and

$$\|v * u - w\|_{B_\rho(G)} \leq \|\eta\| \cdot \left\| \int_G u(y^{-1})\pi(y)\xi dy - \sum_{j=1}^n c_j \pi(y_j)\xi \right\| \leq \epsilon \|\eta\|.$$

This finishes the proof. □

It is worth pointing out that, according to the following lemma, the topological Paley-Wiener property is equivalent to a dichotomy for the subspaces $\overline{B_\pi(G) \cap A_c(G)}$ of $A(G)$, which has been investigated in Section 5 of [13].

Lemma 1.3. *For any locally compact group G , the following two conditions are equivalent:*

- (i) *G has the topological Paley-Wiener property.*
- (ii) *For any unitary representation π of G , either $B_\pi(G) \cap A_c(G) = \{0\}$ or $\overline{B_\pi(G) \cap A_c(G)} = A(G)$.*

Proof. (i) \Rightarrow (ii) Let π be any representation of G , and suppose there exists $f \in B_\pi(G) \cap A_c(G)$, $f \neq 0$. By (i), Z_f has empty interior and hence f generates $B_\rho(G)$ in the w^* -topology (Lemma 1.2). Thus

$$B_\rho(G) \subseteq \overline{B_\pi(G) \cap A_c(G)}^{w^*} \subseteq B_\pi(G),$$

and this in turn implies that

$$A(G) = \overline{B_\rho(G) \cap A_c(G)} \subseteq \overline{B_\pi(G) \cap A_c(G)},$$

as required.

(ii) \Rightarrow (i) Let f be a non-zero function in $A_c(G)$, and let E denote the closed two-sided invariant subspace of $A(G)$ generated by f . There exists a representation π of G such that $E = A_\pi(G)$ [2, Théorème 3.17]. By hypothesis, either $B_\pi(G) \cap A_c(G) = \{0\}$ or $\overline{B_\pi(G) \cap A_c(G)} = A(G)$. However, $f \in B_\pi(G) \cap A_c(G)$. Therefore $\overline{B_\pi(G) \cap A_c(G)} = A(G)$ and hence

$$\overline{E}^{w^*} \subseteq B_\rho(G) = \overline{A(G)}^{w^*} = \overline{B_\pi(G) \cap A_c(G)}^{w^*} \subseteq \overline{E}^{w^*},$$

so that f generates $B_\rho(G)$ in the w^* -topology. By Lemma 1.2, $\overset{\circ}{Z}_f \neq \emptyset$. \square

An interesting consequence of Lemma 1.3 is that the topological Paley-Wiener property can be interpreted as an approximation property for $A(G)$.

A locally compact group G is called a SIN-group (a group with small invariant neighbourhoods) if there exists a neighbourhood basis of the identity consisting of sets V such that $x^{-1}Vx = V$ for all $x \in G$. SIN-groups provide a large class of locally compact groups comprising all abelian groups, compact groups and discrete groups. For the structure theory of SIN-groups, see [11].

Theorem 1.4. *For a SIN-group G , the following two conditions are equivalent:*

- (i) *G contains no non-trivial compact normal subgroup.*
- (ii) *G has the topological Paley-Wiener property.*

Proof. This is an immediate consequence of Lemma 1.3 and Theorem 5.6 of [13]. \square

We conclude this section with the observation that the topological Paley-Wiener property is in general not inherited by normal subgroups. Indeed, an easy example showing this can be constructed as follows. Let F be a finite abelian group and let $N = \sum_{j=1}^{\infty} F_j$ be the direct sum of copies F_j of F . Let $G = S_\infty \ltimes N$ be the semidirect product, where the infinite symmetric group S_∞ acts on N by permuting the index set \mathbb{N} . Obviously, G is a discrete ICC group and hence has the weak Paley-Wiener property (Theorem 1.4), whereas the normal subgroup N is an abelian torsion-group.

2. EXTENSIONS AND APPLICATIONS

In this section we are going to prove two theorems of the nature that if a locally compact group G contains a closed normal subgroup that has the topological Paley-Wiener property, then under certain additional hypotheses, G also has the topological Paley-Wiener property. Both of these results apply to various (classes of) locally compact groups. A similar, but much less applicable, result has been shown in [13, Proposition 5.3].

Let H be a closed subgroup of G and let τ be a representation of H . The representation of G induced by τ is denoted $\text{ind}_H^G \tau$. In the sequel we shall use that $\pi \otimes \text{ind}_H^G \tau = \text{ind}_H^G(\pi|_H \otimes \tau)$ and that $\pi \prec \text{ind}_H^G(\pi|_H)$ when G is amenable [10, Theorem 5.1].

Lemma 2.1. *Let H be a closed subgroup of the locally compact group G . Let U be a non-empty open subset of \widehat{G}_r , and let*

$$V = \{\tau \in \widehat{H}_r : \text{supp}(\text{ind}_H^G \tau) \not\subseteq \widehat{G}_r \setminus U\}.$$

Then V is non-empty and open in \widehat{H}_r .

Proof. To show that $\widehat{H}_r \setminus V$ is closed in \widehat{H}_r , let $(\tau_\alpha)_\alpha$ be a net in $\widehat{H}_r \setminus V$ such that $\tau_\alpha \rightarrow \tau$ for some $\tau \in \widehat{H}_r$. Since inducing is continuous in Fell's subgroup representation topology [6, Theorem 4.2], $\text{ind}_H^G \tau_\alpha \rightarrow \text{ind}_H^G \tau$. Since $\text{ind}_H^G \tau_\alpha \prec \widehat{G}_r \setminus U$ for all α and $\widehat{G}_r \setminus U$ is closed in \widehat{G}_r , it follows that $\text{supp}(\text{ind}_H^G \tau) \subseteq \widehat{G}_r \setminus U$ and hence $\tau \in \widehat{H}_r \setminus V$.

Assume that $V = \emptyset$, that is, $\text{supp}(\text{ind}_H^G \tau) \subseteq \widehat{G}_r \setminus U$ for all $\tau \in \widehat{H}_r$. Then, since $\rho_G = \text{ind}_H^G \rho_H$, we obtain that

$$\widehat{G}_r \sim \{\text{ind}_H^G \tau : \tau \in \widehat{H}_r\} \sim \bigcup_{\tau \in \widehat{H}_r} \text{supp}(\text{ind}_H^G \tau) \subseteq \widehat{G}_r \setminus U.$$

This contradiction shows that V is non-empty. □

Recall that a locally compact group G is said to be *compact-free* if the identity is the only element of G generating a relatively compact subgroup.

Theorem 2.2. *Let G be a locally compact group containing a closed normal subgroup N such that G/N is abelian and compact-free. If N has the topological Paley-Wiener property, then so does G .*

Proof. Let $f \in C_c(G)$ be such that $\overset{\circ}{Z}_f \neq \emptyset$. We have to show that $L_x f|_N = 0$ for all $x \in G$. Temporarily, fix $\pi \in \overset{\circ}{Z}_f$ and $\xi, \eta \in \mathcal{H}_\pi$, and consider on $\widehat{G/N}$ the function

$$\chi \rightarrow \langle (\pi \otimes \chi)(f)(\xi \otimes 1), \eta \otimes 1 \rangle.$$

Define $h \in C_c(G/N)$ by

$$h(xN) = \int_N f(xn) \langle \pi(xn)\xi, \eta \rangle dn,$$

$x \in G$. Then, for $\chi \in \widehat{G/N}$,

$$\begin{aligned} \widehat{h}(\chi) &= \int_{G/N} \chi(xN) \left(\int_N f(xn) \langle \pi(xn)\xi, \eta \rangle \right) d(xN) \\ &= \langle (\pi \otimes \chi)(f)(\xi \otimes 1), \eta \otimes 1 \rangle. \end{aligned}$$

Since G/N is abelian and compact-free, \widehat{h} can vanish only on a set of Haar measure zero in $\widehat{G/N}$ [13, Lemma 5.1]. However, \widehat{h} vanishes on the set

$$X = \{\chi \in \widehat{G/N} : \pi \otimes \chi \in \overset{\circ}{Z}_f\},$$

and X is a non-empty open subset of $\widehat{G/N}$ since $\pi \in X$ and the mapping $\chi \rightarrow \pi \otimes \chi$ from $\widehat{G/N}$ into \widehat{G} is continuous. We conclude that $h = 0$. Since $\xi, \eta \in \mathcal{H}_\pi$ are arbitrary, it follows that $(\pi \otimes \chi)(f) = 0$ for all $\pi \in \overset{\circ}{Z}_f$ and $\chi \in \widehat{G/N}$. Now,

$$\pi \otimes \widehat{G/N} = \{\pi \otimes \chi : \chi \in \widehat{G/N}\} \sim \pi \otimes \text{ind}_N^G 1_N = \text{ind}_N^G(\pi|_N),$$

and since $\pi \prec \rho_G$ implies that $\pi|_N \prec \rho_N$, we get that

$$\text{ind}_N^G(\pi|_N) \prec \text{ind}_N^G \rho_N = \rho_G$$

and therefore $\pi \otimes \widehat{G/N} \subseteq \widehat{G}_r$. Thus we have seen that $\overset{\circ}{Z}_f \otimes \widehat{G/N} \subseteq Z_f$. Since $\sigma \rightarrow \sigma \otimes \chi$ is a homeomorphism of \widehat{G}_r ,

$$\overset{\circ}{Z}_f \otimes \widehat{G/N} = \bigcup_{\chi \in \widehat{G/N}} \overset{\circ}{Z}_f \otimes \{\chi\}$$

is open in \widehat{G}_r , whence $\overset{\circ}{Z}_f \otimes \widehat{G/N} = \overset{\circ}{Z}_f$.

Now, let $x \in G$ and define $g \in C_c(N)$ by $g(n) = f(xn)$. We have to show that $\overset{\circ}{Z}_g \neq \emptyset$ for all such functions g . To that end, define a subset V of \widehat{N}_r by

$$V = \{\tau \in \widehat{N}_r : \text{supp}(\text{ind}_N^G \tau) \not\subseteq \widehat{G}_r \setminus \overset{\circ}{Z}_f\}.$$

By Lemma 2.1, V is non-empty and open in \widehat{N}_r . It remains to verify that $V \subseteq Z_g$. Fix $\tau \in V$ and choose a net $(\pi_\alpha)_\alpha$ in \widehat{G}_r such that $\pi_\alpha|_N \rightarrow \tau$ in Fell's topology. We claim that $\pi_\alpha \in \overset{\circ}{Z}_f$ eventually. Towards a contradiction, after passing to a subnet if necessary, let us assume that $\pi_\alpha \notin \overset{\circ}{Z}_f$ for all α . Since $\overset{\circ}{Z}_f = \overset{\circ}{Z}_f \otimes \widehat{G/N}$, it follows that

$$\text{ind}(\pi_\alpha|_N) \sim \pi_\alpha \otimes \widehat{G/N} \subseteq \widehat{G}_r \setminus \overset{\circ}{Z}_f$$

for all α . Therefore, for each $\sigma \in \text{supp}(\pi_\alpha|_N)$,

$$\text{supp}(\text{ind}_N^G \sigma) \subseteq \text{supp}(\text{ind}_N^G(\pi_\alpha|_N)) \subseteq \widehat{G}_r \setminus \overset{\circ}{Z}_f,$$

and hence $\sigma \in \widehat{N}_r \setminus V$. Thus $\pi_\alpha|_N \prec \widehat{N}_r \setminus V$ and so $\tau \in \widehat{N}_r \setminus V$. This contradiction shows that $\pi_\alpha \in \overset{\circ}{Z}_f$ eventually. Since

$$\text{ind}_N^G(\pi_\alpha|_N) \sim \pi_\alpha \otimes \widehat{G/N} \subseteq \overset{\circ}{Z}_f \otimes \widehat{G/N} \subseteq Z_f,$$

and $\text{ind}(\pi_\alpha|_N) \rightarrow \text{ind}_N^G \tau$, it follows that $\text{supp}(\text{ind}_N^G \tau) \subseteq Z_f$. But $\text{ind}_N^G \tau(f) = 0$ implies that $\tau(L_x f|_N) = 0$ (compare [16, Lemma 2.1]), that is, $\tau \in Z_g$. This finishes the proof. \square

Corollary 2.3. *Suppose that G possesses a sequence $H_0 = \{e\} \subseteq H_1 \subseteq \dots \subseteq H_m = G$ of closed subgroups such that H_{j-1} is normal in H_j and H_j/H_{j-1} is compact-free abelian ($1 \leq j \leq m$). Then G has the topological Paley-Wiener property.*

Proof. Using Theorem 2.2, the statement follows immediately by induction on m . \square

Corollary 2.4. *Let G be a nilpotent locally compact group. Then G has the topological Paley-Wiener property if and only if G is compact-free.*

Proof. Suppose that G has the topological Paley-Wiener property. Then, by Lemma 1.1, G cannot have a non-trivial compact normal subgroup. It is easy to see that this implies that $G^c = \{e\}$. Indeed, let $\{e\} = Z_0(G) \subseteq Z_1(G) \subseteq \dots$ denote the ascending central series of G and, assuming that $G^c \neq \{e\}$, let $m \geq 0$ be maximal with the property that $G^c \cap Z_m(G) = \{e\}$. Then pick $a \in G^c \cap Z_{m+1}(G), a \neq e$, and let H be the closed subgroup of G generated by $Z_m(G)$ and a . Then H is normal in G and hence H^c is a closed normal subgroup of G . However, H^c is the closed subgroup generated by a , which is compact.

Conversely, if G is compact-free, then G_0 , the connected component of the identity, is simply connected and $D = G/G_0$ is discrete and torsion-free [9, Theorem 8.3]. We now apply Corollary 2.3 as follows. If $G = G_0$, then $Z_m(G)$ and the quotient group $G/Z_m(G)$ are simply connected for all m . Then induction yields that G has the topological Paley-Wiener property. In the general case, note that $D/Z_m(D)$ is torsion-free for all m [3, Corollary 2.11]. Thus, a further induction on the length of nilpotency of D shows that G has the topological Paley-Wiener property. □

Corollary 2.5. *Let G be a simply connected solvable Lie group. Then G has the topological Paley-Wiener property.*

Proof. By the structure theory of simply connected solvable Lie groups, there exists a sequence $H_0 = \{e\} \subseteq H_1 \subseteq \dots \subseteq H_m = G$ of closed subgroups such that H_{j-1} is normal in H_j and $H_j/H_{j-1} = \mathbb{R}$ (see [12, Satz III.3.30]). Thus G has the topological Paley-Wiener property by Corollary 2.3. □

Theorem 2.6. *Let G be a locally compact group and let N be a closed normal subgroup of G such that G/N is amenable. Suppose there exists a dense subset T of \widehat{N}_r such that for each $\tau \in T$, $\text{ind}_N^G \tau$ is weakly equivalent to some irreducible representation. If N has the topological Paley-Wiener property, then so does G .*

Proof. Let $f \in C_c(G)$ such that $\overset{\circ}{Z}_f \neq \emptyset$. By Lemma 2.1, the set

$$V = \{\tau \in \widehat{N}_r : \text{supp}(\text{ind}_N^G \tau) \not\subseteq \widehat{G}_r \setminus \overset{\circ}{Z}_f\}$$

is non-empty and open in \widehat{N}_r . Since T is dense in \widehat{N}_r , $T \cap V$ is dense in V . By hypothesis, for every $\tau \in T$ there exists $\pi_\tau \in \widehat{G}$ such that $\pi_\tau \sim \text{ind}_N^G \tau$. Since G/N is amenable, $\pi_\tau \in \widehat{G}_r$. Then $\pi_\tau \in \overset{\circ}{Z}_f$ for each $\tau \in T \cap V$. Indeed, for such τ ,

$$\pi_\tau \sim \text{supp}(\text{ind}_N^G \tau) \not\subseteq \widehat{G}_r \setminus \overset{\circ}{Z}_f$$

and hence $\pi_\tau \in \overset{\circ}{Z}_f$. Now, for any $\tau \in V$ there exists a net $(\tau_\alpha)_\alpha$ in $T \cap V$ converging to τ . Then

$$\pi_{\tau_\alpha} \sim \text{ind}_N^G \tau_\alpha \rightarrow \text{ind}_N^G \tau,$$

and since $\pi_{\tau_\alpha}(f) = 0$ for all α , it follows that $\text{ind}_N^G \tau(f) = 0$. This in turn implies that $\tau(L_x f|_N) = 0$ for all $x \in G$. Since N has the topological Paley-Wiener property, we conclude that $f = 0$. □

We finish the paper with an example to which Theorem 2.6 applies, but neither Theorem 2.2 nor Proposition 5.3 of [13] applies.

Example 2.7. Let $\mathbb{Z}_2 = \{1, -1\}$, $K = \prod_{n=1}^{\infty} \mathbb{Z}_2$ and $N = \sum_{n=1}^{\infty} \mathbb{Z}$, the restricted direct sum of copies of \mathbb{Z} . Let G be the semidirect product $G = K \ltimes N$, where K acts on N by $(\epsilon \cdot x)_n = \epsilon_n x_n$ ($\epsilon = (\epsilon_n)_n \in K, x = (x_n)_n \in N$). Identifying $\widehat{\mathbb{Z}}$ with \mathbb{T} , we have $\widehat{N} = \prod_{n=1}^{\infty} \mathbb{T}$ and, for $z = (z_n)_n \in \widehat{N}$ and $\epsilon = (\epsilon_n)_n \in K$, $\epsilon \cdot z = z$ if and only if $z_n = \overline{z_n}$ for all $n \in \mathbb{N}$ such that $\epsilon_n = -1$. Now, $T = \bigcap_{n=1}^{\infty} \{z \in \widehat{N} : \overline{z_n} \neq z_n\}$ is dense in \widehat{N} , and each $z \in T$ has a trivial stability group in K and hence the associated induced representation of G is irreducible. By Theorem 2.6, G has the topological Paley-Wiener property.

In [13, Example 5.4] we have given several other examples of locally compact groups having the topological Paley-Wiener property.

REFERENCES

- [1] D. Arnal and J. Ludwig, *Q.U.P. and and Paley-Wiener properties of unimodular, especially nilpotent, Lie groups*, Proc. Amer. Math. Soc. **125** (1997), 1071-1080. MR1353372 (97f:43004)
- [2] G. Arsac, *Sur l'espace de Banach engendré par les coefficients d'une représentation unitaire*, Publ. Dép. Math. (Lyon) **13** (1976), 1-101. MR0444833 (56:3180)
- [3] G. Baumslag, *Lectures on nilpotent groups*, Amer. Math. Soc. Regional Conference Series No. 2 (1971). MR0283082 (44:315)
- [4] J. Dixmier, *C*-algebras*, North-Holland, 1977. MR0458185 (56:16388)
- [5] P. Eymard, *L'algèbre de Fourier d'un groupe localement compact*, Bull. Soc. Math. France **92** (1964), 181-236. MR0228628 (37:4208)
- [6] J.M.G. Fell, *Weak containment and induced representations. II*, Trans. Amer. Math. Soc. **110** (1964), 424-447. MR0159898 (28:3114)
- [7] G. Garimella, *Un théorème de Paley-Wiener pour les groupes de Lie nilpotents*, J. Lie Theory **5** (1995), 165-172. MR1389426 (97f:22016)
- [8] G. Garimella, *Weak Paley-Wiener property for completely solvable Lie groups*, Pacific J. Math. **187** (1999), 51-63. MR1674293 (2000a:22011)
- [9] V.M. Gluskov, *Locally nilpotent locally bicomact groups*, Trudy Moscov. Obshch. **4** (1955), 291-332. (Russian) MR0072422 (17:281b)
- [10] F.F. Greenleaf, *Amenable actions of locally compact groups*, J. Funct. Anal. **4** (1969), 295-315. MR0246999 (40:268)
- [11] S. Grosser and M. Moskowitz, *Compactness conditions in topological groups*, J. reine angew. Math. **246** (1971), 1-40. MR0284541 (44:1766)
- [12] J. Hilgert and K.-H. Neeb, *Lie-Gruppen und Lie-Algebren*, Vieweg, 1991.
- [13] E. Kaniuth, A.T. Lau and G. Schlichting, *Lebesgue type decomposition of subspaces of Fourier-Stieltjes algebras*, Trans. Amer. Math. Soc. **355** (2003), 1467-1490. MR1946400 (2004c:43004)
- [14] V. Kisil, *A theorem of Paley-Wiener type for nilpotent Lie groups*, Ukrainian Math. J. **50** (1998), 1786-1788. MR1706645 (2000m:22010)
- [15] R.L. Lipsman and J. Rosenberg, *The behavior of Fourier transforms for nilpotent Lie groups*, Trans. Amer. Math. Soc. **348** (1996), 1031-1050. MR1370646 (97d:22008)
- [16] J. Ludwig, *Good ideals in the group algebra of a nilpotent Lie group*, Math. Z. **161** (1983), 195-210. MR0498970 (58:16958)

INSTITUT FÜR MATHEMATIK, UNIVERSITÄT PADERBORN, D-33095 PADERBORN, GERMANY
E-mail address: kaniuth@math.uni-paderborn.de

DEPARTMENT OF MATHEMATICAL AND STATISTICAL SCIENCES, UNIVERSITY OF ALBERTA, EDMONTON, ALBERTA, CANADA T6G 2G1
E-mail address: tlau@math.ualberta.ca

FAKULTÄT FÜR MATHEMATIK, TECHNISCHE UNIVERSITÄT MÜNCHEN, D-80290 MÜNCHEN, GERMANY
E-mail address: schlicht@ma.tum.de