

HOPF ALGEBRAS OF DIMENSION $2p$

SIU-HUNG NG

(Communicated by Martin Lorenz)

ABSTRACT. Let H be a finite-dimensional Hopf algebra over an algebraically closed field of characteristic 0. If H is not semisimple and $\dim(H) = 2n$ for some odd integer n , then H or H^* is not unimodular. Using this result, we prove that if $\dim(H) = 2p$ for some odd prime p , then H is semisimple. This completes the classification of Hopf algebras of dimension $2p$.

0. INTRODUCTION

In recent years, there has been some progress on the classification problems of finite-dimensional Hopf algebras over an algebraically closed field k of characteristic 0 (cf. [Mon98], [And02]). It is shown in [Zhu94] that Hopf algebras of dimension p , where p is a prime, are isomorphic to the group algebra $k[\mathbb{Z}_p]$. In [Ng02b] and [Ng02a], the author completed the classification of Hopf algebras of dimension p^2 , which started in [AS98] and [Mas96]. They are group algebras and Taft algebras of dimension p^2 (cf. [Taf71]). However, the classification of Hopf algebras of dimension pq , where p, q are distinct prime numbers, remains open in general.

It is shown in [EG98], [GW00] that semisimple Hopf algebras over k of dimension pq are trivial (i.e. isomorphic to either group algebras or the dual of group algebras). Most recently, Etingof and Gelaki proved that if p, q are odd primes such that $p < q \leq 2p + 1$, then any Hopf algebra over k of dimension pq is semisimple [EG]. Meanwhile, the author proved the same result, using a different method, for the case that p, q are twin primes [Ng]. In addition to that, Williams settled the case of dimensions 6 and 10 in [Wil88], and Beattie and Dăscălescu did dimensions 14, 65 in [BD]. Hopf algebras of dimensions 6, 10, 14 and 65 are semisimple, and so they are trivial.

In this paper, we prove that any Hopf algebra of dimension $2p$, where p is an odd prime, over an algebraically closed field k of characteristic 0, is semisimple. By [Mas95], semisimple Hopf algebras of dimension $2p$ are isomorphic to

$$k[\mathbb{Z}_{2p}], \quad k[D_{2p}] \quad \text{or} \quad k[D_{2p}]^*$$

where D_{2p} is the dihedral group of order $2p$. Hence, our main result, Theorem 3.3, completes the classification of Hopf algebras of dimension $2p$.

Received by the editors November 24, 2003 and, in revised form, April 7, 2004.
2000 *Mathematics Subject Classification*. Primary 16W30.
Key words and phrases. Finite-dimensional Hopf algebras.

1. NOTATION AND PRELIMINARIES

Throughout this paper, p is an odd prime, k denotes an algebraically closed field of characteristic 0, and H denotes a finite-dimensional Hopf algebra over k with antipode S . Its comultiplication and counit are respectively denoted by Δ and ϵ . We will use Sweedler's notation [Swe69]:

$$\Delta(x) = \sum x_{(1)} \otimes x_{(2)}.$$

A non-zero element $a \in H$ is called group-like if $\Delta(a) = a \otimes a$. The set of all group-like elements $G(H)$ of H is a linearly independent set, and it forms a group under the multiplication of H . For the details of elementary aspects for finite-dimensional Hopf algebras, readers are referred to the references [Swe69] and [Mon93].

Let $\lambda \in H^*$ be a non-zero right integral of H^* and $\Lambda \in H$ a non-zero left integral of H . There exists $\alpha \in \text{Alg}(H, k) = G(H^*)$, independent of the choice of Λ , such that $\Lambda a = \alpha(a)\Lambda$ for $a \in H$. Likewise, there is a group-like element $g \in H$, independent of the choice of λ , such that $\beta\lambda = \beta(g)\lambda$ for $\beta \in H^*$. We call g the distinguished group-like element of H and α the distinguished group-like element of H^* . Then we have Radford's formula [Rad76] for S^4 :

$$(1.1) \quad S^4(a) = g(\alpha \rightharpoonup a \leftarrow \alpha^{-1})g^{-1} \quad \text{for } a \in H,$$

where \rightharpoonup and \leftarrow denote the natural actions of the Hopf algebra H^* on H described by

$$\beta \rightharpoonup a = \sum a_{(1)}\beta(a_{(2)}) \quad \text{and} \quad a \leftarrow \beta = \sum \beta(a_{(1)})a_{(2)}$$

for $\beta \in H^*$ and $a \in H$. In particular, we have the following proposition.

Proposition 1.1 ([Rad76]). *Let H be a finite-dimensional Hopf algebra with antipode S over the field k . Suppose that g and α are distinguished group-like elements of H and H^* respectively. Then the order of S^4 divides the least common multiple of the order of g and the order of α .* \square

For any $a \in H$, the linear operator $r(a) \in \text{End}_k(H)$ is defined by $r(a)(b) = ba$ for $b \in H$. The semisimplicity of a finite-dimensional Hopf algebra can be characterized by the antipode.

Theorem 1.2 ([LR87], [LR88], [Rad94]). *Let H be a finite-dimensional Hopf algebra with antipode S over the field k . Then the following statements are equivalent:*

- (1) H is not semisimple;
- (2) H^* is not semisimple;
- (3) $S^2 \neq id_H$;
- (4) $\text{Tr}(S^2) = 0$;
- (5) $\text{Tr}(S^2 \circ r(a)) = 0$ for all $a \in H$. \square

Proposition 1.3 ([Ng, Corollary 2.2]). *Let H be a finite-dimensional Hopf algebra over k with antipode S and g the distinguished group-like element of H . If $\text{lcm}(o(S^4), o(g)) = n$ is an odd integer greater than 1, then the subspace*

$$H_- = \{u \in H \mid S^{2n}(u) = -u\}$$

has even dimension. \square

The following lemma is useful in our remaining discussion.

Lemma 1.4. *Let V be a finite-dimensional vector space over the field k . If T is a linear automorphism on V such that $\text{Tr}(T) = 0$ and $o(T) = q^n$ for some prime q and positive integer n , then*

$$q \mid \dim(V).$$

Proof. Let $\omega \in k$ be a primitive q^n th root of unity and

$$V_i = \{u \in V \mid T(u) = \omega^i u\} \quad \text{for } i = 0, \dots, q^n - 1.$$

Consider the integral polynomial

$$f(x) = \sum_{i=0}^{q^n-1} \dim(V_i)x^i.$$

Since

$$0 = \text{Tr}(T) = f(\omega),$$

there exists $g(x) \in \mathbb{Z}[x]$ such that

$$f(x) = \Phi_{q^n}(x)g(x)$$

where $\Phi_{q^n}(x)$ is the q^n th cyclotomic polynomial. Hence,

$$\dim(V) = f(1) = \Phi_{q^n}(1)g(1).$$

Since $\Phi_{q^n}(x) = \Phi_q(x^{q^{n-1}})$, $\Phi_{q^n}(1) = \Phi_q(1) = q$. Thus we have

$$q \mid \dim(V). \quad \square$$

2. UNIMODULARITY OF HOPF ALGEBRAS OF DIMENSION $2n$

In this section, we prove that if H is a non-semisimple Hopf algebra over k of dimension $2n$, where n is an odd integer, then H or H^* is not unimodular. This result is essential to the proof of our main result in the next section.

Proposition 2.1. *Let H be a finite-dimensional Hopf algebra over the field k with antipode S . If H is unimodular and $o(S^2) = 2$, then*

$$4 \mid \dim(H).$$

Proof. Let λ be a non-zero right integral of H^* . Since H is unimodular, by [Rad94, Theorem 3],

$$\lambda(ab) = \lambda(S^2(b)a)$$

for all $a, b \in H$. Let

$$H_i = \{u \in H \mid S^2(u) = (-1)^i u\} \quad \text{for } i = 0, 1.$$

We claim that $(a, b) = \lambda(ab)$ defines a non-degenerate alternating form on H_1 . For any $a, b \in H_1$,

$$\lambda(ab) = \lambda(S^2(b)a) = -\lambda(ba).$$

Since $\lambda(u) = \lambda(S^2(u)) = -\lambda(u)$ for all $u \in H_1$, $\lambda(H_1) = \{0\}$. Let $a \in H_1$ such that $\lambda(ab) = 0$ for all $b \in H_1$. Then for all $b \in H_0$, $ab \in H_1$ and so $\lambda(ab) = 0$. By the non-degeneracy of λ on H , $a = 0$. Therefore, $(a, b) = \lambda(ab)$ defines a non-degenerate alternating bilinear form on H_1 and hence $\dim(H_1)$ is even. Since $o(S^2) = 2$, by Theorem 1.2, $\text{Tr}(S^2) = 0$ and so $\dim(H_0) = \dim(H_1)$. Therefore

$$\dim(H) = \dim(H_0) + \dim(H_1) = 2 \dim(H_1)$$

is a multiple of 4. □

Corollary 2.2. *Let H be a Hopf algebra over k of dimension $2n$ where n is an odd integer. If H is not semisimple, then H or H^* is not unimodular.*

Proof. If both H and H^* are unimodular, by Proposition 1.1, $S^4 = id_H$. Since H is not semisimple, by Theorem 1.2, $o(S^2) = 2$. It follows from Proposition 2.1 that $\dim(H)$ is then a multiple of 4, which contradicts $\dim(H) = 2n$. \square

3. HOPF ALGEBRAS OF DIMENSION $2p$

In this section, we prove, by contradiction, that non-semisimple Hopf algebras over k of dimension $2p$, p an odd prime, do not exist. By [Mas95], semisimple Hopf algebras of dimension $2p$ are

$$k[\mathbb{Z}_{2p}], \quad k[D_{2p}] \quad \text{and} \quad k[D_{2p}]^*$$

where D_{2p} is the dihedral group of order $2p$. Our main result completes the classification of Hopf algebras of dimension $2p$. We begin to prove our main result with the following lemma.

Lemma 3.1. *Let H be a non-semisimple finite-dimensional Hopf algebra over k of dimension $2p$ where p is an odd prime. Suppose that g and α are the distinguished group-like elements of H and H^* respectively. Then*

$$\text{lcm}(o(g), o(\alpha)) = 2 \text{ or } p.$$

Proof. Since H is not semisimple, by Theorem 1.2, H^* is also not semisimple. Therefore $|G(H)|$ and $|G(H^*)|$ are strictly less than $2p$. By the Nichols-Zoeller theorem [NZ89],

$$|G(H)|, |G(H^*)| \in \{1, 2, p\}.$$

It follows from [Ng02b, Lemma 5.1] that

$$\text{lcm}(|G(H)|, |G(H^*)|) = 1, 2 \text{ or } p.$$

Since $\text{lcm}(o(g), o(\alpha))$ divides $\text{lcm}(|G(H)|, |G(H^*)|)$, we obtain

$$\text{lcm}(o(g), o(\alpha)) = 1, 2 \text{ or } p.$$

By Corollary 2.2, $\text{lcm}(o(g), o(\alpha)) > 1$, and so the result follows. \square

Lemma 3.2. *Let H be a finite-dimensional Hopf algebra over k and $a \in G(H)$ of order d . Let $\omega \in k$ be a primitive d th root of unity and*

$$e_i = \frac{1}{d} \sum_{j=0}^{d-1} \omega^{-ij} a^j \quad (i = 0, \dots, d-1).$$

Then $\dim(H)/d$ is an integer, $\dim(He_i) = \dim(H)/d$ and $S^2(He_i) = He_i$ for $i = 0, \dots, d-1$. In addition, if H is not semisimple, then

$$\text{Tr}(S^2|_{He_i}) = 0$$

for $i = 0, \dots, d-1$.

Proof. Let $B = k[a]$. Then B is a Hopf subalgebra of H and $\dim(B) = d$. By the Nichols-Zoeller theorem, H is a free B -module. In particular, $\dim(H)$ is a multiple of d and

$$H \cong B^{\dim(H)/d}$$

as right B -modules. Note that e_0, \dots, e_{d-1} are orthogonal idempotents of B such that

$$1 = e_0 + \dots + e_{d-1},$$

and $Be_i = ke_i$. Therefore,

$$He_i \cong B^{\dim(H)/d}e_i = (Be_i)^{\dim(H)/d} = (ke_i)^{\dim(H)/d},$$

and so $\dim(He_i) = \dim(H)/d$ for $i = 0, \dots, d-1$. Since $S^2(a) = a$, $S^2(e_i) = e_i$ for $i = 0, \dots, d-1$. Therefore,

$$S^2(He_i) = HS^2(e_i) = He_i.$$

If, in addition, H is not semisimple, by Theorem 1.2,

$$\text{Tr}(S^2|_{He_i}) = \text{Tr}(S^2 \circ r(e_i)) = 0 \quad \text{for } i = 0, \dots, d-1. \quad \square$$

Theorem 3.3. *If p is an odd prime, then any Hopf algebra of dimension $2p$ over the field k is semisimple.*

Proof. Suppose there exists a non-semisimple Hopf algebra H of dimension $2p$. Let g and α be the distinguished group-like elements of H and H^* respectively. By Corollary 2.2, g and α cannot both be trivial. By Theorem 1.2, we may simply assume that g is not trivial and $o(g) = d$. Let $\omega \in k$ be a primitive d th root of unity and

$$e_i = \frac{1}{d} \sum_{j=0}^{d-1} \omega^{-ij} g^j \quad (i = 0, \dots, d-1).$$

By Lemma 3.1,

$$\text{lcm}(o(g), o(\alpha)) = 2 \text{ or } p.$$

If $\text{lcm}(o(g), o(\alpha)) = 2$, then $d = 2$ and $S^8 = id_H$ by Proposition 1.1. It follows from Theorem 1.2 that

$$o(S^2) = 2 \quad \text{or} \quad 4.$$

It follows from Lemma 3.2 that

$$\dim(He_i) = p \quad \text{and} \quad \text{Tr}(S^2|_{He_i}) = 0 \quad (i = 0, 1).$$

Since $o(S^2) = 2$ or 4 , there exists $j \in \{0, 1\}$ such that

$$o(S^2|_{He_j}) = 2 \quad \text{or} \quad 4.$$

By Lemma 1.4, $\dim(He_j)$ is even, which contradicts that $\dim(He_j) = p$.

If $\text{lcm}(o(g), o(\alpha)) = p$, then $d = p$ and $S^{4p} = id_H$. By Proposition 1.3, the subspace

$$H_- = \{u \in H \mid S^{2p}(u) = -u\}$$

has even dimension. On the other hand, by Lemma 3.2, we have

$$\dim(He_i) = 2 \quad \text{and} \quad \text{Tr}(S^2|_{He_i}) = 0 \quad \text{for } i = 0, \dots, p-1.$$

Thus, for $i \in \{0, \dots, p-1\}$, there is a basis $\{u_i^+, u_i^-\}$ for He_i such that

$$S^2(u_i^+) = \zeta_i u_i^+ \quad \text{and} \quad S^2(u_i^-) = -\zeta_i u_i^-$$

for some p th root of unity ζ_i . Thus, $\{u_0^-, u_1^-, \dots, u_{p-1}^-\}$ forms a basis of H_- and so

$$\dim(H_-) = p,$$

a contradiction! □

ACKNOWLEDGEMENT

The author would like to thank L. Long for her useful suggestions to this paper.

REFERENCES

- [And02] Nicolás Andruskiewitsch, *About finite dimensional Hopf algebras*, Quantum symmetries in theoretical physics and mathematics (Bariloche, 2000), Contemp. Math., vol. 294, Amer. Math. Soc., Providence, RI, 2002, pp. 1–57. MR1907185 (2003f:16059)
- [AS98] Nicolás Andruskiewitsch and Hans-Jürgen Schneider, *Hopf algebras of order p^2 and braided Hopf algebras of order p* , J. Algebra **199** (1998), no. 2, 430–454. MR1489920 (99c:16033)
- [BD] M. Beattie and S. Dăscălescu, *Hopf algebras of dimension 14*, J. London Math. Soc. (2) **69** (2004), no. 1, 65–78. MR2025327
- [EG] Pavel Etingof and Shlomo Gelaki, *On Hopf algebras of dimension pq* , J. Algebra **277** (2004), no. 2, 668–674. MR2067625
- [EG98] ———, *Semisimple Hopf algebras of dimension pq are trivial*, J. Algebra **210** (1998), no. 2, 664–669. MR1662308 (99k:16079)
- [GW00] Shlomo Gelaki and Sara Westreich, *On semisimple Hopf algebras of dimension pq* , Proc. Amer. Math. Soc. **128** (2000), no. 1, 39–47. MR1618670 (2000c:16050)
- [LR87] Richard G. Larson and David E. Radford, *Semisimple cosemisimple Hopf algebras*, Amer. J. Math. **109** (1987), no. 1, 187–195. MR0926744 (89a:16011)
- [LR88] ———, *Finite-dimensional cosemisimple Hopf algebras in characteristic 0 are semisimple*, J. Algebra **117** (1988), no. 2, 267–289. MR0957441 (89k:16016)
- [Mas95] Akira Masuoka, *Semisimple Hopf algebras of dimension $2p$* , Comm. Algebra **23** (1995), no. 5, 1931–1940. MR1323710 (96e:16050)
- [Mas96] ———, *The p^n theorem for semisimple Hopf algebras*, Proc. Amer. Math. Soc. **124** (1996), no. 3, 735–737. MR1301036 (96f:16046)
- [Mon93] Susan Montgomery, *Hopf algebras and their actions on rings*, CBMS Regional Conference Series in Mathematics, vol. 82, Published for the Conference Board of the Mathematical Sciences, Washington, DC, 1993. MR1243637 (94i:16019)
- [Mon98] ———, *Classifying finite-dimensional semisimple Hopf algebras*, Trends in the representation theory of finite-dimensional algebras (Seattle, WA, 1997), Contemp. Math., vol. 229, Amer. Math. Soc., Providence, RI, 1998, pp. 265–279. MR1676225 (99k:16084)
- [Ng] Siu-Hung Ng, *Hopf algebras of dimension pq* , J. Algebra **276** (2004), no. 1, 399–406. MR2054403
- [Ng02a] ———, *Hopf algebras of dimension p^2* , Proceedings of the International Conference on Hopf Algebras, DePaul University, Chicago (2002).
- [Ng02b] ———, *Non-semisimple Hopf algebras of dimension p^2* , J. Algebra **255** (2002), no. 1, 182–197. MR1935042 (2003h:16067)
- [NZ89] Warren D. Nichols and M. Bettina Zoeller, *A Hopf algebra freeness theorem*, Amer. J. Math. **111** (1989), no. 2, 381–385. MR0987762 (90c:16008)
- [Rad76] David E. Radford, *The order of the antipode of a finite dimensional Hopf algebra is finite*, Amer. J. Math. **98** (1976), no. 2, 333–355. MR0407069 (53:10852)
- [Rad94] ———, *The trace function and Hopf algebras*, J. Algebra **163** (1994), no. 3, 583–622. MR1265853 (95e:16039)
- [Swe69] Moss E. Sweedler, *Hopf algebras*, W. A. Benjamin, Inc., New York, 1969, Mathematics Lecture Note Series. MR0252485 (40:5705)
- [Taf71] Earl J. Taft, *The order of the antipode of finite-dimensional Hopf algebra*, Proc. Nat. Acad. Sci. U.S.A. **68** (1971), 2631–2633. MR0286868 (44:4075)
- [Wil88] R. Williams, *Finite dimensional hopf algebras*, Ph.D. Thesis, Florida State University, 1988.
- [Zhu94] Yongchang Zhu, *Hopf algebras of prime dimension*, Internat. Math. Res. Notices (1994), no. 1, 53–59. MR1255253 (94j:16072)

DEPARTMENT OF MATHEMATICS, IOWA STATE UNIVERSITY, AMES, IOWA 50011
E-mail address: rng@math.iastate.edu