

## SEIBERG-WITTEN INVARIANTS AND BRANCHED COVERS ALONG TORI

B. DOUG PARK

(Communicated by Ronald A. Fintushel)

ABSTRACT. We compute the Seiberg-Witten invariants of double covers of smooth four-manifolds branched along tori of self-intersection zero.

### 1. INTRODUCTION

This short paper is a modest generalization of [RW]. In that paper Ruan and Wang computed (mod 2) the Seiberg-Witten invariants of double covers of 4-manifolds branched along a connected orientable Riemann surface of genus greater than one. In what follows, we shall present a similar (mod 2) computation when the branched locus is a torus. Our shared philosophy is to express the Seiberg-Witten invariants of the closed 4-manifolds in terms of the relative invariants of the complements of the branch locus. The computation depends essentially on the gluing formulae for the Seiberg-Witten invariant found in [P2] or [T]. In the last section, we answer in the affirmative a conjecture made by Ruan and Wang in [RW]. More examples of our formulae will appear in [P3].

### 2. PRELIMINARY SETUP AND DEFINITIONS

We shall try to be faithful to the notation in [RW] and [P2]. Let  $\tilde{X}, X$  be closed smooth 4-manifolds and  $\Sigma \subset X$  be a smoothly embedded connected orientable surface. We say that a smooth map  $p : \tilde{X} \rightarrow X$  is a cyclic  $m$ -fold cover branched along  $\Sigma$  when the following holds: if  $\tilde{\Sigma} = p^{-1}(\Sigma)$ , then the restriction  $p : \tilde{X} \setminus \tilde{\Sigma} \rightarrow X \setminus \Sigma$  is an unbranched  $m$ -fold cover, and  $p$  has the form  $z \mapsto z^m$  locally on the normal complex planes of  $\tilde{\Sigma}$  and  $\Sigma$ .

$$\begin{array}{ccc}
 \tilde{X} & = & \tilde{Y}_0 \cup \nu(\tilde{\Sigma}) \\
 \downarrow p & & \downarrow m:1 \quad \downarrow z \mapsto z^m \\
 X & = & Y_0 \cup \nu(\Sigma)
 \end{array}$$

Let  $\nu(\Sigma)$  denote a tubular neighborhood of  $\Sigma$  and let  $Y_0 := X \setminus \nu(\Sigma)$ ,  $\tilde{Y}_0 := p^{-1}(Y_0)$ . Again,  $p$  restricts to an unbranched cover  $p : \tilde{Y}_0 \rightarrow Y_0$ . When  $m = 2$ , let  $\sigma : \tilde{Y}_0 \rightarrow \tilde{Y}_0$  be the covering involution. We shall reserve the  $\sim$  symbol for indicating some pull-back object by  $p$ . As in [RW] we will always assume that  $[\Sigma]^2 \geq 0$ .

---

Received by the editors February 5, 2004 and, in revised form, May 3, 2004.  
 2000 *Mathematics Subject Classification*. Primary 57M12, 57R57.

©2005 American Mathematical Society

**Definition 2.1.** (i) Given any complex line bundle  $L \rightarrow X$ , the virtual dimension with respect to  $L$  is defined to be

$$d_L := \frac{1}{4}[c_1(L)^2 - (2e_X + 3s_X)],$$

where  $e_X$  and  $s_X$  are respectively the Euler characteristic and the signature of  $X$ .

(ii) The adjunction term of  $\Sigma$  with respect to  $L$  is defined to be

$$J_L(\Sigma) := |c_1(L) \cdot PD[\Sigma]| + [\Sigma] \cdot [\Sigma] + e_\Sigma,$$

where  $PD : H_2(X; \mathbb{Z}) \rightarrow H^2(X; \mathbb{Z})$  is the Poincaré duality isomorphism.

By changing  $[\Sigma]$  to  $-[\Sigma]$  if necessary, we can always assume that  $c_1(L) \cdot PD[\Sigma] \leq 0$ . With that understood, we have the following

**Lemma 2.2.** (i)  $[\tilde{\Sigma}]^2 = \frac{1}{m}[\Sigma]^2$ .

(ii)  $e_{\tilde{X}} = me_X - (m - 1)e_\Sigma$ , and  $s_{\tilde{X}} = ms_X - \frac{m^2-1}{3}[\tilde{\Sigma}]^2$ .

(iii) For any complex line bundle  $L \rightarrow X$ , we have  $c_1(p^*L)^2 = mc_1(L)^2$ .

(iv) Let  $\hat{L}$  be a line bundle such that  $c_1(\hat{L}) = c_1(p^*L) - (m - 1)PD[\tilde{\Sigma}]$ . Then  $J_{\hat{L}}(\tilde{\Sigma}) = J_L(\Sigma)$ .

(v)  $d_{\hat{L}} = md_L + \frac{1}{2}(m - 1)J_L(\Sigma)$ .

*Proof.* We refer to the excellent book [GS] for some of the proofs and [RW] for the special case when  $m = 2$ . The signature formula in part (ii) can be found in [H]. The rest is an easy exercise, which we omit. □

From now on we will restrict our attention to the case when the genus of  $\Sigma$  is one and  $[\Sigma]^2 = 0$ . We now briefly recall the setup in [P2]. Suppose we are given a compact 4-manifold  $Y_0$  with boundary  $\partial Y_0 \cong T^3$  and a fixed factorization  $\partial Y_0 = \Sigma \times S^1$ , where  $\Sigma = T^2$ . Let  $\gamma = [\{\text{pt}\} \times S^1] \in H_1(\partial Y_0)$ , and let  $\gamma^* \in H^1(\partial Y_0)$  be the Poincaré dual of  $[\Sigma] \in H_2(\partial Y_0)$ . Let  $X$  be the closed 4-manifold  $Y_0 \cup_{\partial Y_0} (\Sigma \times D^2)$ , where the boundaries are identified using the above factorization.

**Definition 2.3.** Let  $\gamma$  denote the homology class of the meridian of  $\Sigma$  in  $\partial Y_0$  as above, and let  $i : \partial Y_0 \hookrightarrow Y_0$  be the inclusion map. We shall say that the pair  $(Y_0, \Sigma)$  is *admissible* if the following three conditions are satisfied:

- (i)  $n\gamma \in \text{Ker}[i_* : H_1(\partial Y_0; \mathbb{Z}) \rightarrow H_1(Y_0; \mathbb{Z})]$  for some positive integer  $n$ .
- (ii)  $\text{Coker}[i^* : H^1(Y_0; \mathbb{Z}) \rightarrow H^1(\partial Y_0; \mathbb{Z})]$  is torsion-free.
- (iii) The intersection form of  $Y_0$  is not negative definite.

For an admissible pair  $(Y_0, \Sigma)$ , we define

$$n_\gamma := \min\{n > 0 \mid n\gamma \in \text{Ker}[i_* : H_1(\partial Y_0; \mathbb{Z}) \rightarrow H_1(Y_0; \mathbb{Z})]\},$$

$$n_\Sigma := \max\{n > 0 \mid [\Sigma] = n\alpha \text{ for some } \alpha \in H_2(X; \mathbb{Z})\}.$$

We shall say that the pair  $(Y_0, \Sigma)$  is *strongly admissible* if it is admissible and  $n_\gamma = n_\Sigma$ .

Note that condition (i) above implies the existence of an embedded surface  $\Gamma_0 \subset Y_0$  such that  $[\partial\Gamma_0] = n_\gamma\gamma$ . Let  $\Gamma$  denote the closed connected surface

$$(2.1) \quad \Gamma_0 \cup \left( \prod_{i=1}^{n_\gamma} \{\text{pt}_i\} \times D^2 \right) \subset X.$$

It follows that  $[\Gamma] \in H_2(X; \mathbb{Z})$  is primitive and  $[\Gamma] \cdot [\Sigma] = n_\gamma$ .  $\Gamma$  is not unique, but we may choose  $\Gamma$  so that its genus is minimal among all such surfaces.

Note that for an  $m$ -fold branched cover,  $m$  divides  $n_\Sigma$ . Let  $\Theta$  denote a smoothly embedded surface in  $X$  such that  $[\Sigma] = n_\Sigma[\Theta]$ . Let  $N$  denote the regular neighborhood of the union  $\Theta \cup \Gamma$  inside  $X$ . If  $(Y_0, \Sigma)$  is strongly admissible, then  $[\Gamma] \cdot [\Theta] = 1$ , and hence there is an orthogonal decomposition

$$H_2(X; \mathbb{Z}) = \langle [\Theta], [\Gamma] \rangle \oplus H_2(X \setminus N; \mathbb{Z})$$

and the corresponding splitting of the intersection form:

$$(2.2) \quad Q_X = \begin{bmatrix} 0 & 1 \\ 1 & c \end{bmatrix} \oplus Q_{X \setminus N},$$

where  $c = [\Gamma] \cdot [\Gamma]$ .

If  $(Y_0, \Sigma)$  is admissible, then we can define a relative Seiberg-Witten invariant of  $Y_0$  as follows. Let  $\mathcal{S}(Y_0)$  denote the set of isomorphism classes of  $\text{Spin}^c$  structures on  $Y_0$  that restricts to the trivial (canonical)  $\text{Spin}^c$  structure on the boundary  $\partial Y_0$ . Let  $\mathcal{H}_{Y_0}$  denote the cokernel of the homomorphism  $i^* : H^1(Y_0; \mathbb{Z}) \rightarrow H^1(\partial Y_0; \mathbb{Z})$ . In [P2], we defined a function

$$SW_{Y_0} : \mathcal{S}(Y_0) \times \mathcal{H}_{Y_0} \longrightarrow \mathbb{Z},$$

which algebraically counts the number of solutions to the twice-perturbed Seiberg-Witten equations for a  $\text{Spin}^c$  structure  $\mathcal{L} \in \mathcal{S}(Y_0)$  on the cylindrical end 4-manifold,  $Y := Y_0 \cup_{\partial Y_0} (\partial Y_0 \times [0, \infty))$ :

$$(2.3) \quad \begin{cases} \not\partial_A \phi = 0, \\ \rho(F_A + \eta) = q(\phi), \\ \eta = f \cdot (ih_{(A, \phi)}^*(\omega) - ir\pi_1^* \pi^* \mu_\Sigma). \end{cases}$$

We refer the reader to [P2] for the exact definition of the terms in (2.3). The cylindrical end moduli space  $\mathcal{M}_Y^r(\mathcal{L}, g, \omega)$  is defined by dividing the space of finite energy solutions to (2.3) by the action of the  $L^2_{5, \text{loc}}$  gauge group  $\mathcal{G}(Y)$ . Note that every solution to (2.3) is irreducible, i.e.  $\phi \neq 0$ . Recall that there exists a continuous map

$$\partial_\infty : \mathcal{M}_Y^r(\mathcal{L}, g, \omega) \longrightarrow \mathcal{H}_{Y_0}.$$

If  $\mathcal{M}_Y^r(\mathcal{L}, g, \omega)$  is 0-dimensional, then we define

$$SW_{Y_0}(\mathcal{L}, x) := \#[\mathcal{M}_Y^r(\mathcal{L}, g, \omega) \cap \partial_\infty^{-1}(x)].$$

**Definition 2.4.** (i) Let  $\mathcal{K}_{Y_0}$  be the set of isomorphism classes of complex line bundles on  $Y_0$  that pull back to a trivial bundle on  $p^{-1}(Y_0)$ . Let  $\mathcal{K}_{Y_0}^* = \mathcal{K}_{Y_0} \setminus \{\underline{\mathbb{C}}\}$  be the subset of non-trivial line bundles.

(ii) Suppose  $(Y_0, \Sigma)$  is strongly admissible. For any  $\text{Spin}^c$  structure  $\xi$  on  $X$  with  $\det \xi = L$  and  $J_L(\Sigma) = -c_1(L) \cdot PD[\Sigma] = 0$ , we let

$$m_L := c_1(L) \cdot PD[\Gamma],$$

where  $\Gamma \subset X$  is defined as in (2.1). We also define

$$k_\xi(X, \Sigma) := \sum_{\kappa \in \mathcal{K}_{Y_0}^*} \sum_{n=1}^{\infty} SW_{Y_0}(\xi|_{Y_0} \otimes \kappa, (\llbracket m_L/n_\gamma \rrbracket - n)\gamma^*),$$

where  $\llbracket m_L/n_\gamma \rrbracket$  denotes the greatest integer less than or equal to  $m_L/n_\gamma$ , and  $\gamma^*$  is the non-zero element of  $\mathcal{H}_{Y_0}$  coming from  $PD[\Sigma] \in H^1(\partial Y_0)$ .

Note that  $\mathcal{K}_{Y_0}$  is isomorphic to the kernel of the homomorphism  $p^* : H^2(Y_0; \mathbb{Z}) \rightarrow H^2(\tilde{Y}_0; \mathbb{Z})$ . Also recall that  $c_1(\kappa)$  is  $m$ -torsion for every line bundle  $\kappa \in \mathcal{K}_{Y_0}$ . (See the proof of Theorem 3.8 in [RW] for the case of double covers.) For any complex line bundle  $L \rightarrow X$ , let us write  $c_1(L) = aPD[\Theta] + bPD[\Gamma] + \beta$ , with  $\beta \in H^2(X \setminus N; \mathbb{Z})$  according to the decomposition (2.2). Then  $c_1(L) \cdot PD[\Sigma] = 0$  implies that  $b = 0$  and  $a = m_L$ . When  $b_2^+(X) > 1$ , one can show that the sum defining  $k_\xi(X, \Sigma)$  is finite as in [P1].

### 3. MAIN FORMULA

We now restrict ourselves to the case when  $m = 2$ . The following is a direct generalization of Theorem 6.8 in [RW].

**Theorem 3.1.** *Let  $p : \tilde{X} \rightarrow X$  be a double cover branched along a homologically non-trivial torus  $\Sigma \subset X$ . Let  $Y_0 = X \setminus \nu(\Sigma)$  be the complement of the tubular neighborhood of  $\Sigma$ , and let  $\tilde{Y}_0 = p^{-1}(Y_0)$ ,  $\tilde{\Sigma} = p^{-1}(\Sigma)$ . Assume that  $b_2^+(X), b_2^+(\tilde{X}) > 1$  and  $[\Sigma]^2 = 0$ . Suppose that  $\xi$  is a  $\text{Spin}^c$  structure on  $X$  whose determinant line bundle  $L$  satisfies  $c_1(L) \cdot PD[\Sigma] = 0$  and  $d_L = 0$ . Let  $\mathcal{L}$  denote the restriction  $\xi|_{Y_0}$ . If both  $(\tilde{Y}_0, \tilde{\Sigma})$  and  $(Y_0, \Sigma)$  are admissible, then we have*

$$SW_{\tilde{Y}_0}(p^*(\mathcal{L}), p^*(x)) \equiv SW_{Y_0}(\mathcal{L}, x) + \sum_{\kappa \in \mathcal{K}_{Y_0}^*} SW_{Y_0}(\mathcal{L} \otimes \kappa, x) \pmod{2}.$$

Moreover let  $\hat{\xi}$  be a  $\text{Spin}^c$  structure on  $\tilde{X}$  whose determinant bundle is  $\hat{L} = p^*L \otimes PD[\tilde{\Sigma}]^{-1}$  and whose restriction to  $\tilde{Y}_0$  is  $p^*(\mathcal{L})$ . Also define

$$\Lambda_L(\Sigma) := \frac{m_L}{n_\Sigma} - \left\lfloor \frac{m_L}{n_\Sigma} \right\rfloor.$$

Suppose both  $(\tilde{Y}_0, \tilde{\Sigma})$  and  $(Y_0, \Sigma)$  are strongly admissible. If  $\Lambda_L(\Sigma) \neq \frac{1}{2}$ , then we have

$$(3.1) \quad SW_{\tilde{X}}(\hat{\xi}) \equiv SW_X(\xi) + k_\xi(X, \Sigma) \pmod{2}.$$

If  $\Lambda_L(\Sigma) = \frac{1}{2}$ , then we have

$$SW_X(\xi) + k_\xi(X, \Sigma) \equiv 0 \pmod{2}.$$

*Proof.* Let  $\tilde{Y} := \tilde{Y}_0 \cup_{\partial \tilde{Y}_0} (\partial \tilde{Y}_0 \times [0, \infty))$ . Consider the following system of equations for the  $\text{Spin}^c$  structure  $p^*(\mathcal{L})$  on  $\tilde{Y}$ :

$$(3.2) \quad \begin{cases} \tilde{\partial}_A \phi = 0, \\ \rho(F_A + \tilde{\eta}) = q(\phi), \\ \tilde{\eta} = (f \circ p) \cdot (ih_{(A, \phi)}^*(p^*\omega) - irp^*\pi_1^*\pi^*\mu_\Sigma). \end{cases}$$

Here,  $\tilde{\partial}_A$  is a twisted Dirac operator constructed from the Levi-Civita connection of the pull-back metric  $p^*g$  on  $\tilde{Y}$ . ( $g$  is a generic cylindrical end metric on  $Y$  used to define (2.3).) Let  $\tilde{\mathcal{M}}_{\tilde{Y}}^r(p^*\mathcal{L}, p^*g, p^*\omega)$  denote the solution space of (3.2) divided by the action of a suitable gauge group  $\mathcal{G}(\tilde{Y})$ . We can choose  $\omega$  such that both  $\omega$  and  $p^*\omega$  are supported away from the critical values of the Chern-Simons-Dirac functional of [P2]. Hence we can prove the compactness the same way and obtain a continuous map

$$\partial_\infty : \tilde{\mathcal{M}}_{\tilde{Y}}^r(p^*\mathcal{L}, p^*g, p^*\omega) \longrightarrow \mathcal{H}_{\tilde{Y}_0}.$$

Note that  $d_L = 0$  implies that the virtual dimension of  $\mathcal{M}_Y^r(\mathcal{L}, g, \omega)$  is zero. Since we have also assumed that  $J_L(\Sigma) = 0$ , it follows that  $d_{\tilde{L}} = 0$ , and hence the virtual dimension of  $\widetilde{\mathcal{M}}_{\tilde{Y}}^r(p^*\mathcal{L}, p^*g, p^*\omega)$  is also zero. We introduce the notation

$$\begin{aligned} \widetilde{\mathcal{N}}_{\tilde{Y}_0}(p^*\mathcal{L}, p^*x; r, p^*g, p^*\omega) &:= \widetilde{\mathcal{M}}_{\tilde{Y}}^r(p^*\mathcal{L}, p^*g, p^*\omega) \cap \partial_{\infty}^{-1}(p^*x), \\ \mathcal{N}_{Y_0}(\mathcal{L}, x; r, g, \omega) &:= \mathcal{M}_Y^r(\mathcal{L}, g, \omega) \cap \partial_{\infty}^{-1}(x). \end{aligned}$$

Note that there is a well-defined homomorphism  $p^* : \mathcal{H}_{Y_0} \rightarrow \mathcal{H}_{\tilde{Y}_0}$ .

Next let  $P_{SO(4)}$  denote the frame bundle of  $\tilde{Y}_0$ , and let  $\tilde{P}$  be the principal  $\text{Spin}^c(4)$  bundle corresponding to  $p^*\mathcal{L}$ . As in [RW] we choose a lifting  $\tau$  of the covering involution  $\sigma$ :

$$\begin{array}{ccc} \tilde{P} & \xrightarrow{\tau} & \tilde{P} \\ \downarrow & & \downarrow \\ P_{SO(4)} & \xrightarrow{\sigma^*} & P_{SO(4)} \end{array}$$

Let  $\tau^*$  denote the induced action on the Seiberg-Witten configuration space  $\mathcal{B}(p^*\mathcal{L})$ . Given any subset  $S \subset \mathcal{B}(p^*\mathcal{L})$ , we let  $S^\tau \subset S$  be the fixed point set of  $\tau^* : \mathcal{B}(p^*\mathcal{L}) \rightarrow \mathcal{B}(p^*\mathcal{L})$ . Since every solution to (3.2) is irreducible, Theorem 3.8 of [RW] gives us a homeomorphism

$$\begin{aligned} &\widetilde{\mathcal{N}}_{\tilde{Y}_0}(p^*\mathcal{L}, p^*x; r, p^*g, p^*\omega)^\tau \\ &\approx \mathcal{N}_{Y_0}(\mathcal{L}, x; r, g, \omega) \prod \left( \prod_{\kappa \in \mathcal{K}_{Y_0}^*} \mathcal{N}_{Y_0}(\mathcal{L} \otimes \kappa, x; r, g, \omega) \right). \end{aligned}$$

Now let  $g'$  be a generic cylindrical end metric on  $\tilde{Y}$ . We need to compare the moduli spaces of (2.3) and (3.2) on  $\tilde{Y}$ . Since  $p^*\pi_1^*\pi^*\mu_\Sigma = 2\pi_1^*\pi^*\mu_{\tilde{\Sigma}}$  and we are free to vary the parameter  $r \in \mathbb{R} \setminus \{0\}$  in (2.3) (as long as  $|r|$  is very small and the sign of  $r$  remains the same), we easily see that the only difference between  $\mathcal{M}_Y^{2r}(p^*\mathcal{L}, g', p^*\omega)$  and  $\widetilde{\mathcal{M}}_{\tilde{Y}}^r(p^*\mathcal{L}, p^*g, p^*\omega)$  is in the non-generic choice of the metric  $p^*g$  in the definition of (3.2). Hence Theorem 2.2 of [RW] implies that

$$\begin{aligned} SW_{\tilde{Y}_0}(p^*\mathcal{L}, p^*x) &= \#[\mathcal{N}_{\tilde{Y}_0}(p^*\mathcal{L}, p^*x; 2r, g', p^*\omega)] \\ &\equiv \#[\widetilde{\mathcal{N}}_{\tilde{Y}_0}(p^*\mathcal{L}, p^*x; r, p^*g, p^*\omega)^\tau] \pmod{2}. \end{aligned}$$

This proves our first congruence. To prove the second, recall from [P2] (Corollary 20) that

$$(3.3) \quad \overline{SW}_X = \overline{SW}_{Y_0} \cdot ([\Sigma] + [\Sigma]^3 + [\Sigma]^5 + \dots),$$

where  $\overline{SW}$  denotes the Seiberg-Witten series. Hence it follows from the product formula in [P2] that

$$(3.4) \quad SW_X(\xi) = \sum_{n=1}^{\infty} SW_{Y_0}(\mathcal{L}, ([m_L/n_\Sigma] - 2n + 1)\gamma^*).$$

(Note that  $PD[\Theta] \in H^2(Y_0; \mathbb{Z}) \cong H_2(Y_0, \partial Y_0; \mathbb{Z})$  is  $n_\Sigma$ -torsion.)  $\gamma^*$  is indivisible and since the difference (modulo torsion) of any two characteristic elements of  $H^2(Y_0, \partial Y_0; \mathbb{Z})$  is divisible by two (cf. [MMS], [P1]), we must have by default

$$(3.5) \quad SW_{Y_0}(\mathcal{L}, ([m_L/n_\Sigma] - 2n)\gamma^*) = 0$$

for every integer  $n$ . Thus we may rewrite (3.4) as

$$SW_X(\xi) = \sum_{n=1}^{\infty} SW_{Y_0}(\mathcal{L}, (\lfloor m_L/n_\Sigma \rfloor - n)\gamma^*).$$

Summing our first congruence over  $n$ , we conclude that

$$\begin{aligned} & SW_X(\xi) + k_\xi(X, \Sigma) \\ &= \sum_{n=1}^{\infty} \left( SW_{Y_0}(\mathcal{L}, (\lfloor \frac{m_L}{n_\Sigma} \rfloor - n)\gamma^*) + \sum_{\kappa \in \mathcal{K}_{Y_0}^*} SW_{Y_0}(\mathcal{L} \otimes \kappa, (\lfloor \frac{m_L}{n_\Sigma} \rfloor - n)\gamma^*) \right) \\ &\equiv \sum_{n=1}^{\infty} SW_{\tilde{Y}_0}(p^*\mathcal{L}, p^*(\lfloor m_L/n_\Sigma \rfloor - n)\gamma^*) \pmod{2}. \end{aligned}$$

Let  $\delta \in H_1(\partial\tilde{Y}_0)$  denote the homology class of the meridian of  $\tilde{\Sigma}$ , and let  $\delta^* := PD[\tilde{\Sigma}] \in H^1(\partial\tilde{Y}_0)$ . As in (2.1), let  $\Delta \subset \tilde{X}$  be a minimal genus surface such that  $[\tilde{\Sigma}] \cdot [\Delta] = n_\delta = n_{\tilde{\Sigma}}$ . Note that  $[\Sigma] \cdot p_*[\Delta] = 2n_\delta = n_\gamma$ . Since  $p^*(\gamma^*) = 2\delta^*$ , we get

$$(3.6) \quad SW_X(\xi) + k_\xi(X, \Sigma) \equiv \sum_{n=1}^{\infty} SW_{\tilde{Y}_0}(p^*\mathcal{L}, 2(\lfloor m_L/n_\Sigma \rfloor - n)\delta^*) \pmod{2}.$$

As in (3.4), we have

$$SW_{\tilde{X}}(\hat{\xi}) = \sum_{n=1}^{\infty} SW_{\tilde{Y}_0}(p^*\mathcal{L}, (\lfloor m_{\hat{L}}/n_{\tilde{\Sigma}} \rfloor - 2n + 1)\delta^*).$$

But now note that

$$m_{\hat{L}} = c_1(\hat{L}) \cdot PD[\Delta] = -n_\delta + c_1(p^*L) \cdot PD[\Delta] = -n_\delta + m_L.$$

Since  $n_\Sigma = 2n_{\tilde{\Sigma}}$  (cf. [GS]), we must have

$$\frac{m_{\hat{L}}}{n_{\tilde{\Sigma}}} = \frac{-n_\delta + m_L}{n_\Sigma/2} = -1 + \frac{2m_L}{n_\Sigma}.$$

Hence it follows that

$$(3.7) \quad SW_{\tilde{X}}(\hat{\xi}) = \sum_{n=1}^{\infty} SW_{\tilde{Y}_0}(p^*\mathcal{L}, (\lfloor 2m_L/n_\Sigma \rfloor - 2n)\delta^*).$$

Let  $\bar{\xi}$  be the complex conjugate  $\text{Spin}^c$  structure on  $X$  with  $\det \bar{\xi} = -L$ . We have  $SW_X(\bar{\xi}) = (-1)^{(e_X + s_X)/4} SW_X(\xi)$ . Hence by changing  $\xi$  to  $\bar{\xi}$  if necessary, we may assume that

$$0 \leq \frac{m_L}{n_\Sigma} - \left\lfloor \frac{m_L}{n_\Sigma} \right\rfloor \leq \frac{1}{2}.$$

Now if  $m_L/n_\Sigma - \lfloor m_L/n_\Sigma \rfloor \neq 1/2$ , then we get

$$\lfloor 2m_L/n_\Sigma \rfloor = 2\lfloor m_L/n_\Sigma \rfloor.$$

Thus the combination of (3.6) and (3.7) proves our second congruence, provided that  $m_L/n_\Sigma - \lfloor m_L/n_\Sigma \rfloor \neq 1/2$ . Finally suppose that

$$(3.8) \quad m_L = \left\lfloor \frac{m_L}{n_\Sigma} \right\rfloor n_\Sigma + \frac{n_\Sigma}{2}.$$

Then we have  $\lfloor 2m_L/n_\Sigma \rfloor = 2\lfloor m_L/n_\Sigma \rfloor + 1$ , and thus

$$SW_{\widehat{X}}(\widehat{\xi}) = \sum_{n=1}^{\infty} SW_{\widehat{Y}_0}(p^*\mathcal{L}, (2\lfloor m_L/n_\Sigma \rfloor + 1 - 2n)\delta^*).$$

As in (3.5), it follows that

$$SW_{\widehat{Y}_0}(p^*\mathcal{L}, (2\lfloor m_L/n_\Sigma \rfloor - 2n)\delta^*) = 0$$

for every integer  $n$ . Hence (3.6) implies that

$$SW_X(\xi) + k_\xi(X, \Sigma) \equiv 0 \pmod{2}. \quad \square$$

*Remark 3.2.* (i) If  $b_2^+(X) > 1$ , then by the adjunction inequality (cf. [OS]),  $c_1(L) \cdot PD[\Sigma] \neq 0$  implies that  $SW_X(\xi) = 0$ . Since  $c_1(\widehat{L}) \cdot PD[\widehat{\Sigma}] = c_1(L) \cdot PD[\Sigma]$ , it also implies that if  $b_2^+(\widehat{X}) > 1$ , then we must have  $SW_{\widehat{X}}(\widehat{\xi}) = 0$  as well.

(ii) If  $c_1(L) \cdot PD[\Sigma] = 0$  but  $d_L \neq 0$ , then by Lemma 2.2 (v) we have  $d_{\widehat{L}} \neq 0$ . Hence if  $\widehat{X}$  is of Seiberg-Witten simple-type (e.g. if  $\widehat{X}$  is a symplectic manifold), then  $SW_{\widehat{X}}(\widehat{\xi}) = 0$ .

(iii) Let  $\lambda \rightarrow X$  be the complex line bundle with  $c_1(\lambda) = \frac{1}{2}PD[\Sigma]$  determined by the branched cover  $p : \widetilde{X} \rightarrow X$ . If  $(Y_0, \Sigma)$  is admissible, then  $[\Sigma] \in H_2(X; \mathbb{Z})$  is not torsion. Hence Proposition 5.3 of [RW] implies that  $\lambda|_{Y_0} \in \mathcal{K}_{Y_0}^*$ , and if  $H_1(X; \mathbb{Z})$  contains no 2-torsion, then  $\mathcal{K}_{Y_0}^* = \{\lambda|_{Y_0}\}$ .

#### 4. EXAMPLES

We start out with the following conjecture made by Ruan and Wang:

**Conjecture 4.1** (cf. [RW], p. 502). *The difference of the SW-invariants,*

$$SW_{\widehat{X}}(\widehat{\xi}) - SW_X(\xi) \pmod{2},$$

*is not always zero.*

We prove the conjecture (at least when the genus of  $\Sigma$  is one) by providing examples with  $k_\xi(X, \Sigma) \equiv 1 \pmod{2}$ .

For each integer  $n > 0$ , let  $E(n)$  be a simply-connected elliptic surface with no multiple fibers and with geometric genus  $p_g = n - 1$ . Let  $F$  denote a generic torus fiber of  $E(n)$ , and let  $E(n)_q$  be the result of a logarithmic transformation of multiplicity  $q > 1$  along  $F$ . Let  $F_q \subset E(n)_q$  denote the multiple fiber. Recall that  $[F] = q[F_q]$  in  $H_2(E(n)_q; \mathbb{Z})$ . If  $q = 2\ell$  is even, then let  $\widetilde{E}(n)_q$  be the double cover of  $E(n)_q$  branched along the torus  $\Sigma = F$ .

We may take  $\Gamma$  to be the union of a normal disk of  $F_q$  and  $q$  punctured sections in  $E(n) \setminus \nu(F)$ , and set  $\Theta = F_q$ . It is not very hard to check that the pairs  $(\widetilde{E}(n)_q \setminus \nu(\widetilde{F}), \widetilde{F})$  and  $(E(n)_q \setminus \nu(F), F)$  are strongly admissible. (Note that  $\mathcal{H}_{\widetilde{E}(n)_q \setminus \nu(\widetilde{F})} \cong \mathcal{H}_{E(n)_q \setminus \nu(F)} \cong \mathbb{Z}^3$ .)

**Lemma 4.2.** *Let  $t = [F]$  and  $t_q = [F_q]$  in  $H_2(E(n)_q; \mathbb{Z})$ . Then we have*

$$\begin{aligned} \overline{SW}_{E(n)_q} &= (t^{-1} - t)^{n-2} \cdot (t_q^{q-1} + t_q^{q-3} + \dots + t_q^{-(q-3)} + t_q^{-(q-1)}), \\ \overline{SW}_{E(n)_q \setminus \nu(F)} &= (t^{-1} - t)^{n-1} \cdot (t_q^{q-1} + t_q^{q-3} + \dots + t_q^{-(q-3)} + t_q^{-(q-1)}). \end{aligned}$$

*Proof.*  $\overline{SW}_{E(n)_q}$  was computed in [FS]. Apply Corollary 20 of [P2] to it. (See formula (3.3) above.)  $\square$

**Theorem 4.3.** *Let  $\xi_j$  denote the  $\text{Spin}^c$  structure on  $E(2)_{2\ell}$  with  $c_1(\det \xi_j) = PD(j[F_{2\ell}])$ . If  $j$  is odd and satisfies  $1 \leq j \leq 2\ell - 1$ , then*

$$k_{\xi_j}(E(2)_{2\ell}, F) = 1.$$

Moreover, if  $j \neq \ell$ , then

$$SW_{\widetilde{E(2)_{2\ell}}}(\widehat{\xi}_j) - SW_{E(2)_{2\ell}}(\xi_j) \equiv 1 \pmod{2}.$$

Therefore Conjecture 4.1 is true.

*Proof.* As before let  $Y_0 = E(2)_{2\ell} \setminus \nu(F)$  and let  $L_j = \det \xi_j$ . Let  $t = [F]$  and  $t_q = [F_{2\ell}]$  with  $q = 2\ell$ . Note that  $n_F = 2\ell$  and  $m_{L_j} = j$ . Thus  $\llbracket m_{L_j}/n_F \rrbracket = 0$ . Since  $E(n)_q$  is simply-connected,  $\mathcal{K}_{Y_0}^* = \{\lambda|_{Y_0}\}$ , where  $c_1(\lambda) = \ell PD(t_q)$ . Thus if  $1 \leq j \leq 2\ell - 1$ , we have

$$k_{\xi_j}(E(2)_{2\ell}, F) = \sum_{n=1}^{\infty} SW_{Y_0}(\xi_j|_{Y_0} \otimes \lambda|_{Y_0}, (-n)\gamma^*).$$

Now recall from [P2] that

$$SW_{Y_0}(\xi_j|_{Y_0} \otimes \lambda|_{Y_0}, m\gamma^*) = SW_{Y_0}(\xi_j|_{Y_0}, (m - 1)\gamma^*)$$

for every integer  $m$ . Hence we conclude that

$$k_{\xi_j}(E(2)_{2\ell}, F) = \sum_{n=1}^{\infty} SW_{Y_0}(\xi_j|_{Y_0}, (-n - 1)\gamma^*).$$

From Lemma 4.2, we have

$$\overline{SW}_{Y_0} = (t^{-1} - t) \cdot (t_q^{q-1} + t_q^{q-3} + \dots + t_q^{-(q-3)} + t_q^{-(q-1)}).$$

Let  $a_{n,j}$  denote the coefficient of  $t^n t_q^j$  in  $\overline{SW}_{Y_0}$  with  $|j| \leq q - 1$ . From the definition of a Seiberg-Witten series in [P2], we deduce that if  $|j| \leq q - 1$ , then

$$a_{n,j} = SW_{Y_0}(\xi_j|_{Y_0}, n\gamma^*).$$

It follows that if  $j$  is odd and  $1 \leq j \leq 2\ell - 1$ , then

$$k_{\xi_j}(E(2)_{2\ell}, F) = \sum_{n=1}^{\infty} a_{-n-1,j} = a_{-2,j} = a_{-1,j-2\ell} = 1.$$

For the last statement note that if  $1 \leq j \leq 2\ell - 1$  and  $j \neq \ell$ , then

$$\Lambda_{L_j}(F) = \frac{m_{L_j}}{n_F} - \left\llbracket \frac{m_{L_j}}{n_F} \right\rrbracket = \frac{j}{2\ell} - \left\llbracket \frac{j}{2\ell} \right\rrbracket = \frac{j}{2\ell} \neq \frac{1}{2}.$$

Now apply congruence (3.1) of Theorem 3.1. □

*Remark 4.4.* (i) For  $n > 2$ , let  $\xi_{i,j}$  be the  $\text{Spin}^c$  structure on  $E(n)_{2\ell}$  such that  $c_1(\det \xi_{i,j}) = PD(it + jt_q)$  with  $|j| < 2\ell$ . As in the last proof, we can express  $k_{\xi_{i,j}}(E(n)_{2\ell}, F)$  as a sum of coefficients of  $\overline{SW}_{E(n)_{2\ell} \setminus \nu(F)}$ . In this way we can find a plethora of instances where  $k_{\xi_{i,j}}(E(n)_{2\ell}, F) \equiv SW_{\widetilde{E(n)_{2\ell}}}(\widehat{\xi}_{i,j}) - SW_{E(n)_{2\ell}}(\xi_{i,j}) \equiv 1 \pmod{2}$ .

(ii) If  $\ell$  is odd and  $j = \ell$ , then  $\Lambda_{L_j}(F) = \frac{1}{2}$  and  $SW_{E(2)_{2\ell}}(\xi_j) = 1$ . Thus we have

$$SW_{E(2)_{2\ell}}(\xi_j) + k_{\xi_j}(E(2)_{2\ell}, F) = 2 \equiv 0 \pmod{2},$$

which is the final congruence of Theorem 3.1.

(iii) Many more examples with  $k_{\xi}(X, \Sigma) \equiv 1 \pmod{2}$  can be found in [P3].

## ACKNOWLEDGMENTS

The author would like to thank Ian Hambleton for introducing him to the field of equivariant gauge theory and providing inspiration. He would also like to thank Tian-Jun Li and Michael McCooey for very helpful discussions.

## REFERENCES

- [FS] R. Fintushel and R. J. Stern: Rational blowdowns of smooth 4-manifolds, *J. Differential Geom.* **46** (1997), 181–235. MR1484044 (98j:57047)
- [GS] R. E. Gompf and A. I. Stipsicz: *4-Manifolds and Kirby Calculus*, Graduate Studies in Mathematics, No. 20, Amer. Math. Soc., 1999. MR1707327 (2000h:57038)
- [H] F. Hirzebruch: The signature of ramified coverings, in *Global Analysis (Papers in Honor of K. Kodaira)*, pp. 253–265, Univ. Tokyo Press, Tokyo, 1969. MR0258060 (41:2707)
- [MMS] J. W. Morgan, T. S. Mrowka and Z. Szabó: Product formulas along  $T^3$  for Seiberg-Witten invariants, *Math. Res. Lett.* **4** (1997), 915–929. MR1492130 (99f:57039)
- [OS] P. Ozsváth and Z. Szabó: Higher type adjunction inequalities in Seiberg-Witten theory, *J. Differential Geom.* **55** (2000), 385–440. MR1863729 (2002j:57061)
- [P1] B. D. Park: A product formula for the Seiberg-Witten invariant along certain Seifert fibered manifolds, *Asian J. Math.* **6** (2002), 37–60. MR1902646 (2003b:57047)
- [P2] B. D. Park: A gluing formula for the Seiberg-Witten invariant along  $T^3$ , *Michigan Math. J.* **50** (2002), 593–611. MR1935154 (2003i:57051)
- [P3] B. D. Park: Constraints on Alexander polynomials of certain two-component links, *Topology Appl.* **144** (2004), 161–171. MR2097134
- [RW] Y. Ruan and S. Wang: Seiberg-Witten invariants and double covers of 4-manifolds, *Comm. Anal. Geom.* **8** (2000), 477–515. MR1775135 (2001h:57039)
- [T] C. H. Taubes: The Seiberg-Witten invariants and 4-manifolds with essential tori, *Geom. Topol.* **5** (2001), 441–519. MR1833751 (2002d:57025)

DEPARTMENT OF PURE MATHEMATICS, UNIVERSITY OF WATERLOO, WATERLOO, ONTARIO, CANADA N2L 3G1

*E-mail address:* [bdpark@math.uwaterloo.ca](mailto:bdpark@math.uwaterloo.ca)

*URL:* <http://www.math.uwaterloo.ca/~bdpark/>