

JORDAN ISOMORPHISMS OF TRIANGULAR RINGS

TSAI-LIEN WONG

(Communicated by David R. Larson)

ABSTRACT. We investigate Jordan isomorphisms of triangular rings and give a sufficient condition under which they are necessarily isomorphisms or anti-isomorphisms. As corollaries we obtain generalizations of two recent results: the one concerning Jordan isomorphisms of triangular matrix algebras by Beidar, Brešar and Chebotar, and the one concerning Jordan isomorphisms of nest algebras by Lu.

1. INTRODUCTION

Let \mathcal{T} and \mathcal{S} be rings. A bijective additive map $\varphi : \mathcal{T} \rightarrow \mathcal{S}$ is called a *Jordan isomorphism* if $\varphi(xy+yx) = \varphi(x)\varphi(y)+\varphi(y)\varphi(x)$ for all $x, y \in \mathcal{T}$. Isomorphisms and anti-isomorphisms are obvious examples, and the usual goal is to describe a Jordan isomorphism through these two examples. This problem has a long history; the initial results were obtained already in the 40s and the 50s [1, 2, 9, 10, 11, 12, 17]. From a classical theorem of Herstein [9] (together with a technical improvement by Smiley [17]) it follows that every Jordan isomorphism between prime rings of characteristic not 2 is either an isomorphism or an anti-isomorphism. The situation where the rings are semiprime is more involved, but also well understood [3, 5, 6], and so the problem is now interesting for rings containing nonzero nilpotent ideals.

We denote by $\mathcal{T}_r(R)$ the ring of all $r \times r$ upper triangular matrices over a ring R . In 1998, Molnár and Šemrl [14] proved that automorphisms and antiautomorphisms are the only Jordan automorphisms of $\mathcal{T}_r(F)$, where F is a field containing at least three elements. This result was generalized by Beidar, Brešar and Chebotar [4] who proved that every Jordan isomorphism of $\mathcal{T}_r(C)$ onto an arbitrary algebra over C is either an isomorphism or an anti-isomorphism, provided that C is a unital 2-torsionfree commutative ring whose only idempotents are 0 and 1. Further, recently Lu [13] proved that every Jordan isomorphism between nest algebras is either an isomorphism or an anti-isomorphism. Our aim in this paper is to unify and generalize these results. In Section 2 we shall introduce the concept of an indecomposable triangular ring and give some examples of such rings; in particular, algebras $\mathcal{T}_r(C)$ from [4] and nontrivial nest algebras are such examples. In Section 3 we shall prove our main result stating that every Jordan isomorphism from a

Received by the editors June 29, 2004.

2000 *Mathematics Subject Classification*. Primary 47L35; Secondary 16S50.

Key words and phrases. Jordan isomorphisms, triangular rings, triangular matrix algebras, nest algebras.

This research was supported by NSC Grants NSC 91-2115-M-110-005.

©2005 American Mathematical Society
Reverts to public domain 28 years from publication

2-torsionfree unital indecomposable triangular ring onto another ring is either an isomorphism or an anti-isomorphism.

2. INDECOMPOSABLE TRIANGULAR RINGS

We fix some notation first. Let A and B be rings and let M be an (A, B) -bimodule which is faithful both as a left A -module and as a right B -module. Note that

$$Tri(A, B, M) = \left\{ \begin{pmatrix} a & m \\ 0 & b \end{pmatrix} \mid a \in A, b \in B, m \in M \right\}$$

is a ring under the usual matrix operations. Following Cheung [7] we shall call $Tri(A, B, M)$ a *triangular ring*.

We shall consider $A, B,$ and M as subsets of $\mathcal{T} = Tri(A, B, M)$, i.e. we shall identify them by their copies inside \mathcal{T} . Note that $AB = BA = MA = BM = M^2 = 0$.

We recall that a ring R is said to be unital if it contains unity, which will be denoted by 1_R , and to be 2-torsionfree if it does not contain a nonzero element a such that $2a = 0$. Suppose that \mathcal{T} is a unital ring. Note that $1_{\mathcal{T}}$ is necessarily of the form

$$1_{\mathcal{T}} = \begin{pmatrix} 1_A & 0 \\ 0 & 1_B \end{pmatrix}$$

where unities 1_A and 1_B of A and B also satisfy $1_A m = m = m 1_B$ for all $m \in M$.

In what follows, we write submodule in short for (A, B) -subbimodule of M . We shall say that a ring R is an *indecomposable triangular ring* if it is isomorphic to a triangular ring $Tri(A, B, M)$ such that M cannot be written as a direct sum of two nonzero submodules. We will give two examples of such rings in Theorems 2.1 and 2.7.

Theorem 2.1. *Suppose that R is a unital ring which contains no nontrivial central idempotents. Then $\mathcal{T}_r(R)$ is an indecomposable triangular ring for every $r \geq 2$.*

Proof. Pick any positive integers s, t such that $r = s + t$. Set $A = \mathcal{T}_s(R), B = \mathcal{T}_t(R)$, and $M = M_{s \times t}(R)$. Note that $\mathcal{T}_r(R) \cong Tri(A, B, M)$.

The matrix units of A, B and M will be denoted by e'_{ij}, e''_{ij} and e_{ij} respectively. Suppose $M = P \oplus Q$, where P, Q are submodules of M . Write $e_{s1} = p + q$ where $p \in P$ and $q \in Q$. Then $p + q = e_{s1} = e'_{ss} e_{s1} e''_{11} = e'_{ss} (p + q) e''_{11} = e'_{ss} p e''_{11} + e'_{ss} q e''_{11}$. Since $e'_{ss} \in A$ and $e''_{11} \in B$ we have $p - e'_{ss} p e''_{11} = -q + e'_{ss} q e''_{11} \in P \cap Q = 0$ and so $p = e'_{ss} p e''_{11} = \alpha e_{s1}$ and $q = e'_{ss} q e''_{11} = \beta e_{s1}$ for some $\alpha, \beta \in R$ such that $\alpha + \beta = 1_R$. Let $x \in R$. Note that, on the one hand,

$$\begin{aligned} x e_{s1} &= (x e'_{ss}) e_{s1} = x e'_{ss} (\alpha e_{s1} + \beta e_{s1}) \\ &= x \alpha e_{s1} + x \beta e_{s1} \end{aligned}$$

and, on the other hand,

$$x e_{s1} = e_{s1} (x e''_{11}) = (\alpha e_{s1} + \beta e_{s1}) x e''_{11} = \alpha x e_{s1} + \beta x e_{s1}.$$

Since $x e'_{ss} \in A$ and $x e''_{11} \in B$, we have $x \alpha = \alpha x$ and $x \beta = \beta x$ for all $x \in R$, i.e. α, β lie in the center of R . If $r \in \alpha R \cap \beta R$, then $r e_{s1} \in P \cap Q = 0$ and so $r = 0$. Therefore $R = \alpha R \oplus \beta R$ as ideals. From $\alpha \beta, \beta \alpha \in \alpha R \cap \beta R = 0$ and $1 = \alpha + \beta = (\alpha + \beta)^2 = \alpha^2 + \beta^2$ it follows that $\alpha^2 = \alpha$ and $\beta^2 = \beta$ and so α, β are central idempotents of R . By the assumption, we have either $\alpha = 0$ or $\beta = 0$, say $\beta = 0$, and so $e_{s1} \in P$. Let $x \in R, 1 \leq i \leq s, 1 \leq j \leq t$. Since $x e'_{is} \in A$

and $e''_{1j} \in B$, we have $xe_{ij} = xe'_{is}e_{s1}e''_{1j} \in P$ and so $P = M$. Thus, $\mathcal{T}_r(R)$ is an indecomposable triangular ring. \square

A nest \mathcal{N} is a totally ordered set of closed subspaces of a Hilbert space H such that $0, H \in \mathcal{N}$ and \mathcal{N} is closed under arbitrary intersections and closed linear spans of its elements. By $\mathcal{B}(H)$ we denote the algebra of all bounded linear operators on H . The nest algebra $\mathcal{T}(\mathcal{N})$ associated to \mathcal{N} is a subalgebra of $\mathcal{B}(H)$ consisting of those operators that leave N invariant for every $N \in \mathcal{N}$, i.e. $\mathcal{T}(\mathcal{N}) = \{T \in \mathcal{B}(H) \mid TN \subseteq N \text{ for all } N \in \mathcal{N}\}$. We follow [8] for the following notation and definitions. Let $N \in \mathcal{N}$. By N^\perp we denote the orthogonal complement of N , by E_N we denote the orthogonal projection of H onto N and by $x \otimes y^*$, for elements $x, y \in H$, we denote the rank one operator $w \mapsto \langle w, y \rangle x$ for all $w \in H$. Further, we define $N_- = \sup\{N' \in \mathcal{N} \mid N' \subset N\}$, $N_+ = \inf\{N' \in \mathcal{N} \mid N \subset N'\}$ and $N' \ominus N = N' \cap N^\perp$ for $N \subseteq N' \in \mathcal{N}$.

Suppose that \mathcal{N} is nontrivial, i.e. \mathcal{N} contains other spaces than 0 and H . The associated nest algebra $\mathcal{T}(\mathcal{N})$ is then called nontrivial. We now fix $N_0 \in \mathcal{N}$ in the following way: $N_0 = 0_+$ if $0_+ \neq 0$, and if $0_+ = 0$, then let N_0 be just any element in \mathcal{N} different from $0, H$. We denote E_{N_0} by E ; hence $E_{N_0^\perp} = 1 - E$. Let $A = \{T \in \mathcal{B}(H) \mid T = ETE\}$, $B = \{T \in \mathcal{B}(H) \mid T = (1 - E)T(1 - E)\}$ and $M = \{T \in \mathcal{B}(H) \mid T = ET(1 - E)\}$. Let $T \in \mathcal{T}(\mathcal{N})$. Note that $(1 - E)TE = 0$. Consider the map $T \mapsto \begin{pmatrix} ETE & ET(1 - E) \\ 0 & (1 - E)T(1 - E) \end{pmatrix}$. We can see that it is an isomorphism and so $\mathcal{T}(\mathcal{N}) \cong Tri(A, B, M)$. We remark that this was noted, in a somewhat different form, in [7, Proposition 5].

We remark that $x \otimes y^* \in \mathcal{T}(\mathcal{N})$ if there exists $N \in \mathcal{N}$ such that $x \in N$ and $y \in N^\perp$ (see [8, Lemma 2.8] or [15, Lemma 3.3]). In what follows, N always denotes some element of \mathcal{N} . According to the definitions of A and B , we have

- Lemma 2.2.** (i) *If $N \subseteq N_0$, then $x \otimes y^* \in A$ for all $x \in N, y \in N_0 \ominus N_-$.*
- (ii) *If $N \supseteq N_0$, then $x \otimes y^* \in B$ for all $x \in N \ominus N_0, y \in N^\perp$.*

Lemma 2.3. *Suppose $0_+ \neq 0$. Then $x \otimes y^* \in A$ for all $x, y \in N_0$.*

Proof. Note that $(0_+)_- = 0$ if $0_+ \neq 0$. Since $N_0 = 0_+$ in this case, $(N_0)_- = 0$. By substituting N by $N_0 = 0_+$ in Lemma 2.2(i), we have the desired result. \square

Lemma 2.4. *Suppose $0_+ = 0$. If $0 \neq x \in N_0$, then there exists $0 \neq N \in \mathcal{N}$, $N \subset N_0$ such that $x \notin N$.*

Proof. Since $x \neq 0$ and $0_+ = 0$, we have $x \notin 0_+ = \inf\{N \in \mathcal{N} \mid N \neq 0\}$ and so the desired result follows. \square

Lemma 2.5. *Let L be a nonzero submodule of M . Then there exist rank one elements in L .*

Proof. Let $0 \neq T \in L$. Then there exists $v \in N_0^\perp$ such that $0 \neq u = Tv \in N_0$. If $0_+ \neq 0$, then by Lemma 2.3 we have $u \otimes u^* \in A$ and so $0 \neq (u \otimes u^*)T \in L$ which is a rank one element. If $0_+ = 0$, then by Lemma 2.4 we have $u \notin N$ for some $0 \neq N \in \mathcal{N}$ and $N \subset N_0$. Let $x = E_{N^\perp}u \neq 0$ and choose $0 \neq w \in N$. From Lemma 2.2(i), we have $w \otimes x^* \in A$, and so $0 \neq (w \otimes x^*)T \in L$ which is an element of rank one. \square

Lemma 2.6. *Let L be a submodule of M and $0 \neq u \otimes v^* \in L$ for some $u \in N_0$, $v \in N_0^\perp$.*

- (i) *If $u \notin N$ for some $N \subset N_0$, then $w \otimes v^* \in L$ for all $w \in N$.*
- (ii) *If $v \in N$ for some $N \supset N_0$, then $u \otimes w^* \in L$ for all $w \in N^\perp$.*
- (iii) *$u \otimes (E_{N^\perp} v)^* \in L$ for all $N \supseteq N_0$.*

Proof. (i) Let $x = E_{N^\perp} u$. Note that $x \neq 0$. Since $w \otimes x^* \in A$ by Lemma 2.2(i), we have $(w \otimes x^*)(u \otimes v^*) = \|x\|^2(w \otimes v^*) \in L$ and so $w \otimes v^* \in L$. (ii) By Lemma 2.2(ii) we have $v \otimes w^* \in B$ and so $(u \otimes v^*)(v \otimes w^*) = \|v\|^2(u \otimes w^*) \in L$, which in turn implies $u \otimes w^* \in L$. (iii) Let $x = E_{N^\perp} v$. Our claim trivially holds if $x = v$. In case $x \neq v$, then $y = E_N v \neq 0$. It follows from Lemma 2.2(ii) that $y \otimes x^* \in B$ and so $(u \otimes v^*)(y \otimes x^*) = \|y\|^2(u \otimes x^*) \in L$, which in turn implies $u \otimes x^* \in L$. \square

Theorem 2.7. *Nontrivial nest algebras are indecomposable triangular rings.*

Proof. We have $\mathcal{T}(\mathcal{N}) \cong \text{Tri}(A, B, M)$ where A, B, M are as above. Suppose that M can be decomposed into a direct sum of two nonzero submodules P and Q . We show that this will lead to a contradiction. First, for $w \in H$, we define N_w as the least element of \mathcal{N} containing w , i.e. $N_w = \inf\{N \in \mathcal{N} \mid w \in N\}$. Suppose $0 \neq u \otimes v^* \in P$ and $0 \neq u' \otimes v'^* \in Q$ for some $u, u' \in N_0$ and $v, v' \in N_0^\perp$. If $0_+ \neq 0$, then by Lemma 2.3 we have $u \otimes u'^* \in A$ and so $(u \otimes u'^*)(u' \otimes v'^*) = \|u'\|^2(u \otimes v'^*) \in Q$, which in turn implies $u \otimes v'^* \in Q$. If $0_+ = 0$ and if $N_u = N_{u'}$, then by Lemma 2.4 we have $u \notin N$ for some $0 \neq N \in \mathcal{N}$ and $N \subset N_0$. Choose $0 \neq w \in N$. By Lemma 2.6(i) we have $0 \neq w \otimes v^* \in P$ and $N_w \subseteq N \neq N_u = N_{u'}$. That is, we have that if $0 \neq u \otimes v^* \in P$ and $0 \neq u' \otimes v'^* \in Q$, then

$$(1) \quad u \otimes v'^* \in Q \text{ in case } 0_+ \neq 0;$$

and

$$(2) \quad \text{there exists } 0 \neq w \otimes v^* \in P \text{ such that } N_w \neq N_{u'} \text{ in case } 0_+ = 0.$$

Next, suppose $u = u'$ and $N_v \neq N_{v'}$, say $N_v \subset N_{v'}$. Note that $v' \notin N_v$ and so $E_{N_v^\perp} v' \neq 0$. It follows from Lemma 2.6(ii) and (iii) that $0 \neq u \otimes (E_{N_v^\perp} v')^* \in P \cap Q$, a contradiction. Therefore, we have

$$(3) \quad \text{if } 0 \neq u \otimes v^* \in P \text{ and } 0 \neq u \otimes v'^* \in Q, \text{ then } N_v = N_{v'}.$$

Suppose $N_u \neq N_{u'}$, say $N_u \subset N_{u'}$. Then by Lemma 2.6(i) we have $0 \neq u \otimes v'^* \in Q$. By the result in (3), we have $N_v = N_{v'}$. That is, we have

$$(4) \quad \text{if } 0 \neq u \otimes v^* \in P \text{ and } 0 \neq u' \otimes v'^* \in Q, \text{ then either } N_u = N_{u'} \text{ or } N_v = N_{v'}.$$

In case $0_+ \neq 0$, by (1) we have $0 \neq u \otimes v'^* \in Q$ and so $N_v = N_{v'}$ by (3). Suppose $0_+ = 0$. If $N_u \neq N_{u'}$, then by (4) we have $N_v = N_{v'}$. If $N_u = N_{u'}$, then by (2) there exists $0 \neq w \otimes v^* \in P$ and $N_w \neq N_{u'}$ and so, by (4) again, we have $N_v = N_{v'}$. Therefore, it follows that

$$(5) \quad \text{if } 0 \neq u \otimes v^* \in P \text{ and } 0 \neq u' \otimes v'^* \in Q, \text{ then } N_v = N_{v'}.$$

Suppose that $N_v \neq H$. Choose $0 \neq w \in N_v^\perp$. Since by Lemma 2.6(ii) we have $0 \neq u \otimes w^* \in P$, then by (5) we have $N_w = N_{v'}$. However, it follows from $w \notin N_v$ that $N_w \neq N_v = N_{v'}$, a contradiction. Therefore, we have

$$(6) \quad \text{if } 0 \neq u \otimes v^* \in P, \text{ then } N_v = H.$$

Suppose that $v \notin N^\perp$ for some $N_0 \subseteq N \neq H$. Let $x = E_N v \neq 0$ and $y = E_{N^\perp} v$. From Lemma 2.6(iii) it follows that $u \otimes y^* \in P$ and so $0 \neq u \otimes x^* = u \otimes v^* - u \otimes y^* \in P$.

By (6), we have $N_x = H$, but $N_x \subseteq N \neq H$, a contradiction. Therefore $v \in H^\perp$. By Lemma 2.6(ii), we have $u' \otimes v^* \in Q$. Suppose $N_u \neq N_{u'}$, say $N_u \subset N_{u'}$. Since $u' \notin N_u$, from Lemma 2.6(i) it follows that $0 \neq u \otimes v^* \in P \cap Q$, a contradiction. Therefore we have

$$(7) \quad \text{if } 0 \neq u \otimes v^* \in P \text{ and } 0 \neq u' \otimes v'^* \in Q, \text{ then } u' \otimes v^* \in Q \text{ and } N_u = N_{u'}.$$

If $0_+ = 0$, then by (2) there exist $0 \neq w \otimes v^* \in P$ such that $N_w \neq N_{w'}$; this contradicts (7) and so we have

$$(8) \quad 0_+ \neq 0.$$

Now by Lemma 2.5, there exist $0 \neq u \otimes v^* \in P$ and $0 \neq u' \otimes v'^* \in Q$ for some $u, u' \in N_0$ and $v, v' \in N_0^\perp$. By (7) we have $u' \otimes v^* \in Q$. Now, by (8), $0_+ \neq 0$ and so it follows from (1) that $0 \neq u \otimes v^* \in P \cap Q$, a contradiction. This completes the proof. \square

3. JORDAN ISOMORPHISMS

Theorem 3.1. *Let \mathcal{T} be a 2-torsionfree unital indecomposable triangular ring. Then every Jordan isomorphism from \mathcal{T} onto another ring is either an isomorphism or an anti-isomorphism.*

Proof. In the proof we use some ideas from [4].

We may assume that $\mathcal{T} = \text{Tri}(A, B, M)$ where M cannot be decomposed. Let φ be a Jordan isomorphism from \mathcal{T} onto a ring \mathcal{S} . Since \mathcal{T} is 2-torsionfree, so is \mathcal{S} , and so φ clearly satisfies $\varphi(x^2) = \varphi(x)^2$ for all $x \in \mathcal{T}$. We denote $xy + yx$ by $x \circ y$. Since $2xyx = x \circ (x \circ y) - x^2 \circ y$, we see that φ also satisfies $\varphi(xy x) = \varphi(x)\varphi(y)\varphi(x)$ for all $x, y \in \mathcal{T}$. This obviously yields $\varphi(xyz + zyx) = \varphi(x)\varphi(y)\varphi(z) + \varphi(z)\varphi(y)\varphi(x)$ for all $x, y, z \in \mathcal{T}$. In what follows we shall often use these identities without explicit mention.

Let $e = \varphi(1_A)$ and $f = \varphi(1_B)$. Obviously, e and f are idempotents in \mathcal{S} . From $e \circ f = \varphi(1_A \circ 1_B) = 0 = \varphi(1_A 1_B 1_A) = e f e$ it follows that $e f = 0 = f e$. Further, let $a \in A$, $\varphi(a) = \varphi(1_A a 1_A) = e \varphi(a) e$ and $2\varphi(a) = \varphi(1_A \circ a) = e \circ \varphi(a)$ imply that $\varphi(a) = e \varphi(a) = \varphi(a) e$, i.e. $e = 1_{\varphi(A)}$. Similarly we see that $f = 1_{\varphi(B)}$ and $e + f = \varphi(1_{\mathcal{T}}) = 1_{\mathcal{S}}$.

Let $a \in A, b \in B, m \in M$. It follows from $1_A b 1_A = 0 = 1_A m 1_A$ that $e \varphi(b) e = 0 = e \varphi(m) e$. Similarly, $f \varphi(a) f = 0 = f \varphi(m) f$. Therefore $\varphi(A) = e \varphi(\mathcal{T}) e = e \mathcal{S} e$ and $\varphi(B) = f \varphi(\mathcal{T}) f = f \mathcal{S} f$ are subrings of \mathcal{S} . Since $\varphi(A)\varphi(B) = 0 = \varphi(B)\varphi(A)$, we have

$$\varphi(a)\varphi(m)\varphi(b) = \varphi(a)(\varphi(m) \circ \varphi(b)) = \varphi(a)\varphi(m \circ b) = \varphi(a)\varphi(mb)$$

and

$$\varphi(b)\varphi(m)\varphi(a) = \varphi(b)(\varphi(a) \circ \varphi(m)) = \varphi(b)\varphi(a \circ m) = \varphi(b)\varphi(am).$$

Comparing $\varphi(amb) = \varphi(amb + bma) = \varphi(a)\varphi(m)\varphi(b) + \varphi(b)\varphi(m)\varphi(a) = \varphi(a)\varphi(mb) + \varphi(b)\varphi(am)$ with $\varphi(amb) = \varphi(am \circ b) = \varphi(am) \circ \varphi(b) = \varphi(am)\varphi(b) + \varphi(b)\varphi(am)$, we have $\varphi(a)\varphi(mb) = \varphi(am)\varphi(b)$. Further, since $a \circ mb = amb = am \circ b$, we have $\varphi(mb)\varphi(a) = \varphi(b)\varphi(am)$, that is,

$$\begin{aligned} \varphi(a)\varphi(mb) &= \varphi(am)\varphi(b), \\ \varphi(mb)\varphi(a) &= \varphi(b)\varphi(am) \end{aligned}$$

for all $a \in A, b \in B, m \in M$. In particular, we have

$$(9) \quad \begin{aligned} \varphi(a)\varphi(m) &= \varphi(am)f & , & \quad \varphi(m)\varphi(a) = f\varphi(am), \\ \varphi(m)\varphi(b) &= e\varphi(mb) & , & \quad \varphi(b)\varphi(m) = \varphi(mb)e, \\ e\varphi(m) &= \varphi(m)f & , & \quad \varphi(m)e = f\varphi(m) \end{aligned}$$

for all $a \in A, b \in B, m \in M$.

We claim first $e\varphi(M) \subseteq \varphi(M)$. Let $m \in M$ and write $e\varphi(m) = \varphi(a) + \varphi(b) + \varphi(n)$ for some $a \in A, b \in B, n \in M$. Multiplying this relation from the right by e and from the left by f separately and using $e\varphi(m)e = 0, ef = 0 = fe$, we have $0 = \varphi(a) + \varphi(n)e$ and $0 = \varphi(b) + f\varphi(n)$. Since, by (9), $\varphi(n)e = f\varphi(n)$, we have $\varphi(a) = \varphi(b) = 0$ and so $e\varphi(m) = \varphi(n) \in \varphi(M)$, as desired. Therefore, we have $e\varphi(M) \subseteq \varphi(M)$ and $f\varphi(M) = (1_S - e)\varphi(M) \subseteq \varphi(M)$ and so $\varphi^{-1}(e\varphi(M)) \subseteq M$ and $\varphi^{-1}(f\varphi(M)) \subseteq M$.

We claim next that M is a direct sum of $\varphi^{-1}(e\varphi(M))$ and $\varphi^{-1}(f\varphi(M))$, which are both submodules. Let $x \in \varphi^{-1}(e\varphi(M)) \subseteq M$. Therefore $xa = 0$ and it follows from (9) that $\varphi(x)\varphi(a) \subseteq e\varphi(M)\varphi(A) \subseteq ef\varphi(M) = 0$ and $\varphi(a)\varphi(x) \in e\varphi(M)$. Therefore $\varphi(ax) = \varphi(a \circ x) = \varphi(a) \circ \varphi(x) = \varphi(a)\varphi(x) \in e\varphi(M)$, i.e. $ax \in \varphi^{-1}(e\varphi(M))$, and so $\varphi^{-1}(e\varphi(M))$ is a left A -submodule of M . Similarly, $\varphi^{-1}(e\varphi(M))$ is a right B -submodule and $\varphi^{-1}(f\varphi(M))$ is a submodule. Therefore $M = \varphi^{-1}(e\varphi(M)) \oplus \varphi^{-1}(f\varphi(M))$ as submodules. According to our assumption we have either $f\varphi(M) = 0$ or $e\varphi(M) = 0$.

Case 1: Suppose $f\varphi(M) = 0$. Then $\varphi(m) = e\varphi(m) = \varphi(m)f$ for all $m \in M$, and so $\varphi(M)\varphi(M) = \varphi(M)f\varphi(M) = 0$ and $\varphi(M)\varphi(A) = 0 = \varphi(B)\varphi(M)$. Hence, by (9)

$$\begin{aligned} \varphi(a)\varphi(m) &= \varphi(am)f = \varphi(am), \\ \varphi(m)\varphi(b) &= e\varphi(mb) = \varphi(mb), \end{aligned}$$

and so, since M is faithful both as a left A -module and as a right B -module, we see that $\varphi(M)$ is faithful as a left $\varphi(A)$ -module as well as a right $\varphi(B)$ -module. Hence, it follows from $\varphi(aa')\varphi(m) = \varphi((aa')m) = \varphi(a(a'm)) = \varphi(a)\varphi(a'm) = \varphi(a)\varphi(a')\varphi(m)$ and $\varphi(m)\varphi(bb') = \varphi(m(bb')) = \varphi((mb)b') = \varphi(mb)\varphi(b') = \varphi(m)\varphi(b)\varphi(b')$ that $\varphi(aa') = \varphi(a)\varphi(a')$ and $\varphi(bb') = \varphi(b)\varphi(b')$. Therefore,

$$\begin{aligned} \varphi((a+b+m)(a'+b'+m')) &= \varphi(aa' + bb' + am' + mb') \\ &= \varphi(aa') + \varphi(bb') + \varphi(am') + \varphi(mb') \\ &= \varphi(a)\varphi(a') + \varphi(b)\varphi(b') + \varphi(a)\varphi(m') + \varphi(m)\varphi(b') \\ &= (\varphi(a) + \varphi(b) + \varphi(m))(\varphi(a') + \varphi(b') + \varphi(m')) \\ &= \varphi(a+b+m)\varphi(a'+b'+m'); \end{aligned}$$

that is, φ is an isomorphism.

Case 2: Suppose $e\varphi(M) = 0$. Then $\varphi(m) = f\varphi(m) = \varphi(m)e$ for all $m \in M$, and so

$$(10) \quad \varphi(M)\varphi(M) = 0 = \varphi(A)\varphi(M) = \varphi(M)\varphi(B).$$

By (9), we have

$$(11) \quad \begin{aligned} \varphi(m)\varphi(a) &= f\varphi(am) = \varphi(am), \\ \varphi(b)\varphi(m) &= \varphi(mb)e = \varphi(mb), \end{aligned}$$

and so, $\varphi(M)$ is faithful as a left $\varphi(B)$ -module as well as a right $\varphi(A)$ -module. Therefore, it follows from $\varphi(m)\varphi(aa') = \varphi((aa')m) = \varphi(a(a'm)) = \varphi(a'm)\varphi(a) = \varphi(m)\varphi(a')\varphi(a)$ and $\varphi(bb')\varphi(m) = \varphi(m(bb')) = \varphi((mb)b') = \varphi(b')\varphi(mb) = \varphi(b')\varphi(b)\varphi(m)$ that $\varphi(aa') = \varphi(a')\varphi(a)$ and $\varphi(bb') = \varphi(b')\varphi(b)$; together with (10) and (11) we have that φ is an anti-isomorphism. \square

Now we have the following two immediate results which generalize [4, Theorem] and [13, Theorem 15] respectively.

Theorem 3.2. *Let R be a 2-torsionfree unital ring and let $r \geq 2$ be an integer. Then R contains no nontrivial central idempotents if and only if every Jordan isomorphism of $\mathcal{T}_r(R)$ onto any ring is either an isomorphism or an anti-isomorphism.*

Proof. If R contains no nontrivial central idempotents, then the desired result follows from Theorem 2.1 and Theorem 3.1. Suppose that R contains a nontrivial central idempotent ε . Then the map $A \mapsto \varepsilon A + (1 - \varepsilon)UA^{\text{tr}}U$ (cf. [4, 14]), where A^{tr} denotes the transpose of A and $U = e_{1r} + e_{2r-1} + \dots + e_{r-12} + e_{r1}$, is a Jordan automorphism of $\mathcal{T}_r(R)$ which is neither an automorphism nor an anti-automorphism. This completes the proof. \square

Theorem 3.3. *Every Jordan isomorphism from a nest algebra onto another complex algebra is either an isomorphism or an anti-isomorphism.*

Proof. Let $\mathcal{T}(\mathcal{N})$ be a nest algebra. If \mathcal{N} is trivial, then $\mathcal{T}(\mathcal{N}) = \mathcal{B}(H)$ is a prime ring and so we have the desired conclusion by Herstein's result [9]. If \mathcal{N} is nontrivial, then we apply Theorems 2.7 and 3.1. \square

We remark that using Theorem 3.3 and the facts that every isomorphism between nest algebras is spatial [16, Theorem 4.2] (see also [8, Corollary 17.13]) and the composition of an anti-isomorphism and $*$ (adjoint) is an isomorphism, we can get [13, Theorem 15], which states that every Jordan isomorphism between nest algebras is of the form $T \mapsto STS^{-1}$ or $T \mapsto ST^*S^{-1}$ for some invertible operator S .

ACKNOWLEDGEMENT

I wish to express my sincere thanks to Professor M. Brešar for suggesting this topic to me and for giving me much helpful advice.

REFERENCES

[1] G. Ancochea, Le théorème de von Staundt en géométrie projective quaternionienne, *J. Reine Angew. Math.* **184** (1942), 192–198. MR0008893 (5:72d)
 [2] G. Ancochea, On semi-automorphisms of division algebras, *Ann. Math.* **48** (1947), 147–153. MR0018642 (8:310c)
 [3] W.E. Baxter and W.S. Martindale 3rd, Jordan homomorphisms of semiprime rings, *J. Algebra* **56** (1979), 457–471. MR0528587 (80f:16008)
 [4] K.I. Beidar, M. Brešar, M.A. Chebotar, Jordan isomorphisms of triangular matrix algebras over a connected commutative ring, *Linear Algebra Appl.* **312** (2000), 197–201. MR1759333 (2001a:16048)
 [5] M. Brešar, Jordan mappings of semiprime rings, *J. Algebra* **127** (1989), 218–228. MR1029414 (91a:16025)
 [6] M. Brešar, Jordan mappings of semiprime rings II, *Bull. Austral. Math. Soc.* **44** (1991), 233–238. MR1126361 (92j:16024)
 [7] W.-S. Cheung, Commuting maps of triangular algebras, *J. London Math. Soc.* **63** (2001), 117–127. MR1802761 (2001k:16053)

- [8] K.R. Davidson, Nest algebras, Pitman Research Notes in Mathematics, Vol. 191, Longman, London/New York, 1988. MR0972978 (90f:47062)
- [9] I.N. Herstein, Jordan homomorphisms, *Trans. Amer. Math. Soc.* **81** (1956), 331–351. MR0076751 (17,938f)
- [10] L.-K. Hua, On the automorphisms of a s-field, *Proc. Nat. Acad. Sci. U.S.A.* **35** (1949), 386–389. MR0029886 (10:675d)
- [11] N. Jacobson and C. Rickart, Jordan homomorphisms of rings, *Trans. Amer. Math. Soc.* **69** (1950), 479–502. MR0038335 (12:387h)
- [12] I. Kaplansky, Semi-automorphisms of rings, *Duke Math. J.* **14** (1947), 521–527. MR0022209 (9:172e)
- [13] F. Lu, Jordan isomorphisms of nest algebras, *Proc. Amer. Math. Soc.* **131** (2002), 147–154. MR1929034 (2003f:47121)
- [14] L. Molnár, P. Šemrl, Some linear preserver problems on upper triangular matrices, *Linear and Multilinear Algebra* **45** (1998), 189–206. MR1671619 (99h:15003)
- [15] J.R. Ringrose, On some algebras of operators, *Proc. London Math. Soc.* **15** (1965), 61–83. MR0171174 (30:1405)
- [16] J.R. Ringrose, On some algebras of operators II, *Proc. London Math. Soc.* **16** (1966), 385–402. MR0196516 (33:4703)
- [17] M.F. Smiley, Jordan homomorphisms onto prime rings, *Trans. Amer. Math. Soc.* **84** (1957), 426–429. MR0083484 (18:715b)

DEPARTMENT OF APPLIED MATHEMATICS, NATIONAL SUN YAT-SEN UNIVERSITY, KAOHSIUNG, TAIWAN, 804

E-mail address: `tlwong@math.nsysu.edu.tw`