

CONTINUOUS SELECTIONS AND FIXED POINTS
OF MULTI-VALUED MAPPINGS
ON NONCOMPACT OR NONMETRIZABLE SPACES

LAI-JIU LIN, NGAI-CHING WONG, AND ZENN-TSUEN YU

(Communicated by Jonathan M. Borwein)

ABSTRACT. In this paper, we obtain several new continuous selection theorems for multi-valued mappings on completely regular spaces and fixed point theorems for multi-valued maps on nonmetrizable spaces. They, in particular, provide a partial solution of a conjecture of X. Wu.

1. INTRODUCTION

In [4, 5], Browder first used a continuous selection theorem to prove the Fan-Browder fixed point theorem. Later, Yannelis and N. D. Prabhakar [17], Ben-El-Mechaiekh [2, 3], Ding, Kim and Tan [8], Horvath [11], Wu [16, 15], Park [12, 13], and many others, established several continuous selection theorems with applications. We note that in all the continuous selection theorems studied by the above authors, the multi-valued maps are defined on a compact or paracompact space. In [18], Yu and Lin studied continuous selections of multi-valued mappings defined on noncompact spaces, but they assume some kind of coercivity conditions instead.

In this paper, we establish a continuous selection theorem for a multi-valued map defined on a completely regular topological space. We do not assume the compactness of its domain.

In the second part of this paper, we discuss collectively fixed points of lower semicontinuous multi-valued maps. Recently, many authors studied fixed point theorems of lower semicontinuous multi-valued maps; see for example [14, 6, 15, 1]. In particular, Wu established the following one.

Theorem 1.1 ([15]). *Let X be a nonempty subset of a Hausdorff locally convex topological vector space, let D be a nonempty compact metrizable subset of X , and let $T : X \rightarrow 2^D$ be a multi-valued mapping with the following properties:*

- (a) $T(x)$ is a nonempty closed convex set for each x in X ;
- (b) T is lower semicontinuous.

Then there exists a point \bar{x} in D such that $\bar{x} \in T(\bar{x})$.

Wu conjectured in [15] that the conclusion of Theorem 1.1 remains true even if the metrizability condition of D is dropped. In this paper, we shall use the

Received by the editors July 17, 2003.

2000 *Mathematics Subject Classification*. Primary 54C65, 46H10, 54H25.

Key words and phrases. Multi-valued mappings, continuous selections, fixed points.

approximate continuous selection theorem of Deutsch and Kenderov [7] (see also [20]) to establish an approximate fixed point theorem for a sub-lower semicontinuous multi-valued map. This gives rise to a partial solution of the conjecture of Wu [15]. We shall also provide a simple proof of a Himmelberg type collectively fixed point theorem. We remark that our results differ from the approximate fixed point theorem recently established by Park [13].

We would like to thank the referee for many helpful suggestions on improving the presentation and the bibliography in this paper.

2. PRELIMINARIES

Let X and Y be topological spaces. A multi-valued map $T : X \rightarrow 2^Y$ is a map from X into the power set 2^Y of Y . Let $T^{-1} : Y \rightarrow 2^X$ be defined by the condition that $x \in T^{-1}y$ if and only if $y \in T(x)$. Recall that

- (a) T is said to be *closed* if its graph $G_r(T) = \{(x, y) : x \in X, y \in T(x)\}$ is closed in the product space $X \times Y$;
- (b) T is said to be *upper semicontinuous* (in short, u.s.c.) at x if for every open set V in Y with $T(x) \subset V$, there exists a neighborhood $W(x)$ of x such that $T(W(x)) \subset V$; T is said to be u.s.c. on X if T is u.s.c. at every point of X ;
- (c) T is said to be *lower semicontinuous* (in short, l.s.c.) at x if for every open neighborhood $V(y)$ of every y in $T(x)$, there exists a neighborhood $W(x)$ of x such that $T(u) \cap V(y) \neq \emptyset$ for all u in $W(x)$; T is said to be l.s.c. on X if T is l.s.c. at every point of X ;
- (d) in case Y is a topological linear space, T is said to be *sub-lower semicontinuous* (see, e.g., [20]) at an x in X if for each neighborhood V of 0 in Y , there is a z in $T(x)$ and a neighborhood $U(x)$ of x in X such that $z \in T(y) + V$ for each y in $U(x)$; T is said to be sub-lower semicontinuous on X if T is sub-lower semicontinuous at every point of X . It is plain that if T is lower semicontinuous at x , then T is sub-lower semicontinuous at x .

The following lemmas are needed in this paper.

Lemma 2.1 (Deutsch and Kenderov [7]). *Let X be a paracompact topological space, let Y be a locally convex topological linear space, and let $F : X \rightarrow 2^Y$. Then F is sub-lower semicontinuous if and only if for each neighborhood V of 0 in Y , there is a continuous function $f : X \rightarrow Y$ such that $f(x) \in F(x) + V$ for each x in X .*

Lemma 2.2 (Yuan [19]). *Let X be a topological space, let Y be a nonempty subset of a topological vector space with a base \mathcal{B} for the zero neighborhoods, and let $F : X \rightarrow 2^Y$. For each V in \mathcal{B} , define $F_V : X \rightarrow 2^Y$ by*

$$F_V(x) = (F(x) + V) \cap Y, \quad \forall x \in X.$$

Write $\bar{y} \in \overline{F}(\bar{x})$ if $(\bar{x}, \bar{y}) \in \overline{G_r F}$. Then for any \bar{x} in X and \bar{y} in Y , we have

$$\bar{y} \in \overline{F}(\bar{x}) \quad \text{whenever} \quad \bar{y} \in \bigcap_{V \in \mathcal{B}} \overline{F_V}(\bar{x}).$$

Lemma 2.3 (Himmelberg [10]). *Let X be a nonempty convex subset of a locally convex topological vector space. Let $T : X \rightarrow 2^X$ be a u.s.c. multi-valued map*

with nonempty closed convex values such that $T(X) = \bigcup_{x \in X} T(x)$ is contained in a compact subset of X . Then there exists an \bar{x} in X such that $\bar{x} \in T(\bar{x})$.

Lemma 2.4 (Granas [9]; see also Ding, Kim and Tan [8]). *Let D be a nonempty compact subset of a topological vector space. Then the convex hull $\text{co}D$ of D is σ -compact and hence is paracompact.*

3. CONTINUOUS SELECTION THEOREMS

Note that the set $S^{-1}(y) = \{x \in X : y \in S(x)\}$ below can have empty interior for some y in K .

Theorem 3.1. *Let X be a completely regular space and let K be a nonempty subset of a Hausdorff topological vector space E . Assume a multi-valued function $S : X \rightarrow 2^K$ satisfies the following conditions:*

- (a) *For each x in X , the set $S(x)$ is convex.*
- (b) $X = \bigcup \{\text{int } S^{-1}(y) : y \in K\}$.

Then for any compact subset F of X there is an open dense subset U of X containing F such that S has a continuous selection $f : U \rightarrow K$, that is, $f(x) \in S(x)$ for all x in U .

Proof. By assumption (b), there are finitely many points y_1, \dots, y_n in K such that

$$F \subseteq \text{int } S^{-1}(y_1) \cup \dots \cup \text{int } S^{-1}(y_n).$$

For each $k = 1, \dots, n$ and x in $F \cap \text{int } S^{-1}(y_k)$, there is a continuous function g_x on X such that $0 \leq g_x \leq 1$, $g_x(x) = 1$ and g_x vanishes outside $\text{int } S^{-1}(y_k)$. By the compactness of F , there are finitely many g_x such that for every point in F at least one of them assumes a value not less than $1/2$. Summing them in an appropriate way, we will have nonnegative continuous functions g_1, \dots, g_n on X such that g_k vanishes outside $\text{int } S^{-1}(y_k)$, and $\sum_{k=1}^n g_k(x) \geq 1/2$ for all x in F . Let $V = \{x \in X : \sum_{k=1}^n g_k(x) > 1/3\}$. Set $f_j(x) = g_j(x) / \sum_{k=1}^n g_k(x)$ on V , and $f_j(x) = 3g_j(x)$ on $X \setminus V$. Define a continuous function $f_V : X \rightarrow E$ by

$$f_V(x) = \sum_{k=1}^n f_k(x)y_k, \quad \forall x \in X.$$

For each x in V and for each k with $f_k(x) \neq 0$, we have $x \in \text{int } S^{-1}(y_k)$. Hence, $y_k \in S(x)$. Consequently, $f_V(x) \in \text{co}(S(x)) = S(x) \subseteq K$ for all x in V . In other words, the restriction of f_V to V gives rise to a continuous selection of S on the open set V which contains F .

Denote by

$\mathcal{W} = \{(f_W, W) : \text{where } W \text{ is an open subset of } X \text{ containing } F \text{ and}$

$f_W : W \rightarrow K \text{ gives rise to a continuous selection of } S \text{ on } W\}$.

Then \mathcal{W} is not empty as $(f_V, V) \in \mathcal{W}$. Order \mathcal{W} by extension and we get a nonempty partially ordered set. In other words, $(f_W, W) \leq (f_V, V)$ if $W \subseteq V$ and $f_V|_W = f_W$. Applying Zorn's Lemma, we get a maximal element (f_U, U) of \mathcal{W} .

The last step is to verify that U is dense in X . Suppose not and there were an x in X outside the closure of U . Let $x \in \text{int } S^{-1}(y)$ for some y in K . By setting $f|_W \equiv y$, we get a continuous selection of S on an open neighborhood W of x

disjoint from U by restriction. Then the union $f_{U \cup W} : U \cup W \rightarrow K$ defined in a natural way provides a contradiction to the maximality of (f_U, U) . \square

We call a topological space X *residually paracompact* if for every open dense subset U of X the complement $X \setminus U$ is paracompact.

Theorem 3.2. *In addition to conditions (a) and (b) in Theorem 3.1, if we assume further that*

(c) X is residually paracompact,

then there is a continuous function $f : X \rightarrow K$ such that $f(x) \in S(x)$ for all x in X .

Proof. It follows from Theorem 3.1 that there is a continuous function $f_U : U \rightarrow K$ defined on an open dense subset U of X with $f_U(x) \in S(x)$ for all x in U . For each z in $X \setminus U$, there is a y in K such that $z \in \text{int } S^{-1}(y)$ by condition (b). By setting $f|_{W_z} \equiv y$ we get a continuous selection of S on an open neighborhood W_z of z . The paracompactness of $X \setminus U$ ensures it has a locally finite covering by open sets in X , each of which is contained in some W_z . Adding one more open set U , we have a locally finite open covering of X . This provides us with a family $\{g_\lambda\}_\lambda$ of nonzero continuous functions from X into $[0, 1]$ dominated by the open sets W_z and U such that $g_\lambda(x) = 0$ for all but finitely many λ 's and $\sum_\lambda g_\lambda(x) = 1$ for all x in X . If g_λ vanishes outside U , we set $f_\lambda = f_U$. Otherwise, we fix a choice of z such that g_λ vanishes outside W_z , and set $f_\lambda = f_{W_z}$. Define $f : X \rightarrow K$ by

$$f(x) = \sum_{\lambda} g_{\lambda}(x) f_{\lambda}(x), \quad \forall x \in X.$$

For each x in X , only finitely many $g_\lambda(x)$'s are nonzero in the sum, and the nonzero terms give rise to a convex combination of points in the convex set $S(x)$. Thus $f(x) \in S(x)$ for all x in X . \square

It is easy to see that the following corollary follows from Theorem 3.1.

Corollary 3.3. *The conclusion of Theorem 3.1 remains true if conditions (a) and (b) are replaced by*

- (a)' for each x in X , the set $S(x)$ is a nonempty convex set;
- (b)' for each y in K , the set $S^{-1}(y)$ is open.

Remark 3.4. Corollary 3.3 implies Theorem 3.1.

Proof. Let $T : X \rightarrow 2^K$ be defined by

$$T(x) = \{y \in K : x \in \text{int } S^{-1}(y)\}.$$

Then $T^{-1}(y) = \text{int } S^{-1}(y)$ is open for each y in K . By (b), for each x in X , there exists y in K such that $x \in \text{int } S^{-1}(y)$. Therefore $y \in T(x) \neq \emptyset$ for each x in X . Let $H : X \rightarrow 2^K$ be defined by $H(x) = \text{co } T(x)$. Then $H(x)$ is nonempty for each x in X , and $H^{-1}(y)$ is open for each y in K . By Corollary 3.3, there is an open dense subset U of X , containing any but a fixed compact set D , and there is a continuous function $f : U \rightarrow K$ such that $f(x) \in H(x) = \text{co } T(x) \subset S(x)$ for all x in U . \square

4. FIXED POINT THEOREMS

Theorem 4.1. *For each i in a nonempty index set I , let X_i be a nonempty convex subset of a Hausdorff locally convex topological vector space E_i , and let D_i be a compact subset of X_i . Let $X = \prod_{i \in I} X_i$ be the product space. Let $F_i : X \rightarrow 2^{D_i}$ be sub-lower semicontinuous with nonempty convex values. Then for every neighborhood V_i of 0 in E_i , there exists a point $\bar{x}_V = (x_{V_i})$ in $D = \prod_{i \in I} D_i$ such that $(\bar{x}_{V_i} + V_i) \cap F_i(\bar{x}_V) \neq \emptyset$ for all i in I .*

Proof. Let V_i be a neighborhood of zero in E_i for each i in I . Fix any i in I . There exists an absolutely convex neighborhood W_i of 0 such that $W_i \subset V_i$. Note that D is a compact subset of X . By Lemma 2.4, $\text{co } D$ is a paracompact subset of X . Since $F_i : X \rightarrow 2^{D_i}$ is a sub-lower semicontinuous multi-valued map with nonempty convex values, by Lemma 2.1 there exists a continuous function $f_i : \text{co } D \rightarrow D_i$ such that

$$f_i(x) \in (F_i(x) + W_i) \cap D_i \text{ for each } x \in \text{co } D.$$

Define $f : \text{co } D \rightarrow D$ by $f(x) = \prod_{i \in I} f_i(x)$ for x in $\text{co } D$. By the Himmelberg fixed point theorem (Lemma 2.3), there exists an $\bar{x}_V = (\bar{x}_{V_i})_{i \in I}$ in $\text{co } D$ such that $\bar{x}_V = f(\bar{x}_V) = \prod_{i \in I} f_i(\bar{x}_V)$. That is, $\bar{x}_{V_i} = f_i(\bar{x}_V) \in (F_i(\bar{x}_V) + W_i) \cap D_i$. Thus, $\bar{x}_{V_i} \in D_i$ and $(\bar{x}_{V_i} + W_i) \cap F_i(\bar{x}_V) \neq \emptyset$ for all i in I . Since $W_i \subset V_i$, we have $(\bar{x}_i + V_i) \cap F_i(\bar{x}_V) \neq \emptyset$ for all i in I . □

Theorem 4.2. *Suppose in Theorem 4.1 we assume further that for each $x = (\bar{x}_i)_{i \in I} \in X$, its coordinates $x_i \notin \overline{F_i(x)} \setminus F_i(x)$ for all i in I . Then there exists a point $\bar{x} = (\bar{x}_i)_{i \in I} \in D = \prod_{i \in I} D_i$ such that $\bar{x}_i \in F_i(\bar{x})$ for each i in I .*

Proof. For each i in I , let \mathcal{B}_i be the collection of all absolutely convex open neighborhoods of zero in E_i and $\mathcal{B} = \prod_{i \in I} \mathcal{B}_i$. Given any $V = \prod_{i \in I} V_i$ in \mathcal{B} , let $Q_V = \{x \in D : x_i \in \overline{F_{V_i}(x)} \text{ for all } i \text{ in } I\}$. Then Q_V is a nonempty closed subset of D for each V in \mathcal{B} by Theorem 4.1. Let $\{V^{(1)}, \dots, V^{(n)}\}$ be any finite subset of \mathcal{B} . Write $V^{(i)} = \prod_{j \in I} V_j^{(i)}$, where $V_j^{(i)} \in \mathcal{B}_j$ for each $i = 1, \dots, n$. Let $V' = \prod_{j \in I} (\bigcap_{i=1}^n V_j^{(i)}) \in \mathcal{B}$. Clearly, $\emptyset \neq Q_{V'} \subseteq \bigcap_{i=1}^n Q_{V^{(i)}}$. Therefore, the family $\{Q_V : V \in \mathcal{B}\}$ has the finite intersection property. Since $Q_V \subset D$ for all V in \mathcal{B} and D is compact, $\bigcap_{V \in \mathcal{B}} Q_V \neq \emptyset$. Let $\bar{x} = (\bar{x}_i)_{i \in I} \in \bigcap_{V \in \mathcal{B}} Q_V$. Then $\bar{x}_i \in \overline{F_{V_i}(\bar{x})}$ for all i in I and all V_i in \mathcal{B}_i , i.e., $\bar{x}_i \in \bigcap_{V_i \in \mathcal{B}_i} \overline{F_{V_i}(\bar{x})}$ for all i in I . It follows from Lemma 2.2 that $\bar{x}_i \in \overline{F_i(\bar{x})}$ for all i in I . By assumption, $\bar{x}_i \in F_i(\bar{x})$ for all i in I . □

We remark that if F_i is closed, then $x_i \notin \overline{F_i(x)} \setminus F_i(x)$ for each $x = (x_i)_{i \in I}$ in X . As a special case of Theorem 4.2, we have the following collectively Himmelberg type fixed point theorem.

Corollary 4.3. *For each i in a nonempty index set I , let X_i be a nonempty convex subset of a locally convex topological vector space E_i , let D_i be a nonempty compact subset of X_i , and let $f_i : X = \prod_{i \in I} X_i \rightarrow D_i$ be a continuous function. Then there exists $\bar{x} = (\bar{x}_i)_{i \in I} \in D = \prod_{i \in I} D_i$ such that $\bar{x}_i = f_i(\bar{x})$ for each i in I .*

If the index set I is a singleton, then Theorem 4.2 reduces to the following corollary, which provides a partial solution to a conjecture of Wu [15].

Corollary 4.4. *Let X be a nonempty convex subset of a locally convex topological vector space E , let D be a nonempty compact subset of X , and let $F : X \rightarrow 2^D$ be sub-lower semicontinuous with nonempty convex values. Suppose $x \notin \overline{F}(x) \setminus F(x)$ for each x in X . Then there exists a point \bar{x} in D such that $\bar{x} \in F(\bar{x})$.*

By Theorem 4.1, we have the following almost fixed point theorem.

Corollary 4.5. *The conclusions of Theorems 4.1 and 4.2 remain valid if the condition “ $F_i : X \rightarrow 2^{D_i}$ is sub-lower semicontinuous for each i in I ” is replaced by “ $F_i^{-1}(y_i)$ is open for each y_i in D_i and each i in I .”*

Finally we remark that in case I is a singleton, Theorem 4.1 provides a different result from [12, Theorem 3].

REFERENCES

- [1] R. Agarwal and D. O'Regan, Fixed point theory for maps with lower semicontinuous selections and equilibrium theory for abstract economies, *J. Nonlinear and Convex Analysis*, **2** (2001), 31-46. MR1828157 (2002m:49025)
- [2] H. Ben-El-Mechaiekh, The coincidence problem for compositions of set-valued maps, *Bull. Austral. Math. Soc.*, **41** (1990), 421-434. MR1071044 (91h:47060)
- [3] H. Ben-El-Mechaiekh, Fixed points for compact set-valued maps, *Questions Answers Gen. Topology*, **10**(1992), 153-156. MR1180474
- [4] F. E. Browder, A new generation of the Schauder fixed point theorem, *Math. Ann.*, **174** (1967), 285-290. MR0223944 (36:6991)
- [5] F. E. Browder, The fixed point theory of multi-valued mappings in topological vector spaces, *Math. Ann.*, **177** (1968), 283-301. MR0229101 (37:4679)
- [6] P. Cubiotti, Some remarks on fixed points of lower semicontinuous multifunctions, *J. Math. Anal. Appl.*, **174** (1993), 407-412. MR1215621 (94c:47081)
- [7] F. Deutsch and P. Kenderov, Continuous selections and approximate selection for set-valued mappings and applications to metric projections, *SIAM J. Math. Anal.*, **14** (1983), 185-194. MR0686245 (84c:54026)
- [8] X. P. Ding, W. K. Kim and K. K. Tan, A selection theorem and its applications, *Bull. Austral. Math. Soc.*, **46** (1992), 205-212. MR1183778 (93g:47072)
- [9] A. Granas, *Points fixes pour les applications compactes: espaces de Lefschetz et la théorie de l'indice*, Séminaire de Mathématiques Supérieures **68**, Presses de l'Université de Montréal, Montréal, Que., 1980. MR0569745 (81i:55002)
- [10] C. J. Himmelberg, Fixed points of compact multifunctions, *J. Math. Anal. Appl.*, **38** (1972), 205-207. MR0303368 (46:2505)
- [11] C. D. Horvath, Extension and selection theorems in topological vector spaces with a generalized convexity structure, *Ann. Fac. Sci., Toulouse* **2**, (1993), 253-269. MR1253391 (94i:54043)
- [12] S. Park, Continuous selection theorems in generalized convex spaces, *Numer. Funct. Anal. Optim.*, **25** (1999), 567-583. MR1704961 (2000e:54014)
- [13] S. Park, The Knaster-Kuratowski-Mazurkiewicz Theorem and almost fixed points, *Top. Methods in Nonlinear Anal.*, **16** (2000), 195-200. MR1805047 (2001m:47118)
- [14] G. Tian, Fixed point theorems for mappings with noncompact and nonconvex domains, *J. Math. Anal. Appl.*, **158** (1991), 161-167. MR1113407 (92e:47110)
- [15] X. Wu, A new fixed point theorem and its applications, *Proc. Amer. Math. Soc.*, **125** (1997), 1779-1783. MR1397000 (97h:90014)
- [16] X. Wu and S. Shen, A further generalization of Yannelis-Prabhakar's continuous selection theorem and its applications, *J. Math. Anal. Appl.*, **197** (1996), 61-74. MR1371276 (97a:47085)
- [17] N. C. Yannelis and N. D. Prabhakar, Existence of maximal elements and equilibria in linear topological spaces, *J. Math. Economics*, **12** (1983), 233-245. MR0743037 (87h:90061a)
- [18] Z. T. Yu and L. J. Lin, Continuous selection and fixed point theorems, *Nonlinear Anal.*, **52** (2003), 445-455. MR1937632 (2003g:54040)

- [19] G. X. Z. Yuan, G. Isac, K. K. Tan, and J. Yu, The study of minimax inequalities, abstract economics and applications to variational inequalities and Nash equilibria, *Acta Applicandae Mathematicae*, **54** (1998), 135-166. MR1660931 (99j:49018)
- [20] X. Zheng, Approximate selection theorems and their applications, *J. Math. Anal. Appl.*, **212** (1997), 88-97. MR1460186 (98h:90118)

DEPARTMENT OF MATHEMATICS, NATIONAL CHANGHUA UNIVERSITY OF EDUCATION, CHANGHUA, 50058, TAIWAN

E-mail address: maljlin@math.ncue.edu.tw

DEPARTMENT OF APPLIED MATHEMATICS, NATIONAL SUN YAT-SEN UNIVERSITY, AND NATIONAL CENTER FOR THEORETICAL SCIENCES, KAOHSIUNG, 80424, TAIWAN

E-mail address: wong@math.nsysu.edu.tw

DEPARTMENT OF ELECTRICAL ENGINEERING, NAN-KAI INSTITUTE OF TECHNOLOGY, NANTOUR 542, TAIWAN