

TOTALLY GEODESIC BOUNDARIES OF KNOT COMPLEMENTS

RICHARD P. KENT IV

(Communicated by Ronald A. Fintushel)

for Kimberly

ABSTRACT. Given a compact orientable 3-manifold M whose boundary is a hyperbolic surface and a simple closed curve C in its boundary, every knot in M is homotopic to one whose complement admits a complete hyperbolic structure with totally geodesic boundary in which the geodesic representative of C is as small as you like.

1. INTRODUCTION

Let Σ be a closed hyperbolic surface and let \mathfrak{H} be the set of all orientable 3-manifolds admitting a complete finite volume hyperbolic structure with totally geodesic boundary homeomorphic to Σ .

By Mostow-Prasad rigidity, there is a map $\widehat{\partial}$ from \mathfrak{H} to the moduli space of Σ sending a manifold to the hyperbolic structure that appears on its boundary. It is a theorem of Fujii and Soma that the image of $\widehat{\partial}$ is dense [13].

Given a hyperbolic structure σ , Fujii and Soma construct a sequence of manifolds whose boundaries converge to σ using Brooks' theorem [9] that the set of hyperbolic structures admitting circle packings is dense. Given $\varepsilon > 0$, a circle packing point τ is chosen within $\frac{\varepsilon}{2}$ of σ . Then a finite volume manifold whose totally geodesic boundary consists of two copies of τ and two copies of its mirror image is built explicitly from the circle packing. Two of the boundary components are then identified and hyperbolic Dehn surgeries performed to obtain a manifold with no cusps whose boundary components are within $\frac{\varepsilon}{4}$ of τ and its mirror image. A number of copies of this manifold are then glued end to end and finally attached to a manifold having totally geodesic boundary merely homeomorphic to Σ . A large number may be necessary to ensure that the boundary of this manifold is within ε of σ . In particular, this number tends to infinity as ε tends to zero.

With a view towards grasping the relationship between manifold and boundary, we are motivated by the hope that passage to Fujii and Soma's theorem may be made with greater topological control—note, for instance, that as Fujii and Soma's 3-manifolds approach σ , their first Betti numbers tend to infinity.

Received by the editors May 12, 2004 and, in revised form, August 7, 2004.
2000 *Mathematics Subject Classification*. Primary 57M50.

This work was supported in part by a University of Texas Continuing Fellowship.

©2005 American Mathematical Society
Reverts to public domain 28 years from publication

To this end, let M be a compact orientable 3-manifold whose boundary is homeomorphic to Σ and let \mathfrak{L}_M be the set of all complements of links in M that lie in \mathfrak{H} , \mathfrak{K}_M the set of all such knot complements. Fixing a marking on the boundary of M , rigidity again yields a map ∂ from \mathfrak{L}_M to the Teichmüller space of Σ sending a manifold to the marked hyperbolic structure that appears on its boundary.

Question. Is $\partial\mathfrak{L}_M$ dense? Is $\partial\mathfrak{K}_M$?

We prove the following theorem.

Theorem 1. *Let C be a simple closed curve in ∂M and let $\varepsilon > 0$. Then every knot in M is homotopic to one whose complement admits a hyperbolic structure with totally geodesic boundary in which the length of the geodesic representative of C is less than ε .*

The proof proceeds as follows. A surface F is chosen in M whose boundary contains C . A theorem of Myers allows us to homotope a given knot to one whose complement admits a hyperbolic structure with totally geodesic boundary. We may choose this knot to intersect F a number of times so that the intersection of F with the resulting knot complement is a quasi-Fuchsian surface. A pseudo-Anosov mapping class ψ is then found with the property that cutting open along this quasi-Fuchsian surface and precomposing the gluing map with ψ^n yields the complement of a knot in the same homotopy class. Examination of the final gluing step of the proof of Thurston's Uniformization Theorem and a theorem of J. Brock reveal that these knots are cinching C .

2. NOTATION AND MACHINERY

2.1. Topology. Given two subsets X and Y of a set Z we let $X - Y = X - (X \cap Y)$ denote the set of elements of X that are not elements of Y . If X and Y are disjoint and $f: X \rightarrow W$ and $g: Y \rightarrow W$ are two functions, we let $f \cup g$ denote the function $X \cup Y \rightarrow W$ whose restrictions to X and Y are f and g respectively.

If G is a group and $g, h \in G$, we denote the word $g^{-1}h^{-1}gh$ by $[g, h]$.

Given a topological space X , let $|X|$ denote the number of path components of X .

If M is an n -manifold and F is a union of transverse properly embedded submanifolds in M , we write $M \setminus F = M - \text{int nhd}(F)$. If F is a bicollared $(n-1)$ -manifold, $\text{nhd}(F) \cong [-1, 1] \times F$ and we call $\{-1\} \times F$ and $\{1\} \times F$ the *traces* of F in $M \setminus F$. For two transverse submanifolds F and F' , we often write $F \setminus F'$ for $F \setminus (F \cap F')$.

A 3-manifold is said to be *atoroidal* if every one of its embedded incompressible tori is boundary parallel. A properly embedded annulus in a 3-manifold M is *essential* if it is incompressible and is not isotopic relative to its boundary to an annulus in ∂M . A 3-manifold M is said to be *cylindrical* if it contains an essential annulus. M is said to be *acylindrical* otherwise.

A *pared 3-manifold* is a pair (M, P) where M is an orientable compact irreducible 3-manifold and P is a subset of ∂M consisting of pairwise nonisotopic (in ∂M) embedded incompressible tori and annuli with the property that for $X \in \{S^1 \times S^1, S^1 \times I\}$, every map $(X, \partial X) \rightarrow (M, P)$ is homotopic as a map of pairs into P . We call P the *pared locus*.

If (M, P) is a pared manifold with $R = \overline{\partial M - P} \neq \emptyset$, we let $dM = M \cup_R M$ denote the *double* of M —the gluing map is the identity on R . A double dM admits

an involution i that exchanges the two copies of M . We consider dM as a pared manifold with pared locus $dP = P \cup i(P)$.

2.2. Teichmüller space and Kleinian groups. We refer the reader to [2, 1, 14] for more on quasiconformal mappings and Teichmüller theory, [17] for more on Kleinian groups, and [10] for more on convergence of sequences of hyperbolic manifolds.

An *analytically finite surface* S' is a Riemann surface conformally equivalent to a compact Riemann surface from which a finite set of points has been deleted. The surface $\overline{S'}$ is the Riemann surface obtained from S' by postcomposing every chart with complex conjugation—this canonically reverses the orientation of S' . Fix once and for all a compact surface S whose interior is homeomorphic to S' . A *marking* of S' is a choice of homotopy equivalence $S \rightarrow S'$. The Teichmüller space of S parameterizes the marked complete analytically finite hyperbolic structures on the interior of S up to isotopy and is denoted $\mathcal{T}(S)$.

Let S_1 and S_2 be analytically finite surfaces equipped with markings $f_1: S \rightarrow S_1$ and $f_2: S \rightarrow S_2$. A homeomorphism $f: S_1 \rightarrow S_2$ is K -*quasiconformal* if it is absolutely continuous on lines and $|f_{\bar{z}}| \leq k|f_z|$ in every local coordinate z , where $k = (K - 1)/(K + 1) < 1$. The minimum possible value of K for which f is K -quasiconformal is called the *dilatation* of f , and is denoted $K[f]$. The *Teichmüller distance* between S_1 and S_2 is then

$$d_{\mathcal{T}}(S_1, S_2) = \frac{1}{2} \inf \log K[f]$$

where the infimum is taken over all quasiconformal maps $f: S_1 \rightarrow S_2$ homotopic to $f_2 \circ f_1^{-1}$. There is a unique extremal quasiconformal map $S_1 \rightarrow S_2$ realizing the above distance, called the *Teichmüller mapping*.

We follow the convention that if S is disconnected, then its Teichmüller space is the product of the Teichmüller spaces of its components equipped with the metric obtained by taking the maximum of the Teichmüller metrics on the factors. If S is of genus g and connected, $\mathcal{T}(S)$ is homeomorphic to an open Euclidean ball of dimension $6g - 6 + 2|\partial S|$.

The mapping class group $\text{Mod}(S) = \pi_0(\text{Homeo}^+(\text{int } S))$ acts properly discontinuously on $\mathcal{T}(S)$ by pulling back hyperbolic structures. The moduli space $\mathcal{M}(S)$ is the quotient of $\mathcal{T}(S)$ by this action. A mapping class is said to be *reducible* if it fixes a collection \mathcal{C} of disjoint nonperipheral simple closed curves up to isotopy. Elements of \mathcal{C} are called *reducing curves*. A mapping class φ is said to be *pseudo-Anosov* if no nontrivial power of φ is reducible.

We let $\mathcal{P}(S)$ denote the mapping class group of isotopy classes of homeomorphisms of S that do not permute the components of ∂S . Since we are not requiring boundary components to be fixed pointwise, we may regard $\mathcal{P}(S)$ as a subgroup of $\text{Mod}(S)$.

Let \mathbb{H}^3 denote hyperbolic 3-space, $S_{\infty}^2 = \widehat{\mathbb{C}}$ its sphere at infinity.

For our purposes, a *Kleinian group* Γ is a discrete subgroup of $\text{PSL}_2(\mathbb{C}) \cong \text{Isom}^+(\mathbb{H}^3)$. We write $M_{\Gamma} = \mathbb{H}^3/\Gamma$. The *limit set* Λ_{Γ} of Γ is the smallest nonempty closed subset of S_{∞}^2 that is invariant under the action of Γ .

Let H_{Γ} denote the hyperbolic convex hull of Λ_{Γ} in \mathbb{H}^3 . A Kleinian group Γ is said to be *geometrically finite* if there is an open neighborhood of H_{Γ}/Γ in M_{Γ} of finite volume.

The *domain of discontinuity* of a Kleinian group Γ is the set $\Omega_\Gamma = S_\infty^2 - \Lambda_\Gamma$ on which Γ acts properly discontinuously. By the Ahlfors Finiteness Theorem, the surface Ω_Γ/Γ is analytically finite whenever Γ is nonelementary—see [15], Section 4.19, and the references cited there. This surface inherits a conformal structure from that of $\widehat{\mathbb{C}}$. By the Uniformization Theorem, Ω_Γ/Γ also comes equipped with a hyperbolic structure. The manifold $\dot{M}_\Gamma = (H^3 \cup \Omega_\Gamma)/\Gamma$ is the *Kleinian manifold associated to Γ* .

A *hyperbolic structure on a pared manifold* (M, P) is a discrete faithful representation $\pi_1(M, *) \rightarrow \mathrm{PSL}_2(\mathbb{C})$ with image Γ in which the images of the subgroups corresponding to the components of P consist entirely of parabolic elements and such that $\mathbb{H}^3/\Gamma \cong \mathrm{int} M$. The Kleinian manifold \dot{M}_Γ is homeomorphic to $M - P$. A hyperbolic structure is *geometrically finite* if Γ is geometrically finite.

A hyperbolic structure is said to be a *hyperbolic structure with totally geodesic boundary* if the images of the subgroups corresponding to the components of $\partial M - P$ are Fuchsian.

Two hyperbolic structures are deemed *equivalent* if their images are conjugate in $\mathrm{PSL}_2(\mathbb{C})$, and we let $H(M, P)$ denote the set of equivalence classes. We continue to refer to equivalence classes as hyperbolic structures.

Let $\mathrm{AH}(M, P)$ denote the set $H(M, P)$ equipped with the topology induced by the inclusion

$$H(M, P) \subset \mathrm{Hom}(\pi_1(M, *), \mathrm{PSL}_2(\mathbb{C}))/\mathrm{PSL}_2(\mathbb{C})$$

equipped with the quotient of the compact-open topology—this is the topology induced by convergence on a set of generators after conjugating and is called the *algebraic topology*.

Let M and N be two manifolds equipped with hyperbolic metrics d_M and d_N respectively and let $K > 1$. A diffeomorphism $f: M \rightarrow N$ satisfying

$$K^{-1}d_M(x, y) \leq d_N(f(x), f(y)) \leq Kd_M(x, y),$$

for all x and y in M is called a *K -quasi-isometry*.

2.3. Quasi-Fuchsian groups and iteration near a Bers slice. A Kleinian group Γ is *quasi-Fuchsian* if Λ_Γ is a topological circle and Γ preserves each component Ω_1 and Ω_2 of Ω_Γ setwise. The surfaces $S_1 = \Omega_1/\Gamma$ and $S_2 = \Omega_2/\Gamma$ are homeomorphic by an orientation-reversing homeomorphism and each comes equipped with a hyperbolic structure of its own. So, associated to Γ is a point in $\mathcal{T}(S) \times \mathcal{T}(\overline{S})$ once a marking $S \rightarrow S_1$ has been chosen. We call S the *underlying surface of Γ* . The manifold \mathbb{H}^3/Γ is called the *quasi-Fuchsian manifold associated to Γ* and is homeomorphic to $S \times \mathbb{R}$. Let $\mathrm{QF}(S) \subset \mathrm{AH}(S \times \mathbb{R})$ denote the space of quasi-Fuchsian groups. The Ahlfors-Bers Simultaneous Uniformization Theorem [6] provides a homeomorphism $Q: \mathcal{T}(S) \times \mathcal{T}(\overline{S}) \rightarrow \mathrm{QF}(S)$, natural in the sense that $Q^{-1}(\Gamma)$ is the pair of surfaces Ω_Γ/Γ . If $A \subset \mathcal{T}(S)$ and $B \subset \mathcal{T}(\overline{S})$, we write $Q(A, B) = Q(A \times B)$ and we abbreviate $Q(\{a\}, \{b\})$ by writing $Q(a, b)$.

We will need the following lemma.

Lemma 1. *Let $A \subset \mathcal{T}(S)$ and $B \subset \mathcal{T}(\overline{S})$ be compact sets. Then there is a k such that for each $\varphi \in \mathrm{Mod}(S)$, all quasi-Fuchsian manifolds associated to elements of $Q(\varphi A, B)$ are k -quasi-isometric.*

Proof. Since A is compact and $\text{Mod}(S)$ acts on $\mathcal{T}(S)$ by isometries of the Teichmüller metric, the diameter of φA in this metric is bounded and independent of φ .

Fix $b \in B$, $a_1, a_2 \in A$. Let $\Omega_i \subset \Omega_{Q(\varphi a_i, b)}$ be the components corresponding to φa_i , $\widehat{\Omega}_i \subset \Omega_{Q(\varphi a_i, b)}$ the components corresponding to b and let $h_i: U \rightarrow \Omega_i$ and $g_i: L \rightarrow \widehat{\Omega}_i$ be Riemann mappings from the upper and lower halfplanes in $\widehat{\mathbb{C}}$, respectively, for $i \in \{1, 2\}$. Let Γ_i be the Fuchsian group uniformizing φa_i for each i . We may lift the Teichmüller mapping $\varphi a_1 \rightarrow \varphi a_2$ to obtain a quasiconformal mapping $\omega: U \rightarrow U$ such that $\omega^{-1}\Gamma_2\omega = \Gamma_1$ and such that $h_1\omega^{-1}h_2^{-1} \cup g_1g_2^{-1}$ extends to a quasiconformal mapping $w: \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}$ with $wQ(\varphi a_1, b)w^{-1} = Q(\varphi a_2, b)$ —compare Section 6 of [4].

Since the diameter of φA is uniformly bounded, the dilatation of this map is also uniformly bounded as the a_i range over φA .

Symmetry allows us to vary b in B , and so there is a constant k' depending only on A and B such that any $Q(\varphi a, b)$ and $Q(\varphi a', b')$ are k' -quasiconformally conjugate Kleinian groups whenever $a, a' \in A$ and $b, b' \in B$. Such quasiconformal conjugations extend to equivariant k -quasi-isometries between universal covers of quasi-Fuchsian manifolds—see Theorem 2.5 of [18]—with k uniform over $Q(\varphi A, B)$ and independent of φ . By equivariance, these k -quasi-isometries descend to k -quasi-isometries between quasi-Fuchsian manifolds. \square

Let $\varphi \in \text{Mod}(S)$ be reducible of infinite order. By passing to a suitable power, we may assume that φ fixes every reducing curve so that φ is pseudo-Anosov or trivial when restricted to any component of the complement of the reducing curves and we do so. Consider the sequence of quasi-Fuchsian groups $Q(\varphi^n a, b)$ and their quasi-Fuchsian manifolds M_n . It is a theorem of J. Brock [8] that the lengths of the reducing curves in M_n tend to zero as n approaches infinity.

Lemma 1 will allow us to apply Brock’s theorem to sequences of quasi-Fuchsian groups whose n^{th} terms lie in sets $Q(\varphi^n A, B)$.

2.4. Uniformization and the skinning map. We review here the final gluing step of the proof of Thurston’s Uniformization Theorem for Haken Manifolds and refer the reader to [19], [15] and [17] for more information regarding this theorem, the Bounded Image Theorem, and Maskit combination.

Let S be a two-sided properly embedded surface in a pared 3-manifold (M, P) . Let $M' = M \setminus S$ and let S_1 and S_2 denote the traces of S in M' . Let $f: S_1 \rightarrow S_2$ be the gluing map such that $M = M'/f$. We mark S_1 with $f_1: S \rightarrow S_1$ and S_2 with $f \circ f_1$. Precomposing f with a representative of a mapping class $\varphi \in \text{Mod}(S_1)$ yields a new gluing map g and we let $M(\varphi) = M'/g$.

Given a geometrically finite hyperbolic structure ρ on a pared manifold (M, P) with image Γ , we obtain a point in $\mathcal{T}(\partial M - P)$ by considering Ω_Γ/Γ . Conversely, by a theorem of Bers [5], if (M, P) admits such a hyperbolic structure, then the set of all geometrically finite hyperbolic structures on (M, P) is parameterized by $\mathcal{T}(\partial M - P)$.

So, let $(M, \partial M)$ be a pared 3-manifold that does not admit a Seifert fibration and suppose that S is a two-sided properly embedded incompressible surface such that $(N, P) = (M \setminus S, \partial M \setminus \partial S)$ is acylindrical and admits no Seifert fibration. Let S_1 and S_2 denote the traces of S in N equipped with markings $f_i: S \rightarrow S_i$ as above. Then (N, P) admits a geometrically finite hyperbolic structure ρ by the

Uniformization Theorem for Haken Manifolds and so the set of such structures is parameterized by $\mathcal{T}(\partial N - P) \cong \mathcal{T}(S_1) \times \mathcal{T}(S_2)$.

Given an element of $\mathcal{T}(S_1) \times \mathcal{T}(S_2)$, consider the geometrically finite structure on (N, P) to which it is associated. Now, for any pared 3-manifold (M', P') equipped with a geometrically finite hyperbolic structure, the subgroup corresponding to a component of $\partial M' - P'$ is quasi-Fuchsian or else M' is cylindrical, by Proposition 7.2 of [19] and the Annulus Theorem. Since (N, P) is acylindrical, the images $\Gamma_1 = \rho(\pi_1(S_1))$ and $\Gamma_2 = \rho(\pi_1(S_2))$ are quasi-Fuchsian groups. Now, Γ_1 determines a point $(a, \sigma_1(a)) \in \mathcal{T}(S_1) \times \mathcal{T}(\overline{S_1})$. The first coordinate is the conformal structure that appears on the S_1 boundary component of (N, P) , and $\sigma_1(a)$ is called the *conformal structure that S_1 hides*. The map

$$\sigma_1: \mathcal{T}(S_1) \longrightarrow \mathcal{T}(\overline{S_1})$$

is called the *skinning map (associated to N)*.

Since (N, P) is acylindrical, σ_1 has precompact image—this is the content of Thurston's Bounded Image Theorem. We have a similar discussion and a map σ_2 associated to S_2 . For $i \in \{1, 2\}$, let $\overline{\sigma}_i: \mathcal{T}(S_i) \rightarrow \mathcal{T}(S_i)$ denote σ_i followed by the map induced by reversing the orientation of $\overline{S_i}$.

Let $\varphi \in \text{Mod}(S_1)$. Precomposing the gluing map f by a representative of φ yields two sequences of maps:

$$\mathcal{T}(S_1) \xrightarrow{\overline{\sigma}_1} \mathcal{T}(S_1) \xrightarrow{\varphi_*} \mathcal{T}(S_1) \xrightarrow{f_*} \mathcal{T}(S_2)$$

and

$$\mathcal{T}(S_2) \xrightarrow{\overline{\sigma}_2} \mathcal{T}(S_2) \xrightarrow{f_*^{-1}} \mathcal{T}(S_1) \xrightarrow{\varphi_*^{-1}} \mathcal{T}(S_1).$$

To obtain a hyperbolic structure for $(M, \partial M)$, an application of one of two of the Maskit Combination Theorems is applied. Satisfaction of the hypotheses is easily translated into the existence of a point $(x, y) \in \mathcal{T}(S_1) \times \mathcal{T}(S_2)$ such that

$$\overline{\sigma}_2(y) = f_* \varphi_*(x)$$

and

$$\overline{\sigma}_1(x) = \varphi_*^{-1} f_*^{-1}(y).$$

We call this point the *solution to the gluing problem for $f\varphi$* . The existence of such a point means that there is a geometrically finite hyperbolic structure on (N, P) so that the conformal structure at infinity associated to S_1 is carried by our gluing map to the conformal structure that S_2 hides, and vice versa.

3. KNOTS

Let M be a compact orientable 3-manifold whose boundary is homeomorphic to Σ and let C be a simple closed curve in ∂M . If C is separating in ∂M , let F' be a surface in ∂M that it bounds. If C is nonseparating, it is a boundary component of a pair of pants F' whose boundary components are pairwise nonisotopic in ∂M . In either case, let F be a properly embedded surface in M isotopic to F' .

Lemma 2. *Every homotopy class of simple closed curves in M contains a knot K such that $M \setminus K$ admits a hyperbolic structure with totally geodesic boundary and $M \setminus (F \cup K)$ is boundary incompressible and acylindrical.*

Proof. Let γ be a homotopy class and let k be a simple closed curve representing γ that intersects F transversely in at least two points. Consider $k' = k \setminus F \subset M \setminus F$. By Theorem 1.1 of [21], there is a 1-manifold k'' in the homotopy class of k' relative to its boundary such that each component of $(M \setminus F) \setminus k''$ is boundary incompressible, acylindrical and atoroidal. Let K be the knot in M corresponding to k'' .

The knot exterior $M \setminus K$ is irreducible, atoroidal and acylindrical by Lemma 2.1 of [21]. The double $d(M \setminus K)$ admits a hyperbolic structure by the Uniformization Theorem for Haken Manifolds, and we equip $d(M - K)$ with the associated hyperbolic metric. The natural orientation-reversing involution on $d(M - K)$ is homotopic to an isometry that preserves a totally geodesic surface, by Mostow-Prasad rigidity. A cut and paste argument demonstrates that this surface is isotopic to $\partial(M - K)$ and cutting the double open again yields a hyperbolic structure with totally geodesic boundary on $M \setminus K$. \square

Scholium. *Given M, F and K as in Lemma 2, cutting $M \setminus K$ along $F_K = F \setminus K$ and regluing the traces of F_K in a different fashion yields a hyperbolic manifold with totally geodesic boundary no matter what mapping class we choose for the gluing.*

Proof. The conclusion of Lemma 2.1 of [21] holds for any gluing map. \square

4. PROOF OF THEOREM 1

Proof of Theorem 1. Let M, F and K be as in Lemma 2 and let $F_K = F \setminus K$. Let $(X_K, P) = (M \setminus K, \partial X_K - \partial M)$, $Y_K = dX_K$, $S = dF_K \subset Y_K$, $Z_K = Y_K \setminus S$. Let S_1, S_2 denote the traces of S in Z_K , $f: S_1 \rightarrow S_2$ the gluing map such that $Z_K/f = Y_K$. We mark S_1 with $f_1: S \rightarrow S_1$, S_2 with $f \circ f_1$. Note that Z_K is acylindrical as no component of $\partial M \setminus \partial F$ is an annulus.

Let $\{\delta_1, \dots, \delta_{\ell+1}\}$ be the set of boundary components of F_K and let \widehat{F}_K denote the surface obtained from F_K by capping off a boundary component with a disk. Let $\{\gamma_1, \dots, \gamma_\ell\}$ be the set of boundary components of \widehat{F}_K . For each δ_i , there is a short exact sequence

$$1 \rightarrow \pi_1(\widehat{F}_K) \rightarrow \mathcal{P}(F_K) \xrightarrow{\phi_i} \mathcal{P}(\widehat{F}_K) \rightarrow 1$$

where an element of $\pi_1(\widehat{F}_K)$ corresponds to the mapping class that “spins” the associated boundary component about the homotopy class and ϕ_i “forgets” δ_i , see [7]. Note that for any $\psi \in \bigcap_{i=1}^{\ell+1} \ker \phi_i$, $X_K(\psi)$ is the complement of a knot in the same homotopy class as K . We claim that $\bigcap_{i=1}^{\ell+1} \ker \phi_i$ is nontrivial. To see this, choose a basis $\{x_1, \dots, x_m\}$ for the free group $\pi_1(\widehat{F}_K)$ so that a representative for x_i is freely homotopic to γ_i whenever $1 \leq i \leq \ell - 1$ and a representative for

$$v = x_1 \dots x_{\ell-1} [x_\ell, x_{\ell+1}] [x_{\ell+2}, x_{\ell+3}] \dots [x_{m-1}, x_m]$$

is freely homotopic to γ_ℓ —such a basis is easily obtained from the standard handle decomposition of \widehat{F}_K . Since $\pi_1(\widehat{F}_K)$ is nonabelian, the mapping class associated to the commutator

$$w = [[\dots [[x_1, x_2], x_3], \dots, x_m], v]$$

is a nontrivial element of $\bigcap_{i=1}^{\ell+1} \ker \phi_i$.

We claim that w represents a pseudo-Anosov mapping class ψ . Suppose that this is not the case. By Theorem 2 of [16], w has a representative on \widehat{F}_K that does

not fill \widehat{F}_K . This implies that w is conjugate into a proper free factor of $\pi_1(\widehat{F}_K)$. It is easily verified, using techniques due to J. H. C. Whitehead [24], that this is not the case, as the Whitehead graph associated to w has no cut vertex; see [23].

Let $\varphi = d\psi$ be the double of ψ .

We consider $X_K(\psi^n)$, $Y_K(\varphi^n)$, and Z_K as pared manifolds with pared loci $\partial X_K(\psi^n) - \partial M$, $\partial Y_K(\varphi^n)$, and $\partial Z_K \setminus (S_1 \cup S_2)$ respectively, and henceforth suppress the paring in the notation.

The manifold $X_K(\psi^n)$ admits a hyperbolic structure with totally geodesic boundary by the Scholium and $Y_K(\varphi^n)$ admits a hyperbolic structure by doubling.

The latter structure may be obtained from the unique solution to the gluing problem for $f\varphi^n$ and so we turn our attention to the double Y_K cut open, Z_K .

We have the sequences obtained by precomposing f by φ^n :

$$\mathcal{T}(S_1) \xrightarrow{\bar{\sigma}_1} \mathcal{T}(S_1) \xrightarrow{\varphi_*^n} \mathcal{T}(S_1) \xrightarrow{f_*} \mathcal{T}(S_2)$$

and

$$\mathcal{T}(S_2) \xrightarrow{\bar{\sigma}_2} \mathcal{T}(S_2) \xrightarrow{f_*^{-1}} \mathcal{T}(S_1) \xrightarrow{\varphi_*^{-n}} \mathcal{T}(S_1).$$

Since Z_K is acylindrical, the Bounded Image Theorem demands that the images of the $\bar{\sigma}_i$ be contained in compact A_i for each $i \in \{1, 2\}$. The solution to the gluing problem for f lies in the compact set

$$f_*^{-1}A_2 \times f_*A_1 \subset \mathcal{T}(S_1) \times \mathcal{T}(S_2),$$

and the solution to the gluing problem for $f\varphi^n$ lies in the compact

$$\varphi_*^{-n}f_*^{-1}A_2 \times f_*\varphi_*^nA_1 \subset \mathcal{T}(S_1) \times \mathcal{T}(S_2).$$

Let $A = f_*^{-1}A_2$ and let $B \supset \sigma_1(\mathcal{T}(S_1))$ be compact. Write $\Phi = \varphi^{-1}$. In the resulting hyperbolic structure on $Y_K(\varphi^n)$, $\pi_1(S)$ is a quasi-Fuchsian group $Q(\Phi^n a_n, b_n)$ trapped in $Q(\Phi^n A, B)$.

Let $a \in A$, $b \in B$. Since Φ is reducible and pseudo-Anosov when restricted to either component of $S \setminus \partial F$, the lengths of the geodesic representatives of ∂F in the quasi-Fuchsian manifolds associated to the $Q(\Phi^n a, b)$ tend to zero as n grows, by Theorem 4.5 of [8]. By Lemma 1, the quasi-Fuchsian manifolds associated to $Q(\Phi^n a_n, b_n)$ and $Q(\Phi^n a, b)$ are quasi-isometric independent of n , and so the lengths of the geodesic representatives of ∂F in the former manifolds also tend to zero as n grows. In particular, the lengths of the geodesic representatives of C in the $Y_K(\varphi^n)$ tend to zero. Now $\partial X_K(\psi^n)$ is totally geodesic in $Y_K(\varphi^n)$ and so C is shrinking in the $\partial X_K(\psi^n)$ as well. \square

In the case that C is nonseparating, we have cinched C at the expense of cinching two other curves in ∂M . This may be remedied at the expense of the homotopy class of the knot. Consider two disjoint parallel pushoffs C' and C'' of C just inside ∂M . Pick an annulus F between C and C' disjoint from C'' and proceed as above to obtain a knot K in $M - (C' \cup C'')$ in whose complement C is short. Performing $\frac{1}{m}$ -filling on the cusp corresponding to C' and $-\frac{1}{m}$ -filling on the cusp corresponding to C'' has the effect of shearing K along an annulus whose boundary is $C' \cup C''$, yielding a knot in M . As m tends to infinity, these knot complements converge geometrically to $M - (K \cup C' \cup C'')$ (see Proposition E.6.29 of [3]), and so for large m , C is short in these manifolds. Note that this construction is easily adapted to cinch a collection of disjoint simple closed curves.

5. GEOMETRIC FLEXIBILITY AND ISOLATION

In [22], Neumann and Reid construct infinitely many examples of one-cusped hyperbolic 3-manifolds that contain totally geodesic surfaces that are *geometrically isolated* from the cusp in the sense that performing any hyperbolic Dehn filling leaves the hyperbolic structure on the totally geodesic surface unchanged. In contrast, Fujii [11] constructs an explicit one-cusped hyperbolic manifold M with totally geodesic boundary with the property that the manifolds obtained from M by filling exhibit infinitely many hyperbolic structures on their boundaries and in [12], one-cusped manifolds with totally geodesic boundaries are constructed for which the map from the hyperbolic Dehn filling space to the Teichmüller space of the boundary is actually an embedding near infinity.

We have the

Theorem 2. *Let M be a 3-manifold that admits a hyperbolic structure with totally geodesic boundary homeomorphic to Σ . Then M contains a hyperbolic knot K such that ∂M and $\partial(M - K)$ are as far apart as we like in $\mathcal{M}(\Sigma)$.*

Proof. Theorem 1 provides a knot K such that the injectivity radius of $\partial(M - K)$ is as small as we like, so we may choose K so that $\partial(M - K)$ is as far as we like from ∂M in $\mathcal{M}(\Sigma)$ —since the set of surfaces in \mathcal{M} with injectivity radius greater than or equal to $\varepsilon > 0$ is compact [20]. \square

A hyperbolic manifold obtained by performing Dehn filling on a hyperbolic manifold is said to be obtained by *hyperbolic Dehn filling* if the core of the filling torus is a geodesic—see section E.4 of [3]. Unlike the fillings considered in [22], [11], and [12], the filling on $M - K$ yielding M is typically not a hyperbolic Dehn filling—for example, we may take K to be null-homotopic in M . So, though $M - K$ admits a filling that moves the boundary, it could be that every hyperbolic Dehn filling leaves the hyperbolic structure on the boundary alone.

ACKNOWLEDGEMENTS

The author thanks his advisor Cameron Gordon, Jeff Brock, Chris Leininger, and Alan Reid for useful conversations. He thanks the referee for thoughtful comments.

REFERENCES

- [1] W. Abikoff, The real analytic theory of Teichmüller space, Lecture Notes in Math, vol. 820, Springer, 1980. MR0590044 (82a:32028)
- [2] L. V. Ahlfors, Lectures on quasiconformal mappings, Van Nostrand, 1966. MR0200442 (34:336)
- [3] R. Benedetti and C. Petronio, Lectures on hyperbolic geometry, Springer-Verlag, 1992. MR1219310 (94e:57015)
- [4] L. Bers, On boundaries of Teichmüller spaces and on Kleinian groups: I. Ann. of Math., 91 (1970), 570–600. MR0297992 (45:7044)
- [5] ———, Quasiconformal mappings and Teichmüller’s theorem, Analytic Functions (R. Nevanlinna et al., eds), 89–119. Princeton UP, 1960. MR0114898 (22:5716)
- [6] ———, Simultaneous uniformization, Bull. AMS 66 (1960) 94–97. MR0111834 (22:2694)
- [7] J. Birman, Mapping class groups and their relationship to braid groups, Comm. Pure and Applied Math. 22(1969) 213–238. MR0243519 (39:4840)
- [8] J. F. Brock, Iteration of mapping classes and limits of hyperbolic 3-manifolds, Invent. Math. 143, 523–570 (2001). MR1817644 (2002d:30052)
- [9] R. Brooks, Circle packings and co-compact extensions of Kleinian groups, Invent. Math. 86, 461–469 (1986). MR0860677 (88b:32050)

- [10] R. D. Canary, D. B. A. Epstein and P. Green, Notes on notes of Thurston. Analytical and geometric aspects of hyperbolic space (Coventry/Durham, 1984), 3–92, London Math. Soc. Lecture Note Ser., 111, Cambridge UP, 1987. MR0903850 (89e:57008)
- [11] M. Fujii, On totally geodesic boundaries of hyperbolic 3-manifolds, Kodai Math. J. 15 (1992), 244–257. MR1185422 (93i:57018)
- [12] M. Fujii and S. Kojima, Flexible boundaries in deformations of hyperbolic 3-manifolds, Osaka J. Math. 34 (1997), 541–551. MR1613080 (99a:57013)
- [13] M. Fujii and T. Soma, Totally geodesic boundaries are dense in the moduli space, J. Math. Soc. Japan, 49, no. 3, 1997, 589–601. MR1452704 (99b:57029)
- [14] F. P. Gardiner, Teichmüller theory and quadratic differentials, Pure and Applied Mathematics. A Wiley-Interscience Publication. John Wiley & Sons, Inc., New York, 1987. MR0903027 (88m:32044)
- [15] M. Kapovich, Hyperbolic Manifolds and Discrete Groups, Progress in Math. v. 183, Birkhäuser, 2000. MR1792613 (2002m:57018)
- [16] I. Kra, On the Nielsen–Thurston–Bers type of some self-maps of Riemann surfaces, Acta Math. 146 (1981) 231–270. MR0611385 (82m:32019)
- [17] B. Maskit, Kleinian groups, Grundlehren der mathematischen Wissenschaften 287, Springer, 1988. MR0959135 (90a:30132)
- [18] C. T. McMullen, Renormalization and 3-manifolds which fiber over the circle, Annals of Math. Studies no. 142, Princeton, 1996. MR1401347 (97f:57022)
- [19] J. W. Morgan, On Thurston’s uniformization theorem for three-dimensional manifolds, The Smith Conjecture, Academic Press, 1984, 37–125. MR0758464
- [20] D. Mumford, A remark on Mahler’s compactness theorem, Proc. AMS 28 (1971) 289–294. MR0276410 (43:2157)
- [21] R. Myers, Excellent 1-manifolds in compact 3-manifolds, Topology Appl. 49 (1993), no. 2, 115–127. MR1206219 (94a:57034)
- [22] W. D. Neumann and A. W. Reid, Rigidity of cusps in deformations of hyperbolic 3-orbifolds, Math. Ann. 295, 223–237 (1993). MR1202390 (94c:57025)
- [23] J. R. Stallings, Whitehead graphs on handlebodies. Geometric group theory down under (Canberra, 1996), 317–330, de Gruyter, Berlin, 1999. MR1714852 (2001i:57028)
- [24] J. H. C. Whitehead, On certain sets of elements in a free group. Proc. London Math. Soc. 41 (1936), 48–56.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TEXAS, AUSTIN, TEXAS 78712
E-mail address: rkent@math.utexas.edu