

$\mathcal{B}(\mathcal{H})$ IS A FREE SEMIGROUP ALGEBRA

KENNETH R. DAVIDSON

(Communicated by Joseph A. Ball)

ABSTRACT. We provide a simplified version of a construction of Charles Read. For any $n \geq 2$, there are n isometries with orthogonal ranges with the property that the nonselfadjoint weak- $*$ -closed algebra that they generate is all of $\mathcal{B}(\mathcal{H})$.

A free semigroup algebra is the weak operator topology closed algebra generated by an n -tuple of isometries with pairwise orthogonal ranges. The C^* -algebra generated by these isometries is called the Cuntz algebra \mathcal{O}_n , introduced in [4]. Cuntz algebras arise frequently in the theory of C^* -algebras as fundamental building blocks. They also play a crucial role in the analysis of endomorphisms of $\mathcal{B}(\mathcal{H})$ beginning with the seminal work of Powers [20]. Free semigroup algebras have been studied extensively in recent years [1, 5, 6, 7, 8, 9, 10, 11, 14, 16, 17, 18, 19]. In part, this is because of their intrinsically interesting properties as nonselfadjoint algebras. However it is also the case that these algebras reveal invariants of the corresponding C^* -algebra representations. As such, they have allowed the classification of certain families of representations of the Cuntz algebra [9, 7] which are relevant to recent work of Bratteli and Jorgensen, who use the Cuntz algebras to generate wavelets [2, 3, 12, 13].

In [9], we first raised the question of whether a free semigroup algebra could be a von Neumann algebra. In [6, 8], the special structure theory of these algebras has been revealed to a significant degree, and this question was seen as a central unresolved issue. Recently, Charles Read [21] constructed a representation of \mathcal{O}_2 for which the free semigroup is all of $\mathcal{B}(\mathcal{H})$. The purpose of this note is to provide a more transparent view of his construction.

1. THE CONSTRUCTION

For convenience, we will identify points in the unit interval I by their diadic expansion $x = 0.x_1x_2x_3\dots$ where $x_i \in \{0, 1\}$. Let \mathbb{F}_2^+ denote the free semigroup on two letters, which we will think of as the set of all words in symbols $\{0, 1\}$, including the empty word \emptyset , with multiplication being concatenation of words. Let $|w|$ denote the length of the word w . For each $w = i_1\dots i_k$ in \mathbb{F}_2^+ , define a map τ_w of I into itself by $\tau_w(x) = 0.i_1\dots i_kx_1x_2x_3\dots$ which is a homeomorphism of $[0, 1]$ onto the diadic interval I_w determined by w of length $2^{-|w|}$. Also define the function σ of I into itself by $\sigma(x) = 0.x_2x_3x_4\dots$. Observe that if $k = |w|$, then $\sigma^k\tau_w$ is the identity map, and $\tau_w\sigma^k$ is the identity restricted to I_w .

Received by the editors December 9, 2004 and, in revised form, January 20, 2005.

2000 *Mathematics Subject Classification*. Primary 47L80.

The author was partially supported by an NSERC grant.

©2005 American Mathematical Society
Reverts to public domain 28 years from publication

Let $\{h_i : i \geq 1\}$ denote the Walsh basis for $L^2(0, 1)$, which is obtained as products of the Rademacher functions; see Paley [15] or Zygmund [22, vol.1, p.34, Ex.6]. This is an orthonormal basis with the properties that each h_i takes only the values ± 1 , $h_1 = 1$, and $\text{span}\{h_i : i \leq 2^k\}$ is the set of functions which are constant on each dyadic interval of length 2^{-k} . Thus for each $i \leq 2^k$, there are constants $\mu_{i,w} \in \{\pm 1\}$ for words w with $|w| = k$ so that $h_i = \sum_{|w|=k} \mu_{i,w} \chi_{I_w}$.

For $i \geq 1$, let $b_i = 2^i$ and let $w_i = 011\dots 110$ denote the word of length b_i beginning and ending with a 0 and with $b_i - 2$ ones in between. Observe that the intervals I_{w_i} are pairwise disjoint. Define a function on I by

$$g(x) = \begin{cases} h_i(\sigma^{b_i}x) & \text{if } x \in I_{w_i}, \\ 1 & \text{otherwise.} \end{cases}$$

Note that $|g| = 1$ on I ; and $g = 1$ on I_1 . In particular, multiplication by $g\tau_0$ is unitary on $L^2(I)$; therefore, we can define a new orthonormal basis for $L^2(I)$ by

$$g_i(x) := g(\tau_0x)h_i(x) \quad \text{for } i \geq 1.$$

Define two isometries S_0 and S_1 by

$$S_i f(x) = \begin{cases} \sqrt{2}g(x)f(\sigma x) & \text{if } x \in I_i, \\ 0 & \text{otherwise.} \end{cases}$$

Observe that the range of S_i is $L^2(I_i)$ for $i = 0, 1$, and that these are complementary orthogonal subspaces. Since $g = 1$ on I_1 ,

$$S_1 f(x) = \sqrt{2} \chi_{I_1}(x)f(\sigma x).$$

For each $w = i_1 \dots i_k$ in \mathbb{F}_2^+ , write S_w for the product $S_{i_1} \dots S_{i_k}$. Observe that the isometry S_w carries $L^2(I)$ onto $L^2(I_w)$. In fact, this is the ‘standard’ isometry carrying $L^2(I)$ onto $L^2(I_w)$ multiplied by a certain function of modulus one.

We record a couple of useful identities as a lemma:

Lemma 1.1. *For each word $w \in \mathbb{F}_2^+$,*

$$(1.2) \quad S_w f(x) = 2^{|w|/2} G_{|w|}(x) \chi_{I_w}(x) f(\sigma^{|w|}x)$$

where $G_k(x) = \prod_{0 \leq j < k} g(\sigma^j x)$ is a function of modulus one, and

$$(1.3) \quad S_{w_j} g_i(x) = 2^{b_j/2} \chi_{I_{w_j}}(x) (h_j h_i)(\sigma^{b_j} x).$$

In particular, $S_{w_j} g_j = 2^{b_j/2} \chi_{I_{w_j}}$, and $S_{w_j} g_i$ is a function supported on I_{w_j} that has constant modulus and mean 0 when $i \neq j$.

Proof. The computation of $S_w f$ is routine. Note that

$$S_0 g_i(x) = \sqrt{2} \chi_{I_0}(x) g(\tau_0x)^2 h_i(\sigma x) = \sqrt{2} \chi_{I_0}(x) h_i(\sigma x).$$

Hence if we let v_j be the word w_j with the initial 0 truncated,

$$S_1^{b_j-2} S_0 g_i = 2^{(b_j-1)/2} \chi_{I_{v_j}} h_i(\sigma^{b_j-1} x).$$

Therefore

$$\begin{aligned} S_{w_j} g_i(x) &= 2^{(b_j-1)/2} S_0 \chi_{I_{v_j}} h_i(\sigma^{b_j-1} x) \\ &= 2^{b_j/2} \chi_{I_{w_j}}(x) (h_j h_i)(\sigma^{b_j} x). \end{aligned}$$

The properties of $S_{w_j} g_i$ are immediate. □

Definition 1.4. Let $\mathcal{G}_{i,k} = \{w : |w| = b_k - b_i \text{ and } G_{b_k-b_i}\tau_{ww_i} \text{ is constant}\}$. In this case, let $\lambda_{i,k,w}$ denote the value of $G_{b_k-b_i}$ on the interval I_{ww_i} .

Lemma 1.5. *If $k > i$, then*

$$1 - 2^{-6} < 2^{b_i-b_k} |\mathcal{G}_{i,k}| \leq 1.$$

Moreover, membership in $\mathcal{G}_{i,k}$ is not dependent on the first $b_k/4 = 2^{k-2}$ coefficients of w .

Proof. $G_{b_k-b_i}\tau_{ww_i}(x) = \prod_{0 \leq j < b_k-b_i} g(\sigma^j \tau_{ww_i} x)$ is a product of ± 1 's. The only time there is the possibility that $g(\sigma^j \tau_{ww_i} x)$ is not constant is if $\sigma^j \tau_{ww_i} x$ begins with a w_s . Since w_i begins with a 0 followed by b_i-2 ones, such a word is contained entirely within w or ends with the initial 0 of w_i . In particular, $s < k$ and $j \leq b_k - b_i - b_s + 1$. Moreover, the value of g will then be determined by the next $\lceil \log_2 s \rceil$ coefficients. This extends beyond the end of ww_i only if $j + b_s + \lceil \log_2 s \rceil > b_k$; whence

$$b_k - b_s - \lceil \log_2 s \rceil < j \leq b_k - b_i - b_s + 1.$$

This allows $\lceil \log_2 s \rceil + 1 - b_i$ possibilities for j when this number is positive—namely for $2^{b_i-1} < s < k$.

The probability that $\sigma^j \tau_{ww_i} x$ begins with w_s is 2^{-b_s} except in the case $j = b_k - b_i - b_s + 1$ when it is 2^{1-b_s} because the initial 0 of w_i is fixed. Thus the proportion of omitted words is at most

$$\begin{aligned} \sum_{2^{b_i-1} < s \leq k-1} (\lceil \log_2 s \rceil + 2 - b_i) 2^{-b_s} &\leq \sum_{s \geq 3} \lceil \log_2 s \rceil 2^{-b_s} \\ &= 2(2^{-8} + 2^{-16}) + 3(2^{-32} + \dots + 2^{-256}) + 4(2^{-512} + \dots + 2^{-65536}) + \dots \\ &< 2^{-6}. \end{aligned}$$

The smallest value of j which can achieve one of these inequalities is

$$b_k - b_{k-1} - \lceil \log_2(k-1) \rceil + 1 > 2^{k-2} = b_k/4.$$

So the first 2^{k-2} letters of w are irrelevant to determining membership in $\mathcal{G}_{i,k}$. \square

In view of this lemma, we may define $\mathcal{G}'_{i,k} = \sigma^{b_k/4} \mathcal{G}_{i,k}$ to be the set of tails of $\mathcal{G}_{i,k}$. Note that $|\mathcal{G}'_{i,k}| = 2^{-b_k/4} |\mathcal{G}_{i,k}|$ and

$$\mathcal{G}_{i,k} = \{vw' : |v| = b_k/4 \text{ and } w' \in \mathcal{G}'_{i,k}\}.$$

Lemma 1.6. *Let $i, j \in \mathbb{N}$. For each $k > \max\{i, j\}$, define*

$$T_{i,j,k} = |\mathcal{G}_{i,k}|^{-1/2} \sum_{w \in \mathcal{G}_{i,k}} \bar{\lambda}_{i,k,w} \mu_{j,w} S_{ww_i}.$$

Then $T_{i,j,k}$ is an isometry with the property that

$$\langle T_{i,j,k} g_{i'}, h_{j'} \rangle = \delta_{i'i'} \delta_{j'j'} 2^{-b_i/2} (2^{b_i-b_k} |\mathcal{G}_{i,k}|)^{1/2}$$

for all $i', j' \in \mathbb{N}$ with $j' \leq 2^{k-2}$.

Proof. By equation (1.3), we see that

$$S_{w_i} g_{i'} = 2^{b_i/2} \chi_{I_{w_i}}(x) (h_i h_{i'}) (\sigma^{b_i} x).$$

Hence by equation (1.2) for $w \in \mathcal{G}_{i,k}$,

$$\bar{\lambda}_{i,k,w} \mu_{j,w} S_{ww_i} g_{i'}(x) = 2^{b_k/2} \mu_{j,w} \chi_{I_{ww_j}}(x) (h_i h_{i'}) (\sigma^{b_k} x),$$

which is supported on $I_{ww_j} \subset I_w$. If $i' \neq i$, then since this function has mean zero for each $w \in \mathcal{G}_{i,k}$ and $h_{j'}$ is constant on the interval I_w , the inner product is 0.

Suppose that $i' = i$. Observe that $\mu_{j',vw'} = \mu_{j',v}$ depends on at most the first $2^{k-2} = b_k/4$ terms v of $w = vw'$. From the orthogonality of h_j and $h_{j'}$, we obtain $\sum_{|v|=b_k/4} \mu_{j,v} \mu_{j',v} = \delta_{jj'} 2^{b_k/4}$. Hence

$$\begin{aligned} \langle T_{i,j,k} g_i, h_{j'} \rangle &= |\mathcal{G}_{i,k}|^{-1/2} \sum_{w \in \mathcal{G}_{i,k}} 2^{b_k/2} \mu_{j,w} \mu_{j',w} \langle \chi_{I_{ww_j}}, \chi_{I_w} \rangle \\ &= |\mathcal{G}_{i,k}|^{-1/2} \sum_{|v|=b_k/4} \mu_{j,v} \mu_{j',v} \sum_{w' \in \mathcal{G}'_{i,k}} 2^{b_k/2} \langle \chi_{I_{vw'w_j}}, \chi_{I_{vw'}} \rangle \\ &= |\mathcal{G}_{i,k}|^{-1/2} \delta_{jj'} 2^{b_k/4} |\mathcal{G}'_{i,k}| 2^{-b_k} \\ &= \delta_{jj'} 2^{-b_i/2} (2^{b_i-b_k} |\mathcal{G}_{i,k}|)^{1/2}. \quad \square \end{aligned}$$

Theorem 1.7. *For each $s \geq 1$, the weak-* closed algebra generated by $\{S_w : |w| = 2^s\}$ is all of $\mathcal{B}(\mathcal{H})$.*

Proof. The previous lemma exhibits a bounded sequence of operators $T_{i,j,k}$ built from words S_{ww_i} of length 2^k which converge in the weak operator topology to a positive multiple of the rank one operator $h_j g_i^*$. Hence the limit also exists in the weak-* topology. For $k \geq s$, these operators lie in the desired algebra. These limit operators span the set of compact operators, which is weak-* dense in $\mathcal{B}(\mathcal{H})$. \square

Corollary 1.8. *For every $n \geq 2$, $\mathcal{B}(\mathcal{H})$ is a free semigroup algebra for a representation of \mathcal{O}_n .*

Proof. Between the algebras \mathcal{O}_{2^k} and $\mathcal{O}_{2^{k+1}}$ generated by words S_w where w has length 2^k and 2^{k+1} respectively, there is a copy of \mathcal{O}_n for $2^k < n < 2^{k+1}$. For example, \mathcal{O}_3 is generated by S_{00}, S_{01} and S_1 . Use the representation constructed above. \square

Remark 1.9. Since Lemma 1.6 is valid for all i' , it follows that

$$T_{i,j,k}^* h_{j'} = \delta_{jj'} \lambda_{i,k} g_i$$

where $\lambda_{i,k} = 2^{-b_i/2} (2^{b_i-b_k} |\mathcal{G}_{i,k}|)^{1/2}$. So these adjoints actually converge in the strong operator topology.

REFERENCES

[1] A. Arias and G. Popescu, *Factorization and reflexivity on Fock spaces*, Integral Equations. Operator Theory **23** (1995), 268–286. MR1356335 (97e:47066)
 [2] O. Bratteli and P. Jorgensen, *Iterated function systems and permutation representations of the Cuntz algebra*, Mem. Amer. Math. Soc. **139** (1999), no. 663. MR1469149 (99k:46094a)
 [3] O. Bratteli and P. Jorgensen, *Wavelet filters and infinite dimensional unitary groups*, Wavelet analysis and applications (Guangzhou, 1999), 35–65, AMS/IP Stud. Adv. Math. **25**, Amer. Math. Soc., Providence, RI, 2002. MR1887500 (2003e:94015)
 [4] J. Cuntz, *Simple C^* -algebras generated by isometries*, Comm. Math. Phys. **57** (1977), 173–185. MR0467330 (57:7189)
 [5] K.R. Davidson, *Free semigroup algebras, a survey*, Systems, approximation, singular integral operators, and related topics (Bordeaux, 2000), 209–240, Oper. Theory Adv. Appl., **129**, Birkhauser, Basel, 2001. MR1882697 (2003d:47104)
 [6] K.R. Davidson, E. Katsoulis and D.R. Pitts, *The structure of free semigroup algebras*, J. reine angew. Math. (Crelle) **533** (2001), 99–125. MR1823866 (2002a:47107)

- [7] K.R. Davidson, D.W. Kribs and M.E. Shpigel, *Isometric dilations of non-commuting finite rank n -tuples*, *Canad. J. Math.* **53** (2001), 506–545. MR1827819 (2002f:47010)
- [8] K.R. Davidson, J. Li and D.R. Pitts, *A Kaplansky Density Theorem for Free Semigroup Algebras*, *J. Func. Anal.*, to appear.
- [9] K.R. Davidson and D.R. Pitts, *Invariant subspaces and hyper-reflexivity for free semi-group algebras*, *Proc. London Math. Soc.* **78** (1999), 401–430. MR1665248 (2000k:47005)
- [10] K.R. Davidson and D.R. Pitts, *The algebraic structure of non-commutative analytic Toeplitz algebras*, *Math. Ann.* **311** (1998), 275–303. MR1625750 (2001c:47082)
- [11] K.R. Davidson and D.R. Pitts, *Nevanlinna–Pick Interpolation for non-commutative analytic Toeplitz algebras*, *Integral Equations Operator Theory* **31** (1998), 321–337. MR1627901 (2000g:47016)
- [12] P. Jorgensen, *Minimality of the data in wavelet filters*, *Adv. Math.* **159** (2001), 143–228. MR1825057 (2002h:46092)
- [13] P. Jorgensen and D. Kribs, *Wavelet representations and Fock space on positive matrices*, *J. Func. Anal.* **197** (2003), 526–559. MR1960424 (2004e:42055)
- [14] D.W. Kribs, *Factoring in non-commutative analytic Toeplitz algebras*, *J. Operator Theory* **45** (2001), 175–193. MR1823067 (2002g:47151)
- [15] R.E.A.C. Paley, *A remarkable system of orthogonal functions*, *Proc. London Math. Soc.* **34** (1932), 241–279.
- [16] G. Popescu, *Isometric dilations for infinite sequences of noncommuting operators*, *Trans. Amer. Math. Soc.* **316** (1989), 523–536. MR0972704 (90c:47006)
- [17] G. Popescu, *Characteristic functions for infinite sequences of noncommuting operators*, *J. Operator Theory* **22** (1989), 51–71. MR1026074 (91m:47012)
- [18] G. Popescu, *Multi-analytic operators and some factorization theorems*, *Indiana Univ. Math. J.* **38** (1989), 693–710. MR1017331 (90k:47019)
- [19] G. Popescu, *Multi-analytic operators on Fock spaces*, *Math. Ann.* **303** (1995), 31–46. MR1348353 (96k:47049)
- [20] R.T. Powers, *An index theory for semigroups of $*$ -endomorphisms of $\mathcal{B}(\mathcal{H})$ and type II_1 factors*, *Canad. J. Math.* **40** (1988), 86–114. MR0928215 (89f:46116)
- [21] C. Read, *A large weak operator closure for the algebra generated by two isometries*, *Journal of Operator Theory*, to appear.
- [22] A. Zygmund, *Trigonometric Series*, 2nd ed., Cambridge University Press, Cambridge, 1959, reprinted 1977. MR0617944 (58:29731)

DEPARTMENT OF PURE MATHEMATICS, UNIVERSITY OF WATERLOO, WATERLOO, ONTARIO,
N2L 3G1 CANADA

E-mail address: krdaids@uwaterloo.ca