

PERTURBATION THEORETIC ENTROPY OF THE BOUNDARY ACTIONS OF FREE GROUPS

RUI OKAYASU

(Communicated by David R. Larson)

ABSTRACT. We compute the exact value of Voiculescu’s perturbation theoretic entropy of the boundary actions of free groups. This result is a partial answer of Voiculescu’s question.

1. INTRODUCTION

In the serial papers [Voi1, Voi2, Voi3] on perturbations of Hilbert space operators, D. Voiculescu studied a certain numerical invariant $k_{\infty}^{-}(\tau)$ for a tuple τ of operators. This number can be viewed as a kind of dimension of τ with respect to the Macaev ideal (see [Voi1] and [DV]). In [Oka1, Oka2], we computed the exact value of $k_{\infty}^{-}(\tau)$ for a family τ of operators in the following two classes: (1) creation operators associated with a subshift, which were introduced by K. Matsumoto [Mat], (2) unitary operators in the left regular representation of a finitely generated group.

In [Voi4, Voi5], Voiculescu defined the perturbation theoretic entropy $H_P(\Omega, \mathfrak{M})$ for a von Neumann algebra \mathfrak{M} and a set Ω of unitary operators normalizing \mathfrak{M} . Thanks to the theory of the standard form of von Neumann algebras, the perturbation theoretic entropy can be defined even for a tuple of automorphisms of a von Neumann algebra without assuming the existence of invariant states. In the case where $\mathfrak{M} = L^{\infty}(X)$ and Ω consists of the unitary operator U_T induced by an ergodic measure preserving transform T on a probability space (X, μ) , he obtained that

$$\frac{1}{2}h(T) \leq H_P(U_T, L^{\infty}(X)) \leq 6h(T),$$

and moreover if T is Bernoulli, then $H_P(U_T, L^{\infty}(X))$ is proportional to $h(T)$, where $h(T)$ is Kolmogorov-Sinai entropy. These results suggest that the perturbation theoretic entropy has a certain relation to Connes-Størmer entropy [CS]. However for the free shift σ on the free group factor $L(\mathbb{F}_{\infty})$, it is quite different. Indeed, Connes-Størmer entropy $H(\sigma) = 0$, but the perturbation theoretic entropy $H_P(\sigma, L(\mathbb{F}_{\infty})) = +\infty$. Thus the perturbation theoretic entropy may be a better concept of entropy in some highly non-commutative cases. We also mention that the perturbation theoretic entropy is well defined for transforms with quasi-invariant measures and depends only on the class of the measure. However, it is an open

Received by the editors October 25, 2004 and, in revised form, January 25, 2005.
2000 *Mathematics Subject Classification*. Primary 46L55; Secondary 28D20, 47B37.
Key words and phrases. Perturbation, entropy, Poisson boundary, free group.

problem whether $H_P(U_T, L^\infty(X))$ can be non-zero and finite in the absence of an invariant measure.

We consider in the present paper the boundary actions of free groups. If $\partial\mathbb{F}_N$ is the boundary of the free group \mathbb{F}_N with the canonical symmetric generating set S , then using the similar technique of [Oka1], we obtain the lower bound

$$H_P(\Omega_N, L^\infty(\partial\mathbb{F}_N)) \geq \log(2N - 1),$$

where $\Omega_N = \{U_a\}_{a \in S}$, U_a is the corresponding unitary on $L^2(\partial\mathbb{F}_N, \mu)$ and μ is a certain harmonic measure on $\partial\mathbb{F}_N$. Unfortunately we cannot get the upper bound of $H_P(\Omega_N, L^\infty(\partial\mathbb{F}_N))$. However if we replace $L^\infty(\partial\mathbb{F}_N)$ by the smaller algebra $C(\partial\mathbb{F}_N)$, then we obtain the equation

$$H_P(\Omega_N, C(\partial\mathbb{F}_N)) = \log(2N - 1).$$

This result is a partial answer of the above-mentioned problem because there is no invariant measure on $\partial\mathbb{F}_N$ under the action of \mathbb{F}_N . We also remark that this number coincides with the topological entropy $h(\mathbb{F}_N; \partial\mathbb{F}_N)$ (see e.g. [BW]).

2. PRELIMINARIES

We quickly introduce several definitions and results of [Voi3, Voi4, Voi5], which we will use. The Macaev ideal $(\mathcal{C}_\infty^-(H), \|\cdot\|_\infty^-)$ in the C^* -algebra $\mathbb{B}(H)$ of bounded operators on a separable infinite-dimensional Hilbert space H is defined by the norm

$$\|T\|_\infty^- = \sum_{j=1}^\infty s_j(T)j^{-1},$$

where $s_1(T) \geq s_2(T) \geq \dots$ are the eigenvalues of $(T^*T)^{1/2}$. Its dual is denoted by $(\mathcal{C}_1^+(H), \|\cdot\|_1^+)$. The set $\mathbb{F}(H)$ of finite rank operators is dense in $\mathcal{C}_\infty^-(H)$ but not in $\mathcal{C}_1^+(H)$.

If $\tau = (T_1, \dots, T_N)$ with $T_k \in \mathbb{B}(H)$ ($1 \leq k \leq N$), the invariant $k_\infty^-(\tau)$ is defined by

$$k_\infty^-(\tau) = \liminf_{A \in \mathbb{F}(H)_1^+} \max_{1 \leq i \leq N} \|[A, T_i]\|_\infty^-,$$

where $\mathbb{F}(H)_1^+ = \{A \in \mathbb{F}(H) \mid 0 \leq A \leq I\}$ with the natural order. We will evaluate $k_\infty^-(\tau)$ using the following facts.

Fact 1. If $A_m \in \mathbb{F}(H)_1^+$ with $A_m \uparrow I$, then we have

$$k_\infty^-(\tau) \leq \liminf_{m \rightarrow \infty} \max_{1 \leq i \leq N} \|[A_m, T_i]\|_\infty^-.$$

Fact 2. Let $\{X_i\}_{1 \leq i \leq N} \subset \mathcal{C}_1^+(H)$. If $\sum_{i=1}^N [T_i, X_i] \in \mathcal{C}_1(H) + \mathbb{B}(H)_+$, then we have

$$\left| \text{Tr} \left(\sum_{i=1}^N [T_i, X_i] \right) \right| \leq k_\infty^-(\tau) \sum_{i=1}^N \|X_i\|_1^{\tilde{+}},$$

where $\mathcal{C}_1(H)$ is the trace class ideal and

$$\|X_i\|_1^{\tilde{+}} = \inf_{Y \in \mathbb{F}(H)} \|X_i + Y\|_1^+.$$

Remark 1. Note that for $X \in \mathcal{C}_1^+(H)$, we have

$$\|X\|_1^{\tilde{+}} = \limsup_{n \rightarrow \infty} \frac{\sum_{j=1}^n s_j(X)}{\sum_{j=1}^n j^{-1}}.$$

Moreover if an operator X has singular values $c\alpha^{-n}$ with multiplicities $d\alpha^n$ ($n \geq 1$), where some constant $c > 0$ and positive integers d, α , then one can obtain that

$$\|X\|_1^{\tilde{+}} = \frac{cd}{\log \alpha}.$$

Definition 2 (cf. [Voi4], [Voi6]). Let \mathfrak{A} be a C^* -algebra acting on a Hilbert space H and let Ω be a set of unitaries on H normalizing \mathfrak{A} , i.e., $U\mathfrak{A}U^* = \mathfrak{A}$ for $U \in \Omega$. The perturbation theoretic entropy $H_P(\Omega, \mathfrak{A})$ is defined by

$$H_P(\Omega, \mathfrak{A}) = \sup\{k_{\infty}^-(\omega \cup \tau) \mid \omega \in \mathcal{F}(\Omega), \tau \in \mathcal{F}(\mathfrak{A}), \dim C^*(\tau) < +\infty\},$$

where $\mathcal{F}(\Omega)$ and $\mathcal{F}(\mathfrak{A})$ are sets of all finite subsets of Ω and \mathfrak{A} , respectively.

Remark 3. The above definition is slightly different from the original one in [Voi4]. Our definition is introduced in [Voi6]. One can easily check that all results in [Voi4] and [Voi5] can be also obtained if the above definition is adopted. We believe that our definition is better, because we do not need to deal with the invariant $k_{\infty}^-(\tau)$ for a family τ of infinitely many operators. We also remark that if we consider the original definition, then our main result can be obtained by the same proof.

We note that $H_P(\Omega, \mathfrak{A}) = +\infty$ unless $k_{\infty}^-(\tau) = 0$ for all $\tau \in \mathcal{F}(\mathfrak{A})$ with $\dim C^*(\tau) < +\infty$. It is clear that if $\mathfrak{M} = \mathfrak{A}''$, then we have that

$$H_P(\Omega, \mathfrak{M}) \geq H_P(\Omega, \mathfrak{A}).$$

3. RESULTS

Let $\mathbb{F}_N = \langle a_1, \dots, a_N \rangle$ be the free group with the canonical generating set $S = \{a_1^{\pm 1}, \dots, a_N^{\pm 1}\}$ and $\partial\mathbb{F}_N$ its boundary with the product topology,

$$\partial\mathbb{F}_N = \{(x_n)_{n=1}^{\infty} \mid x_n \in S, x_{n+1} \neq x_n^{-1}\}.$$

Note that \mathbb{F}_N acts on $\partial\mathbb{F}_N$ by left multiplications. We always assume that the expression such as $x_1 \cdots x_n$ is reduced throughout this paper. We define the probability measure μ on $\partial\mathbb{F}_N$ by

$$\mu(C(x_1 \cdots x_n)) = \frac{1}{2N(2N-1)^{n-1}},$$

where $C(x_1 \cdots x_n)$ is the cylinder set. We consider the C^* -algebra $C(\partial\mathbb{F}_N)$ acting on the Hilbert space $L^2(\partial\mathbb{F}_N, \mu)$ by left multiplications and the unitary operator U_g on $L^2(\partial\mathbb{F}_N, \mu)$ for $g \in \mathbb{F}_N$ given by

$$(U_g f)(\mathbf{x}) = f(g^{-1}\mathbf{x}) \sqrt{\frac{dg_*\mu}{d\mu}}(\mathbf{x}),$$

for $f \in L^2(\partial\mathbb{F}_N, \mu)$. Note that if $\mathbf{x} = (x_n)_{n=1}^{\infty} \in \partial\mathbb{F}_N$, we have

$$\frac{dg_*\mu}{d\mu}(\mathbf{x}) = (2N-1)^{|g_n| - |g^{-1}g_n|}$$

for sufficiently large n , where $g_n = x_1 \cdots x_n$ and $|\cdot|$ is the word-length on \mathbb{F}_N . We denote by $\chi_{x_1 \cdots x_n}$ the characteristic function of $C(x_1 \cdots x_n)$. We remark that for $a \in S$ and $n \geq 2$,

$$U_a \chi_{x_1 \cdots x_n} = \begin{cases} c_N \chi_{ax_1 \cdots x_n} & \text{if } x_1 \neq a^{-1}, \\ c_N^{-1} \chi_{x_2 \cdots x_n} & \text{if } x_1 = a^{-1}, \end{cases}$$

where $c_N = \sqrt{2N - 1}$. Let $\Omega_N = \{U_a\}_{a \in S}$.

Proposition 4.

$$H_P(\Omega_N, C(\partial\mathbb{F}_N)) \geq \log(2N - 1).$$

Hence,

$$H_P(\Omega_N, L^\infty(\partial\mathbb{F}_N)) \geq \log(2N - 1).$$

Proof. We set $H_0 = \mathbb{C}\chi$ where $\chi(\mathbf{x}) = 1$ for $\mathbf{x} \in \partial\mathbb{F}_N$ and for $n \geq 1$,

$$H_n = \text{span}\{\chi_{x_1 \cdots x_n}\}.$$

Note that $H_0 \subset H_1 \subset \cdots \subset H_n \subset \cdots \subset H = L^2(\partial\mathbb{F}_N, \mu)$ and $\dim H_n = 2N(2N - 1)^{n-1}$ for $n \geq 1$. We also define the subspace $K_n = H_n \cap H_{n-1}^\perp$ for $n \geq 1$. One can easily check that

$$K_n = \text{span}\{\chi_{x_1 \cdots x_{n-1}x} - \chi_{x_1 \cdots x_{n-1}y}\}$$

with $\dim K_1 = 2N - 1$ and $\dim K_n = \dim H_n - \dim H_{n-1} = 4N(N - 1)(2N - 1)^{n-2}$ for $n \geq 2$.

We denote by P_n the projection onto K_n . For $a \in S$ we define the operator $X_a \in \mathbb{B}(H)$ by

$$X_a = \sum_{n \geq 1} \sum_{x_1, \dots, x_n} \frac{1}{2N(2N - 1)^{n-1}} P_n U_a^* P_{n+1}.$$

For $a \in S$ and $n \geq 2$, let us denote by $P_n(a)$ the projection onto

$$K_n(a) = \text{span}\{\chi_{x_1 \cdots x_{n-1}x} - \chi_{x_1 \cdots x_{n-1}y} \mid x_1 = a\}.$$

Then it is easy to check that

$$\sum_{a \in S} P_n(a) = P_n$$

and $\dim K_n(a) = 2(N - 1)(2N - 1)^{n-2}$ for $a \in S$. Moreover $P_n U_a^* P_{n+1} = (P_n - P_n(a^{-1}))U_a^*$ for $n \geq 2$ and $P_n U_a^* P_{n+1} = U_a^* P_{n+1}(a)$ for $n \geq 1$. It follows that

$$\sum_{a \in S} U_a X_a = \sum_{n \geq 1} \frac{1}{2N(2N - 1)^{n-1}} P_{n+1},$$

and

$$\sum_{a \in S} X_a U_a = \frac{1}{2N} \sum_{a \in S} U_a^* P_2(a) U_a + \sum_{n \geq 2} \frac{1}{2N(2N - 1)^{n-2}} P_n.$$

Therefore,

$$\left| \text{Tr} \left(\sum_{a \in S} [U_a, X_a] \right) \right| = \frac{1}{2N} \text{Tr} \left(\sum_{a \in S} U_a^* P_2(a) U_a \right) = \frac{1}{2N} \text{Tr}(P_2) = 2(N - 1).$$

For $a \in S$, we next estimate the norm $\|X_a\|_\infty^\pm$. By using the fact in Remark 1 as

$$c = \frac{1}{2N}, d = 2(N - 1), \alpha = 2N - 1,$$

we obtain that

$$\|X_a\|_1^{\tilde{\tau}} = \frac{2(N-1)}{2N \log(2N-1)}.$$

Hence it follows from Fact 2 that

$$H_P(\Omega_N, C(\partial\mathbb{F}_N)) \geq k_{\infty}^-(\Omega_N) \geq \log(2N-1).$$

□

Theorem 5.

$$H_P(\Omega_N, C(\partial\mathbb{F}_N)) = \log(2N-1).$$

Proof. We define the finite-dimensional C^* -subalgebra \mathfrak{A}_n of $\mathfrak{A} = C(\partial\mathbb{F}_N)$ by $\mathfrak{A}_0 = \mathbb{C}\chi$ and

$$\mathfrak{A}_n = \overline{\text{span}}\{\chi_{x_1 \cdots x_n}\}$$

for $n \geq 1$. Note that $\dim \mathfrak{A}_0 = 1, \dim \mathfrak{A}_n = 2N(2N-1)^{n-1}$ and

$$\mathfrak{A} = \overline{\bigcup_{n \geq 0} \mathfrak{A}_n}.$$

We denote by P_n the projection onto the subspace $H_n = \mathfrak{A}_n\chi \subset L^2(\partial\mathbb{F}_N, \mu)$. Note that for $a \in S$ and $n \geq 2$,

$$P_{n-1} \leq U_a P_n U_a^* \leq P_{n+1}.$$

Let $\tau \in \mathcal{F}(\mathfrak{A})$ with $\dim C^*(\tau) < +\infty$. Note that $C^*(\tau)$ is contained in \mathfrak{A}_K for some $K > 1$. Then we define

$$Q_m = \frac{1}{m} \sum_{n=K}^{K+m-1} P_n.$$

Since

$$U_a Q_m U_a^* - Q_m = \frac{1}{m} \sum_{n=K}^{K+m-1} (U_a P_n U_a^* - P_n) \leq \frac{1}{m} \sum_{n=K}^{K+m-1} (P_{n+1} - P_n) \leq \frac{1}{m} P_{K+m}$$

and

$$U_a Q_m U_a^* - Q_m \geq \frac{1}{m} \sum_{n=K}^{K+m-1} (P_{n-1} - P_n) \geq -\frac{1}{m} P_{K+m},$$

we obtain that

$$U_a Q_m U_a^* - Q_m = \frac{1}{m} P_{K+m} C_m P_{K+m}$$

for some contraction C_m . Since $P_n \uparrow I$, we also have that $Q_m \uparrow I$. Moreover each Q_m commutes with τ . Therefore we have

$$\liminf_{m \rightarrow \infty} \|U_a Q_m U_a^* - Q_m\|_{\infty}^- \leq \lim_{m \rightarrow \infty} \frac{1}{m} \log(2N(2N-1)^{K+m-1}) = \log(2N-1),$$

and our assertion follows from Fact 1 and Proposition 2. □

Remark 6. We also prove that $k_{\infty}^-(\Omega_N) = \log(2N-1)$, actually. We wonder how this is related to our previous result $k_{\infty}^-(\lambda_N) = \log(2N-1)$ in [Oka1] for $\lambda_N = \{\lambda_a\}_{a \in S}$, where λ is the left regular representation of \mathbb{F}_N , because Ω_N and λ_N generate the same C^* -algebra, although they are not unitarily equivalent. (See, e.g., [FTP] for the representation U of \mathbb{F}_N arising from the Poisson boundary.)

ACKNOWLEDGMENT

The author would like to thank the referee for valuable suggestions and comments.

REFERENCES

- [BW] Biś, A.; Walczak, P. G. *Entropies of hyperbolic groups and some foliated spaces*. Foliations: geometry and dynamics (Warsaw, 2000), 197–211, World Sci. Publishing, River Edge, NJ, 2002. MR1882770 (2002k:37038)
- [CS] Connes, A.; Størmer, E. *Entropy for automorphisms of II_1 von Neumann algebras*. Acta Math. **134** (1975), no. 3-4, 289–306. MR0454657 (56:12906)
- [DV] David, G.; Voiculescu, D. *s-numbers of singular integrals for the invariance of absolutely continuous spectra in fractional dimensions*. J. Funct. Anal. **94** (1990), no. 1, 14–26. MR1077543 (92f:47014)
- [FTP] Figà -Talamanca, A.; Picardello, M. A. *Spherical functions and harmonic analysis on free groups*. J. Funct. Anal. **47** (1982), no. 3, 281–304. MR0665019 (83m:22018)
- [Mat] Matsumoto, K. *On C^* -algebras associated with subshifts*. Internat. J. Math. **8** (1997), no. 3, 357–374. MR1454478 (98h:46077)
- [Oka1] Okayasu, R. *Entropy of Subshifts and the Macaev Norm*. J. Math. Soc. Japan **56** (2004), no. 1, 177–191. MR2027621 (2004j:47137)
- [Oka2] Okayasu, R. *Gromov hyperbolic groups and the Macaev Norm*. Preprint.
- [Voi1] Voiculescu, D. *Some results on norm-ideal perturbations of Hilbert space operators*. J. Operator Theory **2** (1979), no. 1, 3–37. MR0553861 (80m:47012)
- [Voi2] Voiculescu, D. *Some results on norm-ideal perturbations of Hilbert space operators. II*. J. Operator Theory **5** (1981), no. 1, 77–100. MR0613049 (83f:47014)
- [Voi3] Voiculescu, D. *On the existence of quascentral approximate units relative to normed ideals. Part I*. J. Funct. Anal. **91** (1990), no. 1, 1–36. MR1054113 (91m:46089)
- [Voi4] Voiculescu, D. *Entropy of dynamical systems and perturbations of operators*. Ergodic Theory Dynam. Systems **11** (1991), no. 4, 779–786. MR1145622 (93b:46130)
- [Voi5] Voiculescu, D. *Entropy of dynamical systems and perturbations of operators. II*. Houston J. Math. **17** (1991), no. 4, 651–661. MR1147278 (93b:46131)
- [Voi6] Voiculescu, D. *Entropy-invariants of dynamical systems and perturbations of operators*. Mathematical physics, X (Leipzig, 1991), 303–307, Springer, Berlin, 1992. MR1386418

DEPARTMENT OF MATHEMATICS EDUCATION, OSAKA KYOIKU UNIVERSITY, KASHIWARA, OSAKA 582-8582, JAPAN

E-mail address: rui@cc.osaka-kyoiku.ac.jp