

## SIMPLE REAL RANK ZERO ALGEBRAS WITH LOCALLY HAUSDORFF SPECTRUM

PING WONG NG

(Communicated by David R. Larson)

ABSTRACT. Let  $\mathcal{A}$  be a unital, simple, separable  $C^*$ -algebra with real rank zero, stable rank one, and weakly unperforated ordered  $K_0$  group. Suppose, also, that  $\mathcal{A}$  can be locally approximated by type I algebras with Hausdorff spectrum and bounded irreducible representations (the bound being dependent on the local approximating algebra). Then  $\mathcal{A}$  is tracially approximately finite dimensional (i.e.,  $\mathcal{A}$  has tracial rank zero).

Hence,  $\mathcal{A}$  is an  $AH$ -algebra with bounded dimension growth and is determined by  $K$ -theoretic invariants.

The above result also gives the first proof for the locally  $AH$  case.

### 1. INTRODUCTION

In the  $K$ -theoretic classification program for simple unital separable stably finite nuclear  $C^*$ -algebras, a great deal of progress has been made for those algebras which have stable rank one, real rank zero, and weak unperforation in the ordered  $K_0$ -group (see, for example, [7], [13], [4], [1] and the last paragraph of [10]). One of the fundamental results in this direction is the work in [7], where Elliott and Gong classified (using  $K$ -theoretic invariants) all simple unital  $AH$ -algebras with bounded dimension growth and real rank zero.

We note that the class of algebras in [7] exhausts the current invariant for simple unital stably finite real rank zero nuclear  $C^*$ -algebras. Much work to date has been done to give classification results, for simple, nuclear, stably finite, real rank zero algebras, that do not assume that the  $C^*$ -algebras involved are  $AH$ -algebras (see, for example, [13] and the references therein).

**Definition 1.1.** Let  $\mathcal{A}$  be a simple unital  $C^*$ -algebra. Then  $\mathcal{A}$  is said to be tracially approximately finite dimensional (abbreviated by “TAF”) if for every  $\epsilon > 0$ , for every finite subset  $\mathcal{F}$  of  $\mathcal{A}$  and for every strictly positive element  $a \in \mathcal{A}$ , there is a projection  $p$  which is Murray-von Neumann equivalent to a subprojection in the hereditary subalgebra generated by  $a$  and there exists a finite dimensional  $C^*$ -subalgebra  $\mathcal{B}$  of  $\mathcal{A}$  such that: (a)  $1_{\mathcal{A}} - p = 1_{\mathcal{B}}$  where  $1_{\mathcal{B}}$  is the unit of  $\mathcal{B}$ , (b)  $\|xp - px\| < \epsilon$  for every  $x \in \mathcal{F}$ , and (c)  $(1_{\mathcal{A}} - p)x(1_{\mathcal{A}} - p)$  is within  $\epsilon$  of an element of  $\mathcal{B}$ , for every  $x \in \mathcal{F}$ .

The term “tracial rank zero” is often used in place of “tracially approximately finite dimensional” (see, for instance, [12] and [13]). Hence, by definition, a simple,

---

Received by the editors November 21, 2003 and, in revised form, June 23, 2004.  
2000 *Mathematics Subject Classification.* Primary 46L35.

©2006 American Mathematical Society  
Reverts to public domain 28 years from publication  
2223

unital  $C^*$ -algebra is *TAF* if and only if it has tracial rank zero. (Indeed, Lin has a notion of *tracial rank*, which takes on values other than zero. For example, all nonreal rank zero, simple, unital *AH*-algebras, with bounded dimension growth, have tracial rank one; see [12].)

Lin has shown that the class of simple unital separable nuclear *TAF* algebras which satisfy the universal coefficient theorem is exactly the class of [7] (see [13]).

**Definition 1.2.** Let  $\mathcal{A}$  be a  $C^*$ -algebra. (a) Then  $\mathcal{A}$  is said to be locally type I if for every  $\epsilon > 0$ , for every finite subset  $\mathcal{F}$  of  $\mathcal{A}$ , there is a separable type I  $C^*$ -subalgebra  $\mathcal{B}$  of  $\mathcal{A}$  such that every element of  $\mathcal{F}$  is within  $\epsilon$  of an element of  $\mathcal{B}$ . (b) If in (a), every (local approximating) type I  $C^*$ -algebra  $\mathcal{B}$  has Hausdorff spectrum and there exists an integer  $L$  (dependent on  $\mathcal{B}$ ) such that every irreducible representation of  $\mathcal{B}$  has dimension less than  $L$ , then  $\mathcal{A}$  is said to have locally Hausdorff spectrum. (c) If in (a), every (locally approximating) type I  $C^*$ -algebra has the form  $\bigoplus_{i=1}^N p_i \mathbb{M}_{n_i}(C(X_i)) p_i$ , where each  $X_i$  is a compact metric space and each  $p_i$  is a projection in  $\mathbb{M}_{n_i}(C(X_i))$ , then  $\mathcal{A}$  is said to be locally *AH*.

We note that Dadarlat and Eilers have given an example of a (nonsimple) separable, unital, locally *AH*  $C^*$ -algebra which has real rank zero and stable rank one, but is not an *AH*-algebra (see [3]).

We also note that in [2], Dadarlat has shown that if  $\mathcal{A}$  is a separable nuclear  $C^*$ -algebra which can be locally approximated by  $C^*$ -algebras which satisfy the universal coefficient theorem, then  $\mathcal{A}$  also satisfies the universal coefficient theorem. Hence, every locally type I  $C^*$ -algebra satisfies the universal coefficient theorem.

In [15], Lin proved the following very interesting result (there are several proofs in the literature; other proofs can be found in [1, Corollary 7.11], [14] and [17, Theorem 5.16]):

**Theorem 1.3.** *Let  $\mathcal{A}$  be a unital separable simple locally type I  $C^*$ -algebra with real rank zero, stable rank one, weak unperforation in the  $K_0$ -group. Suppose also that the tracial simplex has countably many extreme points. Then  $\mathcal{A}$  is *TAF*. By a theorem of Lin, this implies that  $\mathcal{A}$  is an *AH*-algebra with bounded dimension growth and is determined by  $K$ -theoretic invariants.*

We note that Lin's result requires a restriction on the tracial simplex of  $\mathcal{A}$  (countably many extreme points). There have also been other interesting results in the literature which require this restriction on the tracial simplex (see, for example [1], the last paragraph of [10], [14], [15] and [17]).

In this paper, we remove the unique trace condition in Lin's result provided that the (local) type I algebras have Hausdorff spectrum and bounded irreducible representations.

**Definition 1.4.**  $\mathcal{LCH}^+$  is the class of simple unital separable  $C^*$ -algebras with real rank zero, stable rank one, weak unperforation in the ordered  $K_0$ -group, and having locally Hausdorff spectrum.

**Theorem 1.5.** *Let  $\mathcal{A}$  be a  $C^*$ -algebra in  $\mathcal{LCH}^+$ . Then  $\mathcal{A}$  is *TAF*. Hence, by a theorem of Lin,  $\mathcal{A}$  is an *AH*-algebra with bounded dimension growth and is determined by  $K$ -theoretic invariants.*

Our result gives the first proof that a simple unital separable locally *AH*  $C^*$ -algebra with real rank zero, stable rank one, weak unperforation in the  $K_0$  group is *TAF*, without any restriction on the tracial simplex.

A modification of our argument gives a short alternative proof of the following result of Lin (which also follows from our result).

**Theorem 1.6** (see [16]). *Let  $\mathcal{A}$  be a simple unital AH-algebra which has stable rank one, real rank zero and weakly unperforated  $K_0$  group. Then  $\mathcal{A}$  is TAF.*

Note that in the hypothesis of the above result, it is not assumed that  $\mathcal{A}$  has bounded dimension growth. Also, Lin’s argument does not generalize to the locally AH case.

In what follows, if  $\mathcal{A}$  is a unital  $C^*$ -algebra, then  $T(\mathcal{A})$  is the simplex of unital traces on  $\mathcal{A}$ .

2. MAIN RESULT

*Proof of Theorem 1.5.* Let  $\{\mathcal{G}_m^{(1)}\}_{m=1}^\infty$  be an increasing sequence of finite subsets of  $\mathcal{A}$  such that  $\mathcal{A} = \overline{\bigcup_{m=1}^\infty \mathcal{G}_m^{(1)}}$ . Let  $f$  be the function on the unit interval  $[0, 1]$  given by  $f(t) = 0$  for  $t < 1/2$  and  $f(t) = 1$  for  $t \geq 1/2$ . For each  $m$ , let  $\mathcal{G}_m^{(2)}$  be the (finite) set of elements of  $\mathcal{A}$  given by  $\mathcal{G}_m^{(2)} =_{df} \{f|a|/\|a\| : a \neq 0, a \in \mathcal{G}_m^{(1)}, f \text{ is continuous on the spectrum of } |a|/\|a\|\}$  (here, given  $a \in \mathcal{A}$ ,  $|a|$  is the absolute value of  $a$  and  $\|a\|$  is the norm of  $a$ ). Note that  $\bigcup_{m=1}^\infty \mathcal{G}_m^{(2)}$  is dense in the set of projections of  $\mathcal{A}$ . Now for each  $m$ , let  $\mathcal{G}_m =_{df} \mathcal{G}_m^{(1)} \cup \mathcal{G}_m^{(2)}$ . Since  $\mathcal{A}$  is in  $\mathcal{LCH}^+$ , let  $\{\mathcal{A}_m\}_{m=1}^\infty$  be a sequence of unital separable subalgebras of  $\mathcal{A}$ , with Hausdorff spectrum and bounded irreducible representations, such that for each  $m$ ,  $a$  is within a distance  $1/2m$  of an element, say  $\phi_m(a)$ , of  $\mathcal{A}_m$  for every  $a \in \mathcal{G}_m$ . If  $a$  is a projection in  $\mathcal{G}_m^{(2)}$ , we further require that  $\phi_m(a)$  be a projection. Now, for each  $m$ ,  $\mathcal{A}_m$  need not be a continuous trace  $C^*$ -algebra, but by [19, Theorem 4],  $\mathcal{A}_m$  is “continuous trace” with respect to the normalized trace; that is, for each  $a \in \mathcal{A}_m$ , the map  $\widehat{\mathcal{A}_m} \rightarrow \mathbb{R}$  given by  $\pi \mapsto tr(\pi(a))$  is continuous (where  $tr$  is the unital, normalized trace on the image of  $\pi$ , and  $\widehat{\mathcal{A}_m}$  is the spectrum space of irreducible representations of  $\mathcal{A}_m$ ). But for each  $m$ , for each  $p \in \mathcal{G}_m^{(2)}$ , the map  $\pi \mapsto tr(\pi(\phi_m(p)))$  can take on only finitely many (rational) values. Hence,  $\widehat{\mathcal{A}_m}$  is the disjoint union of finitely many clopen sets such that for each  $p \in \mathcal{G}_m^{(2)}$ , the map  $\pi \mapsto tr(\pi(\phi_m(p)))$  has constant value on each clopen set. Hence, for each  $m$ ,  $\mathcal{A}_m$  can be realized as a finite direct sum  $\mathcal{A}_m = \bigoplus_{i=1}^{N_m} \mathcal{A}_{m,i}$  where each summand  $\mathcal{A}_{m,i}$  has spectrum being one of the clopen sets. In particular, this means that for every  $m$ , for every projection  $p \in \mathcal{G}_m^{(2)}$ , for  $1 \leq i \leq N_m$ , the map  $\pi \mapsto tr(\pi(1_{\mathcal{A}_{m,i}} \phi_m(p) 1_{\mathcal{A}_{m,i}}))$  is constant on the spectrum  $\widehat{\mathcal{A}_{m,i}}$  (where “ $tr$ ”, as always, denotes the unital normalized trace on the image of  $\pi$ ). We may assume that  $1_{\mathcal{A}_m} = 1_{\mathcal{A}}$  for every  $m$ . Let  $\mathcal{B} =_{df} \sum_{m=1}^\infty \bigoplus_{i=1}^{N_m} \mathcal{A}_{m,i} = \sum_{l=1}^\infty \mathcal{B}_l$ . Then the multiplier algebra of  $\mathcal{B}$  is  $\mathcal{M}(\mathcal{B}) = \prod_{l=1}^\infty \mathcal{B}_l$  such that each  $\mathcal{B}_l$  is one of the  $\mathcal{A}_{m,i}$ s.

For  $a \in \bigcup_{m=1}^\infty \mathcal{G}_m$  and for each strictly positive integer  $l$ , let  $(a, l)$  be an element  $b \in \mathcal{B}_l$  defined in the following manner: Suppose that  $\mathcal{B}_l$  is the summand  $\mathcal{A}_{m,i}$  of  $\mathcal{M}(\mathcal{B})$ . If  $a$  is in  $\mathcal{G}_m$ , then let  $(a, l)$  be  $1_{\mathcal{B}_l} \phi_m(a) 1_{\mathcal{B}_l}$ . Otherwise, let  $(a, l)$  be zero.

We may assume that for every integer  $l$ ,  $(1_{\mathcal{A}}, l) = 1_{\mathcal{B}_l}$ . We have an  $*$ -homomorphism  $\Gamma: \mathcal{A} \rightarrow \mathcal{M}(\mathcal{B})/\mathcal{B}$  which is defined as follows: suppose that  $a \in \mathcal{A}$ . Let  $\{a_n\}_{n=1}^\infty$  be a sequence in  $\bigcup_{m=1}^\infty \mathcal{G}_m$  which converges to  $a$ . Then we let  $\Gamma(a) =_{df} \lim_{n \rightarrow \infty} (a_n, l)/\mathcal{B}$ . One can check that  $\Gamma$  is indeed a well-defined  $*$ -homomorphism.

Now since  $\mathcal{A}$  is simple,  $\Gamma$  is either injective or the zero map. Since  $(1_{\mathcal{A}}, l) = 1_{\mathcal{B}_l}$  for every  $l$ ,  $\Gamma$  is unital and hence must be injective.

For each  $l$ , let  $\tau_l$  be a unital trace on  $\mathcal{B}_l$  obtained by a point evaluation on  $\hat{\mathcal{B}}_l$ , the spectrum of  $\mathcal{B}_l$  (that is,  $\tau_l$  is obtained by composing an irreducible representation of  $\mathcal{B}_l$  with the usual unital trace on matrices).

Let  $\epsilon > 0$  and a finite subset  $\mathcal{F}$  of  $\mathcal{A}$  be given. To show that  $\mathcal{A}$  is TAF, we need to prove that there is a projection  $p \in \mathcal{A}$  and there is a finite dimensional  $C^*$ -subalgebra  $\mathcal{C}$  of  $\mathcal{A}$ , with  $1_{\mathcal{C}} = 1 - p$ , such that:

- (1)  $\sup_{r \in T(\mathcal{A})} \tau(p) < \epsilon$ ,
- (2)  $\|pf - fp\| < \epsilon$  for every  $f \in \mathcal{F}$ , and
- (3)  $(1 - p)f(1 - p)$  is within  $\epsilon$  of an element of  $\mathcal{C}$  for every  $f \in \mathcal{F}$ .

So, let  $\epsilon$  and  $\mathcal{F}$  be given as above. To simplify notation, we may assume that each element of  $\mathcal{F}$  has norm less than or equal to one (adjust  $\epsilon$  if necessary).

**Claim.** There is a strictly positive integer  $L$  such that for each  $l \geq L$ , there is a projection  $p_l$  in  $\mathcal{B}_l$  and a finite dimensional  $C^*$ -subalgebra  $\mathcal{C}_l$  of  $\mathcal{B}_l$  with  $1_{\mathcal{C}_l} = 1_{\mathcal{B}_l} - p_l$  such that:

- (1) if  $\mathcal{B}_l$  is the summand  $\mathcal{A}_{m,i}$ , then  $p_l$  has the form  $1_{\mathcal{B}_l} \phi_m(p) 1_{\mathcal{B}_l}$  for some projection  $p \in \mathcal{G}_m^{(2)}$ ,
- (2)  $\tau_l(p_l) < \epsilon/100$ ,
- (3)  $\|p_l f - f p_l\| < \epsilon/100$  for every  $f \in \mathcal{F}$ , and
- (4)  $1_{\mathcal{C}_l} f 1_{\mathcal{C}_l}$  is within  $\epsilon/100$  of an element of  $\mathcal{C}_l$  for every  $f \in \mathcal{F}$ .

Now suppose, to the contrary, that the claim is not true. Let  $\{l_\alpha\}_{\alpha \in I}$  be a subnet of the sequence of positive integers such that for each  $\alpha \in I$ , the statement of the claim does not hold for  $l = l_\alpha$ . Now for each integer  $k$ , let  $\tilde{\tau}_k$  be the trace on  $\mathcal{M}(\mathcal{B}) = \prod_{l=1}^\infty \mathcal{B}_l$  given by  $\tilde{\tau}_k((a_l)_{l=1}^\infty) = \tau_k(a_k)$  ( $\tau_k$  is defined two paragraphs before the claim, and  $a_l \in \mathcal{B}_l$  for every  $l$ ). Now since  $T(\mathcal{M}(\mathcal{B}))$  is  $w^*$ -compact, the net  $\{\tilde{\tau}_{l_\alpha}\}_{\alpha \in I}$  has a converging subnet. For simplicity, let us assume that  $\{\tilde{\tau}_{l_\alpha}\}_{\alpha \in I}$  converges to, say  $\tilde{\tau}$ . Note that  $\tilde{\tau}$  induces a trace on  $\mathcal{M}(\mathcal{B})/\mathcal{B}$ , which we also denote by “ $\tilde{\tau}$ ”. Since  $\Gamma: \mathcal{A} \rightarrow \mathcal{M}(\mathcal{B})/\mathcal{B}$  is a unital  $*$ -embedding,  $\tilde{\tau} \circ \Gamma$  is a tracial state on  $\mathcal{A}$ . For simplicity, we will also denote  $\tilde{\tau} \circ \Gamma$  by “ $\tilde{\tau}$ ”.

Note that the argument of Theorem 1.3 actually works for any (arbitrary) *single* trace (see either [1, Corollary 7.11], [15] or [17, Theorem 5.16], and in the locally AH case, an elementary proof can be obtained using the argument in the last section of [11]). Hence, we have that there exists a projection  $q \in \mathcal{A}$  and a finite dimensional  $C^*$ -subalgebra  $\mathcal{D}$  of  $\mathcal{A}$  with  $1_{\mathcal{D}} = 1 - q$  such that:

- (1)  $\tilde{\tau}(q) < \epsilon/1000$ ,
- (2)  $\|qf - fq\| < \epsilon/1000$  for every  $f \in \mathcal{F}$ , and
- (3)  $(1 - q)f(1 - q)$  is within  $\epsilon/1000$  of an element of  $\mathcal{D}$  for every  $f \in \mathcal{F}$ .

Now by our choices of the  $\mathcal{A}_m$ s and  $\mathcal{G}_m^{(2)}$ s, there is a positive integer  $M > 0$ , and there is a sequence  $\{\epsilon_m\}_{m=1}^\infty$  of positive real numbers converging to zero, such that for each  $m \geq M$ , we have the following:

- (a) There is a matrix algebra, say  $\mathcal{D}_m$ , which is a subalgebra of  $\mathcal{A}_m$ , and there is a unitary element  $U$  of  $\mathcal{A}$  such that (i)  $\mathcal{D}_m = U\mathcal{D}_m U^*$ , and (ii)  $U$  is within  $\epsilon_m$  of  $1_{\mathcal{A}}$ .
- (b)  $UqU^* = 1_{\mathcal{A}} - 1_{\mathcal{D}_m}$ , and  $1_{\mathcal{A}} - 1_{\mathcal{D}_m}$  is an element of  $\mathcal{G}_m^{(2)}$ . (Recall that  $1_{\mathcal{A}} = 1_{\mathcal{A}_m}$ .)
- (c)  $(1_{\mathcal{A}} - 1_{\mathcal{D}_m})a$  is within  $\epsilon/500$  of  $a(1_{\mathcal{A}} - 1_{\mathcal{D}_m})$ , for every  $a \in \mathcal{F}$ .

Now suppose that for each  $m$ ,  $\mathcal{A}_m = \bigoplus_{l=L_m}^{L_{m+1}-1} \mathcal{B}_l$  (so  $\mathcal{B}_{L_m+k} = \mathcal{A}_{m,k+1}$  and  $L_{m+1} - L_m = N_m$ ). And suppose that for  $m \geq M$ ,  $1_{\mathcal{A}} - 1_{\mathcal{D}_m} = q_{L_m} \oplus q_{L_m+1} \oplus q_{L_m+2} \oplus \dots \oplus q_{L_{m+1}-1}$ , where  $q_{L_m+k}$  is a projection in  $\mathcal{B}_{L_m+k}$ , for each  $k$ . Then  $\Gamma(q) = (0, 0, \dots, 0, q_{L_M}, q_{L_M+1}, q_{L_M+2}, \dots) / \mathcal{B}$ , where  $q_{L_M}$  is in the  $L_M$ th position, and where we view  $\mathcal{B}$  as  $\mathcal{B} = \sum_{l=1}^{\infty} \mathcal{B}_l$ .

By the definition of  $\tilde{\tau}$ , and since  $\tilde{\tau}(q) < \epsilon/1000$ , we must have that  $\lim_{\alpha} \tau_{l_{\alpha}}(q_{l_{\alpha}}) < \epsilon/1000$ . Choose  $\alpha_0$  such that for  $\alpha \geq \alpha_0$ ,  $\tau_{l_{\alpha}}(q_{l_{\alpha}}) < \epsilon/1000$ . Let  $m_0$  be the integer such that  $\mathcal{B}_{l_{\alpha_0}}$  comes from  $\mathcal{A}_{m_0}$ . Choosing  $\alpha_0$  “large” enough if necessary, we may assume that  $m_0 \geq M$  and  $\epsilon_{m_0} < \epsilon/1000$ . Hence, taking  $l = l_{\alpha_0}$ ,  $m = m_0$ ,  $\mathcal{C}_l = 1_{\mathcal{B}_{l_{\alpha_0}}} \mathcal{D}_{m_0} 1_{\mathcal{B}_{l_{\alpha_0}}}$ , and  $p_l = q_{l_{\alpha_0}}$ , we have that clauses (1)–(4) of the Claim are satisfied for  $l = l_{\alpha_0}$ . This is a contradiction. Hence, the Claim must be true.

So let  $L$  and  $\mathcal{C}_l$  and  $p_l \forall l \geq L$  be as in the Claim. Recalling our definition of  $\mathcal{B}$  and the  $\mathcal{B}_l$ s, suppose that  $m, N_m$  are integers such that  $\mathcal{B}_{L+j} = \mathcal{A}_{m,j}$  for  $1 \leq j \leq N_m$  and  $\mathcal{A}_m = \bigoplus_{j=1}^{N_m} \mathcal{A}_{m,j} = \bigoplus_{j=1}^{N_m} \mathcal{B}_{L+j}$ . Let  $r =_{df} \bigoplus_{j=1}^{N_m} p_{L+j}$  and  $\mathcal{E} =_{df} \bigoplus_{j=1}^{N_m} \mathcal{C}_{L+j}$ . Then (a)  $1_{\mathcal{E}} = 1 - r$ , (b)  $\|rf - fr\| < \epsilon$  for every  $f \in \mathcal{F}$ , and (c)  $(1 - r)f(1 - r)$  is within  $\epsilon$  of an element of (the finite-dimensional  $C^*$ -algebra)  $\mathcal{E}$  for every  $f \in \mathcal{F}$ .

Also, by clause (1) of the Claim, it follows that for  $1 \leq j \leq N_m$ , the map on  $\widehat{\mathcal{B}_{L+j}}$  (the spectrum of  $\mathcal{B}_{L+j}$ ), given by  $\pi \mapsto tr(\pi(p_{L+j}))$ , is a constant rational-valued function (here,  $tr$  is the unital trace on matrices). Hence, since  $\tau_{L+j}(p_{L+j}) < \epsilon/100$  for  $1 \leq j \leq N_m$ , it follows that  $\tau(r) < \epsilon$  for every  $\tau \in T(\mathcal{A})$ .

So since  $\epsilon$  and  $\mathcal{F}$  are arbitrary,  $\mathcal{A}$  is TAF. □

REFERENCES

- [1] N. Brown, *Invariant means and finite representation theory of  $C^*$ -algebras*, preprint. Available online at <http://arxiv.org/abs/math.OA/0304009>.
- [2] M. Dadarlat, *Some remarks on the universal coefficient theorem in  $KK$ -theory*, *Constanța*, (2003), 65–74. MR2018224 (2004j:19005)
- [3] M. Dadarlat and S. Eilers, *Approximate homogeneity is not a local property*, *J. Reine Angew. Math.*, **507**, (1999), 1–13. MR1670254 (2000b:46094)
- [4] M. Dadarlat and S. Eilers, *On the classification of nuclear  $C^*$ -algebras*, *Proc. London Math. Soc.* (3), **85**, (2002), no. 1, 168–210. MR1901373 (2003d:19006)
- [5] M. Dadarlat and G. Gong, *A classification result for approximately homogeneous  $C^*$ -algebras of real rank zero*, *Geom. Funct. Anal.*, **7**, (1997), no. 4, 646–711. MR1465599 (98j:46062)
- [6] J. Dixmier,  *$C^*$ -algebras*, Paris, 1964; North Holland, Amsterdam/New York/Oxford.
- [7] G. Elliott and G. Gong, *On the classification of  $C^*$ -algebras of real rank zero II*, *Ann. of Math.* (2), **144**, (1996), no. 3, 497–610. MR1426886 (98j:46055)
- [8] G. Elliott, G. Gong, L. Li, *On the classification of simple inductive limit  $C^*$ -algebras, II: the isomorphism theorem*, preprint.
- [9] J. M. G. Fell, *The structure of algebras of operator fields*, *Acta Math.*, **106**, (1961), 233–280. MR0164248 (29:1547)
- [10] E. Kirchberg and W. Winter, *Covering dimension and quasidiagonality*, *Internat. J. Math.*, **15**, (2004), 63–85. MR2039212 (2005a:46148)
- [11] H. Lin, *Homomorphisms from  $C^*$ -algebras of continuous trace*, *Math. Scand.*, **86**, (2000), no. 2, 249–272. MR1754997 (2001b:46088)
- [12] H. Lin, *The tracial topological rank of  $C^*$ -algebras*, *Proc. London Math. Soc.* (3), **83**, (2001), no. 1, 199–234. MR1829565 (2002e:46063)
- [13] H. Lin, *Classification of simple  $C^*$ -algebras of tracial topological rank zero*, *Duke Math. J.*, **125**, (2004), 91–119. MR2097358
- [14] H. Lin, *Classification of simple  $C^*$ -algebras with unique traces*, *Amer. J. Math.*, **120**, (1998), no. 6, 1289–1315. MR1657178 (2000c:46106)
- [15] H. Lin, *Locally type I simple TAF  $C^*$ -algebras*, preprint.
- [16] H. Lin, *Simple AH-algebras of real rank zero*, preprint. To appear in the Proceedings of the AMS.

- [17] H. Lin, *Traces and simple  $C^*$ -algebras with tracial topological rank zero*. J. Reine Angew. Math., **568**, (2004), 99–137. MR2034925 (2005b:46122)
- [18] I. Raeburn and D. Williams, *Morita equivalence and continuous trace  $C^*$ -algebras*, Mathematical Surveys and Monographs, **60**, AMS, Providence, RI, 1998. MR1634408 (2000c:46108)
- [19] N. B. Vasil'ev,  *$C^*$ -algebras with finite-dimensional irreducible representations*, Uspehi Mat. Nauk, **21**, (1966), no. 1, (127), 135–154. MR0201994 (34:1871)

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF NEW BRUNSWICK, FREDERICTON, NEW BRUNSWICK, CANADA E3B 5A3

*E-mail address:* `pwn@erdos.math.unb.ca`

*Current address:* The Fields Institute for Research in Mathematical Sciences, 222 College Street, Toronto, Ontario, Canada M5T 3J1