

ON BIACCESSIBLE POINTS OF THE MANDELBROT SET

SAEED ZAKERI

(Communicated by Linda Keen)

ABSTRACT. This paper provides a description for the quadratic polynomials on the boundary of the Mandelbrot set \mathcal{M} which are typical in the sense of harmonic measure. In particular, it is shown that a typical point on the boundary of \mathcal{M} has a unique parameter ray landing on it. Applications of this result in the study of embedded arcs in \mathcal{M} and the lamination associated with \mathcal{M} are given.

1. INTRODUCTION

Consider the complex quadratic family $\{f_c : z \mapsto z^2 + c\}_{c \in \mathbb{C}}$ and its connectedness locus \mathcal{M} , commonly known as the “Mandelbrot set.” This paper will give a short proof of the following

Theorem 1. *There exists a Borel set $X \subset \partial\mathcal{M}$ of full harmonic measure such that for every $c \in X$:*

- (i) *the Julia set $J_c = J(f_c)$ is locally connected and full;*
- (ii) *the post-critical orbit $\{c, f_c(c), f_c^{\circ 2}(c), \dots\}$ is evenly distributed with respect to the harmonic measure on J_c (in particular it is dense in J_c);*
- (iii) *\mathcal{M} is locally connected at c , only one parameter ray lands at c , the impression of this ray is trivial, and no other parameter ray contains c in its impression. Moreover, the dynamic ray of the same angle lands at the critical value $c \in J_c$.*

The statement can be recovered from the work of Bruin and Schleicher [BSc] in which they use symbolic dynamics to study various combinatorial and topological aspects of the quadratic family. Modulo the uniqueness of rays in (iii), it also follows from the work of Smirnov [S1] and Graczyk and Świątek [GS]. Instead of using their results, here we present a very elementary argument based on Yoccoz’s local connectivity theorem and the topology of trees in Julia sets.

Recall that $c \in \partial\mathcal{M}$ is *biaccessible* if two or more parameter rays land at c . Thus, we have the following corollary of Theorem 1:

Corollary 2. *The set of all biaccessible points in the Mandelbrot set has harmonic measure zero.*

Received by the editors February 5, 2004 and, in revised form, January 24, 2005.
2000 *Mathematics Subject Classification.* Primary 37F10, 37F20, 37F35, 35F45.

©2006 American Mathematical Society
Reverts to public domain 28 years from publication

The analogue of this result has been known for all connected quadratic Julia sets, with the exception of the Julia set of the Chebyshev map $f_{-2} : z \mapsto z^2 - 2$, which is a straight line segment (see [S2], [Z1], [Zd]).

As another corollary, in §4 we show that

Theorem 3. *Every embedded arc in the Mandelbrot set has harmonic measure zero.*

In particular, the *real slice* $\mathcal{M} \cap \mathbb{R} = [-2, 1/4]$ of the Mandelbrot set has harmonic measure zero. In [Z2], this result came as a byproduct of a completely different method.

Let $\Lambda_{\mathcal{M}} \subset \mathbb{D}$ be the *lamination associated with* \mathcal{M} (see §5 for the definition and compare Figure 3).

Theorem 4. *The area of $\Lambda_{\mathcal{M}}$ is zero, and its Hausdorff dimension is two.*

The proof, presented in §5, will depend on Theorem 1 and the fact that the set of angles of the parameter rays which land on the real slice of \mathcal{M} has Hausdorff dimension one [Z2].

Let us point out that the area statement in Theorem 4 bears resemblance to Thurston's observation that geodesic laminations on hyperbolic surfaces of finite area have measure zero (see for example [Th, Chapter 8]). On the other hand, the dimension statement in Theorem 4 is in sharp contrast with the theorem of Birman and Series according to which such geodesic laminations always have Hausdorff dimension one [BS]. It may be worthwhile to further study the *metric* structure of laminations that arise in complex dynamics (such as $\Lambda_{\mathcal{M}}$ and those associated with Julia sets) and develop a theory parallel to geodesic laminations on hyperbolic surfaces.

2. PRELIMINARIES

First we briefly recall some facts about the conformal geometry of the complement of a planar continuum; see [A], [Mc], [M1] or [P] for details. Let $K \subset \mathbb{C}$ be a non-degenerate continuum. Assume further that K is *full* in the sense that its complement $\mathbb{C} \setminus K$ is connected. By the Riemann mapping theorem, there exists a unique conformal isomorphism $\psi_K : \mathbb{C} \setminus \overline{\mathbb{D}} \xrightarrow{\cong} \mathbb{C} \setminus K$ which satisfies $\lim_{z \rightarrow \infty} \psi_K(z)/z > 0$. By the *external ray at angle* $t \in \mathbb{T} = \mathbb{R}/\mathbb{Z}$ is meant the analytic curve $R_K(t) = \psi_K(\{re^{2\pi it} : r > 1\})$. We say that $R_K(t)$ *lands* at $p \in \partial K$ if $\lim_{r \rightarrow 1} \psi_K(re^{2\pi it}) = p$. According to Carathéodory, ψ_K extends continuously to the boundary circle $\partial\mathbb{D}$ if and only if K is locally connected, in which case all external rays land [M1, Theorem 17.14]. Whether or not K is locally connected, a theorem of Beurling asserts that $R_K(t)$ lands for all t outside a set of logarithmic capacity zero [P, Theorem 9.19].

The (*prime-end*) *impression* $I_K(t)$ of the ray $R_K(t)$ is the set of all $z \in \partial K$ for which there is a sequence $\{w_n\}$ in $\mathbb{C} \setminus \overline{\mathbb{D}}$ such that $w_n \rightarrow e^{2\pi it}$ and $\psi_K(w_n) \rightarrow z$. It is not hard to show that $I_K(t)$ is a non-empty sub-continuum of ∂K , and $\bigcup_{t \in \mathbb{T}} I_K(t) = \partial K$. Here is another easy consequence of the definition of impression that we will need in the proof of Theorem 4 in §5:

$$(2.1) \quad \limsup_{n \rightarrow \infty} I_K(t_n) \subset I_K(t) \quad \text{whenever} \quad t_n \rightarrow t.$$

When $I_K(t)$ reduces to a point, we say that it is a *trivial* impression. This happens precisely when ψ_K has a continuous extension to $e^{2\pi it}$. A ray with trivial impression necessarily lands, but a landing ray may well have a non-trivial impression.

A point $p \in \partial K$ is *accessible* if it is the landing point of an external ray, and is *biaccessible* if at least two external rays land on it. It can be shown that p is biaccessible if and only if $K \setminus \{p\}$ is disconnected. Given a subset $E \subset K$, define

$$(2.2) \quad \psi_K^{-1}(E) = \{t \in \mathbb{T} : R_K(t) \text{ lands at a point of } E\}.$$

The continuum K decomposes into the disjoint union $K = K^0 \cup K^1 \cup K^2 \cup \dots \cup K^\infty$, where

$$K^n = \{p \in K : \psi_K^{-1}(p) \text{ has exactly } n \text{ elements}\}, \quad 0 \leq n \leq \infty.$$

It is easy to verify that each K^n is a Borel measurable subset of K , and K^0 contains the interior of K (possibly empty). Note that in terms of this decomposition, a point is accessible if and only if it belongs to $K \setminus K^0$, and is biaccessible if and only if it belongs to $K \setminus (K^0 \cup K^1)$.

The normalized 1-dimensional Lebesgue measure on $\partial\mathbb{D} \cong \mathbb{T}$ pushes forward by the Riemann map ψ_K to a Borel measure supported on ∂K called the *harmonic measure*. In other words, the harmonic measure of a Borel subset $E \subset K$ is given by the length of the set $\psi_K^{-1}(E)$ defined in (2.2). By the theorem of Beurling mentioned above, the set K^0 has harmonic measure zero. The same is true for $K^3 \cup \dots \cup K^\infty$, since this union is easily seen to be a countable set [P, Corollary 2.19] and single points in K have harmonic measure zero by a theorem of F. and M. Riesz [M1, Theorem A.3]. It follows that $K = K^1 \cup K^2$ up to a set of harmonic measure zero.

We now give a brief account of a few basic results in the dynamics of the complex quadratic family; further details can be found in [DH1], [DH2], [Mc], [M2] and [M3]. Let $c \in \mathbb{C}$. The *filled Julia set* of $f_c : z \mapsto z^2 + c$, denoted by K_c , is the set of all points in the plane with bounded forward orbit under f_c . The topological boundary $J_c = \partial K_c$ is called the *Julia set* of f_c . Both sets are non-empty, compact, and totally invariant, with K_c being full. An external ray for K_c is called a *dynamic ray*. By a *cycle* of f_c is meant a periodic orbit $z \mapsto f_c(z) \mapsto \dots \mapsto f_c^p(z) = z$. The quantity $\lambda = (f_c^{\circ p})'(z)$ is called the *multiplier* of this cycle. The cycle is *attracting*, *repelling*, or *indifferent* according to whether $|\lambda| < 1$, $|\lambda| > 1$, or $|\lambda| = 1$. A parameter c is *Misiurewicz* if the forward orbit under f_c of the critical point 0 is finite but not periodic.

The Mandelbrot set \mathcal{M} consists of all parameters $c \in \mathbb{C}$ for which the (filled) Julia set of f_c is connected. Equivalently, $c \in \mathcal{M}$ if and only if $0 \in K_c$. It can be shown that \mathcal{M} is a full continuum. We will use the term *parameter ray* for an external ray of \mathcal{M} . A parameter $c \in \mathcal{M}$ is called *hyperbolic* if the corresponding quadratic f_c has a necessarily unique attracting cycle in the plane. Hyperbolic parameters appear as connected components of the interior of \mathcal{M} . The main hyperbolic component H_0 consists of all $c \in \mathcal{M}$ for which f_c has an attracting fixed point. It is the prominently visible cardioid in any picture of \mathcal{M} and is defined by the inequality $|1 - \sqrt{1 - 4c}| < 1$ (compare Figure 3 left). The *period* of a hyperbolic component H is the length of the unique attracting cycle of f_c for any $c \in H$. There is a canonical conformal isomorphism $\lambda_H : H \xrightarrow{\cong} \mathbb{D}$ which assigns to each $c \in H$ the multiplier of its attracting cycle. The map λ_H extends to a homeomorphism $\overline{H} \xrightarrow{\cong} \overline{\mathbb{D}}$. The *root* of H is by definition the point $\lambda_H^{-1}(1) \in \partial H$. We call H a *satellite* component if its root is on the boundary of another hyperbolic component. There are exactly

two parameter rays of angles $\theta_- < \theta_+$ landing at the root of H which are rational numbers of the form $n/(2^p - 1)$, where p is the period of H . (When $H = H_0$, the two angles $\theta_- = 0$ and $\theta_+ = 1$ coincide, and only one ray lands at the root $c = 1/4$.) It follows that these two angles have binary expansions of the form

$$(2.3) \quad \theta_- = 0.\overline{\theta_0} \quad \text{and} \quad \theta_+ = 0.\overline{\theta_1},$$

where θ_0 and θ_1 are binary words of length p , and the bars indicate infinite repetition. Conversely, if t is a rational of odd denominator, the parameter ray $R_{\mathcal{M}}(t)$ lands at the root of a unique hyperbolic component.

For every hyperbolic component H , denote by $\iota_H : \mathcal{M} \hookrightarrow \mathcal{M}$ the Douady-Hubbard *tuning* map (see [DH1] or [M3]). Then ι_H is a topological embedding which maps H_0 onto H , respects roots of hyperbolic components, and maps $\partial\mathcal{M}$ into $\partial\mathcal{M}$. The image $M_H = \iota_H(\mathcal{M})$ is a subset of \mathcal{M} homeomorphic to \mathcal{M} , with $\partial H \subset \partial M_H \subset \partial\mathcal{M}$. We have $M_{H_0} = \mathcal{M}$, but when $H \neq H_0$, M_H is a proper subset of \mathcal{M} often described as the *small copy* of \mathcal{M} growing from H .

The tuning operation also acts on parameter rays. Let $\theta_- = 0.\overline{\theta_0} < \theta_+ = 0.\overline{\theta_1}$ be the angles of the two parameter rays landing at the root of a period- p hyperbolic component $H \neq H_0$ as in (2.3). For any angle $t \in \mathbb{T}$ with the binary expansion $0.t_1t_2t_3\cdots$, define the *tuned angle*

$$A_H(t) = 0.\theta_{t_1}\theta_{t_2}\theta_{t_3}\cdots$$

by concatenating blocks of words of length $p > 1$. Note that under this tuning algorithm on angles, a dyadic rational has two distinct images since it has two different binary representations. The image $A_H(\mathbb{T})$ is a self-similar Cantor set and as such has length zero. In fact, let $\Lambda_i : \mathbb{T} \rightarrow \mathbb{T}$ ($i = 0, 1$) be the map defined by $\Lambda_i(0.t_1t_2t_3\cdots) = 0.\theta_{t_1}\theta_{t_2}\theta_{t_3}\cdots$. Then Λ_i is an affine contraction by a factor 2^{-p} and the image $A_H(\mathbb{T})$ is precisely the invariant set generated by the pair Λ_0, Λ_1 [Ma]. It can be shown that if $R_{\mathcal{M}}(t)$ lands at $c \in M$, then the parameter ray $R_{\mathcal{M}}(A_H(t))$ lands at the tuned image $\iota_H(c) \in M_H$ [D1]. Furthermore, with countably many possible exceptions, these are the only parameter rays that land on the small copy M_H . As a result,

$$(2.4) \quad \text{length}(\psi_{\mathcal{M}}^{-1}(M_H)) = \text{length}(A_H(\mathbb{T})) = 0 \quad \text{if } H \neq H_0.$$

Finally, a quadratic f_c with $c \in \mathcal{M}$ is *simply renormalizable* of period $p > 1$ if there are topological disks U and V containing 0 such that (i) U is compactly contained in V ; (ii) $f^{\circ p} : U \rightarrow V$ is a degree 2 branched covering; (iii) $0 \in E$, where E is the set of points in U whose forward orbits under $f^{\circ p}$ are contained in U ; and (iv) the images $E, f(E), \dots, f^{\circ p-1}(E)$ do not disconnect one another. Simply renormalizable quadratics correspond to parameters in small copies of \mathcal{M} . More precisely, if f_c is simply renormalizable of period $p > 1$, then $c \in M_H$ for some period- p hyperbolic component H . Conversely, if H is a period- p hyperbolic component with $p > 1$, every $c \in M_H$ (with the exception of the root of H in the satellite case) gives a quadratic f_c which is simply renormalizable of period p [DH1].

3. PROOF OF THEOREM 1

The proof of Theorem 1 begins as follows. The set $\mathcal{M}_{\text{ren}} \subset \mathcal{M}$ of parameters c for which the quadratic f_c is simply renormalizable is contained in the countable

union $\bigcup_{H \neq H_0} M_H$ of all small copies. Thus,

$$\psi_{\mathcal{M}}^{-1}(\mathcal{M}_{\text{ren}}) \subset \bigcup_{H \neq H_0} \psi_{\mathcal{M}}^{-1}(M_H).$$

Since by (2.4) each $\psi_{\mathcal{M}}^{-1}(M_H)$ has length zero, it follows that \mathcal{M}_{ren} has harmonic measure zero.

For each irreducible fraction $0 < p/q < 1$ there is a period- q hyperbolic component $H_{p/q}$ whose root is $\lambda_{H_0}^{-1}(e^{2\pi ip/q}) \in \partial H_0$ (see Figure 3 left for the case $p/q = 1/3$). The two parameter rays landing at this root have angles of the form $\theta_- = n/(2^q - 1)$ and $\theta_+ = (n + 1)/(2^q - 1)$. For t in the open interval $I_{p/q} = (\theta_-, \theta_+) \subset \mathbb{T}$, the parameter ray $R_{\mathcal{M}}(t)$ does not land on ∂H_0 . It follows that $\psi_{\mathcal{M}}^{-1}(\partial H_0) \subset \mathbb{T} \setminus \bigcup_{0 < p/q < 1} I_{p/q}$. Since the intervals $I_{p/q}$ are disjoint, we obtain

$$\text{length}(\psi_{\mathcal{M}}^{-1}(\partial H_0)) \leq 1 - \sum_{0 < p/q < 1} \frac{1}{2^q - 1}.$$

An easy computation gives $\sum_{0 < p/q < 1} 1/(2^q - 1) = 1$ (compare [D1] or [GM]) and we conclude that ∂H_0 has harmonic measure zero.

Let $\mathcal{M}_{\text{ind}} \subset \mathcal{M}$ be the set of parameters c for which f_c has an indifferent cycle. These are precisely the parameters which lie on the boundary of hyperbolic components of \mathcal{M} , so

$$\mathcal{M}_{\text{ind}} = \partial H_0 \cup \bigcup_{H \neq H_0} \partial H \subset \partial H_0 \cup \bigcup_{H \neq H_0} M_H.$$

Since each set on the right has harmonic measure zero, it follows that the harmonic measure of \mathcal{M}_{ind} is zero.

Now consider the Borel set $Y = \partial \mathcal{M} \setminus (\mathcal{M}_{\text{ren}} \cup \mathcal{M}_{\text{ind}})$ consisting of all c for which the quadratic f_c is non-renormalizable and without indifferent cycles. By the above discussion, Y has full harmonic measure. According to Yoccoz, for every $c \in Y$ the Julia set J_c is a locally connected full continuum. Moreover, \mathcal{M} is locally connected at c , only finitely many parameter rays land at c , and the dynamic rays of the same angles land at the critical value c of f_c (see [H], [M2], [R]). This statement can be improved by showing that the parameter rays landing at c have trivial impressions, and no other ray can contain c in its impression. In the case c is a Misiurewicz parameter, this is shown by Tan Lei, building upon the earlier work of Douady and Hubbard [T1]. In the case c is non-Misiurewicz, the sharper statement follows from Tan Lei's result and the proof of Yoccoz's theorem. In fact, Yoccoz constructs a nested sequence of *para-puzzles* $\{\Pi_n\}$ with the following properties:

- Π_n is a closed topological disk containing c ;
- $\Pi_n \cap \mathcal{M}$ is connected;
- $\partial \Pi_n \cap \mathcal{M}$ consists of finitely many Misiurewicz parameters; and
- $\bigcap \Pi_n = \{c\}$

(compare [R]). Let c belong to some impression $I_{\mathcal{M}}(t)$, and assume $\partial \Pi_n \cap I_{\mathcal{M}}(t) \neq \emptyset$ for some n . Since this intersection is contained in $\partial \Pi_n \cap \mathcal{M}$, it could only be a finite set $\{c_1, \dots, c_k\}$ of Misiurewicz parameters. Tan Lei's result would then show that $I_{\mathcal{M}}(t) = \{c_j\}$ for some j . This would imply $c = c_j$, which would contradict the assumption of c being non-Misiurewicz. Thus, $\partial \Pi_n \cap I_{\mathcal{M}}(t) = \emptyset$ and so $I_{\mathcal{M}}(t) \subset \Pi_n$ for all n . It follows that $I_{\mathcal{M}}(t) = \{c\}$ and in particular $R_{\mathcal{M}}(t)$ lands at c .

Let $c \in Y$ so that J_c is locally connected and full. The normalized Riemann map $\psi_{J_c} : \mathbb{C} \setminus \overline{\mathbb{D}} \rightarrow \mathbb{C} \setminus J_c$ induces a continuous map $\partial\mathbb{D} \cong \mathbb{T} \rightarrow J_c$ which conjugates the action of the doubling map $t \mapsto 2t$ to f_c :

$$\psi_{J_c}(2t) = f_c(\psi_{J_c}(t)).$$

If the parameter ray $R_{\mathcal{M}}(t)$ lands at c , the dynamic ray $R_{J_c}(t)$ lands at the critical value $c \in J_c$, and it follows by induction that $R_{J_c}(2^k t)$ lands at $f_c^{\circ k}(c)$ for all $k \geq 0$. Since the harmonic measure on J_c is the push-forward of Lebesgue measure on \mathbb{T} under ψ_{J_c} , it follows that the post-critical orbit $\{c, f_c(c), f_c^{\circ 2}(c), \dots\}$ is evenly distributed on J_c whenever the orbit $t \mapsto 2t \mapsto 4t \mapsto \dots$ is evenly distributed on \mathbb{T} . (Recall that a sequence $\{x_n\}$ in a topological space X is *evenly distributed* with respect to a probability measure μ if

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \phi(x_n) = \int_X \phi(x) d\mu(x)$$

for every bounded continuous function $\phi : X \rightarrow \mathbb{R}$.) Since the doubling map is ergodic with respect to Lebesgue measure on the circle, the Ergodic Theorem shows that there is a full length Borel set $B \subset \mathbb{T}$ consisting of angles t whose orbits $t \mapsto 2t \mapsto 4t \mapsto \dots$ are evenly distributed on \mathbb{T} . The intersection $\psi_{\mathcal{M}}^{-1}(Y) \cap B$ has full length, so the Borel set $X \subset Y$ which satisfies $\psi_{\mathcal{M}}^{-1}(X) = \psi_{\mathcal{M}}^{-1}(Y) \cap B$ has full harmonic measure. Moreover, the preceding remark shows that for each $c \in X$, the post-critical orbit $\{c, f_c(c), f_c^{\circ 2}(c), \dots\}$ is evenly distributed on J_c , and in particular is dense in J_c .

So far we have verified all the properties of X in Theorem 1 except the uniqueness of rays in (iii). This will be a corollary of the lemma below, which may be of independent interest.

Lemma 5. *Let J_c be locally connected and full. Suppose two or more dynamic rays land at c . Then there exists a forward invariant finite topological tree $T \subset J_c$ which contains the post-critical orbit $\{c, f_c(c), f_c^{\circ 2}(c), \dots\}$.*

It follows from this lemma that every $c \in X$ is the landing point of a unique parameter ray, for otherwise we could find a finite topological tree T in J_c containing the post-critical orbit and use the density of this orbit to conclude that the Julia set $J_c = T$ is a finite tree. The only quadratic with this property is the Chebyshev map $f_{-2} : z \mapsto z^2 - 2$ which is post-critically finite (see for example [M4, Lemma B.4] or [Z1]). This contradiction completes the proof of Theorem 1. \square

Proof of Lemma 5. Since J_c is locally connected and full, each pair z, w of distinct points in J_c can be joined by a unique embedded arc $[z, w] = [w, z]$ in J_c homeomorphic to the interval $[0, 1] \subset \mathbb{R}$. Let β be the fixed point of f_c at which the dynamic ray at angle 0 lands, and let α be the other fixed point. The following facts can be easily verified (see [Z1] for details and compare Figure 1):

- (3.1) If $z \in J_c$ is biaccessible, then $f_c^{\circ k}(z) \in [-\beta, \beta]$ for some $k \geq 0$;
- (3.2) $0 \in [-\beta, \beta]$ and $\alpha \in [-\beta, 0]$;
- (3.3) $f_c([\alpha, -\alpha]) = [c, \alpha]$ and $f_c([-\beta, \beta]) = [c, \alpha] \cup [\alpha, -\alpha] \cup [-\alpha, \beta] = [c, \beta]$;
- (3.4) if $z \in [-\alpha, \beta]$, then $f_c(z) \in [\alpha, z]$; and
- (3.5) for every $z \in J_c$, $f_c([z, \alpha]) \subset [f_c(z), \alpha] \cup [\alpha, c]$.




FIGURE 1. Some embedded arcs in the Julia set J_c .

Let $c_k = f_c^{o k}(c)$. The assumption that two rays land at c implies by (3.1) that $c_m \in [-\beta, \beta]$ for some $m \geq 0$. We claim that there exists $n \geq m$ such that $c_n \in [c, \alpha]$. Otherwise, by (3.3), $c_n \in [-\alpha, \beta]$ for all $n > m$. It follows from (3.4) that $[-\alpha, \beta] \supset [c_{n+1}, \beta] \supset [c_n, \beta]$ for all $n > m$. The limit of the monotone sequence $\{c_n\}_{n>m}$ will then be a fixed point of f_c in $[-\alpha, \beta]$. Since β is the only fixed point of f_c in $[-\alpha, \beta]$, this implies $c_n = \beta$ for all $n > m$. Since c has two rays landing on it and the forward orbit of c never hits the critical point 0, every c_n and hence β must be the landing point of two rays. This is a contradiction since β is the landing point of the unique ray at angle 0.

So choose $n \geq m$ such that $c_n \in [c, \alpha]$, or equivalently $[c_n, \alpha] \subset [c, \alpha]$. The union $T = \bigcup_{k=0}^{n-1} [c_k, \alpha]$ is then a forward invariant finite topological tree. In fact, by (3.5), $f_c([c_k, \alpha]) \subset [c_{k+1}, \alpha] \cup [\alpha, c] \subset T$ for every $0 \leq k \leq n - 1$. Since $c \in T$, the orbit of c is entirely contained in T . \square

4. EMBEDDED ARCS IN \mathcal{M}

In this section we prove Theorem 3 using Theorem 1 and the fact that the boundary of every interior component of \mathcal{M} has harmonic measure zero (Lemma 7 below). The argument is based on the following topological lemma.

Lemma 6. *Let $K \subset \mathbb{C}$ be a full non-degenerate continuum and let $\gamma \subset K$ be an embedded arc with endpoints z, w . Assume there are angles $t, s \in \mathbb{T}$ such that $I_K(t) = \{z\}$ and $I_K(s) = \{w\}$. Then every point of $\gamma \setminus \{z, w\}$ belongs either to the closure of an interior component of K , or to the impression of at least two external rays of K .*

Note that the assumption on the endpoints of γ is necessary. For example, the “comb”

$$K = [-1, 1] \cup [0, i] \cup \bigcup_{n=1}^{\infty} \left[\pm \frac{1}{n}, \pm \frac{1}{n} + i \right]$$

has no interior and the point $p = i/2$ on the embedded arc $\gamma = [0, i]$ belongs to the impression of the unique ray which lands at i (see Figure 2). Note also that the lemma is a generalization of the fact that every point of a “dendrite” (i.e., a locally connected full non-degenerate continuum without interior) is either an end or biaccessible (compare [Z1, Lemma 3]).

Proof. Assume $p \in \gamma \setminus \{z, w\}$ does not belong to the closure of any interior component of K . The union $R_K(t) \cup \gamma \cup R_K(s)$ separates the plane into two connected components C^+ and C^- . Take a sequence of open topological disks D_n around p such that $\text{diam } D_n \rightarrow 0$ and $D_n \setminus (R_K(t) \cup \gamma \cup R_K(s))$ has two connected components D_n^+ and D_n^- , labelled so that $D_n^+ \subset C^+$ and $D_n^- \subset C^-$ for all n . For each n , both topological disks D_n^\pm intersect $\mathbb{C} \setminus K$, for otherwise one of them would be




FIGURE 2.

contained in the interior of K and p would be in the closure of an interior component. Choose sequences $z_n^\pm \in D_n^\pm \setminus K$ and write $z_n^\pm = \psi_K(r_n \exp(2\pi i t_n^\pm))$. After passing to subsequences if necessary, we can assume that $t_n^\pm \rightarrow t^\pm$. It then follows from the definition of impression that $p \in I_K(t^+) \cap I_K(t^-)$.

Since $R_K(t_n^\pm) \subset C^\pm$, we have $R_K(t^\pm) \subset \overline{C^\pm}$. If $t^+ = t^-$, it follows that $R_K(t^+) = R_K(t^-) \subset \overline{C^+} \cap \overline{C^-}$, so $t^+ = t^- = t$ or $t^+ = t^- = s$. But then $p \in I_K(t^+) \cap I_K(t^-) \subset I_K(t) \cup I_K(s)$, which implies $p = z$ or $p = w$. This contradicts our assumption. Thus $t^+ \neq t^-$. \square

Lemma 7. *The union of the boundaries of the interior components of \mathcal{M} has harmonic measure zero.*

Proof. We show that this union is contained in the harmonic measure zero set $E = \partial H_0 \cup \bigcup_{H \neq H_0} M_H$ (see §3). Let W be an interior component of \mathcal{M} . If W is a hyperbolic component, then $\partial W \subset \partial H_0 \cup \bigcup_{H \neq H_0} \partial H \subset E$. Otherwise, W is a “queer” component. In this case, for every $c \in W$ the quadratic f_c has no indifferent cycle and the Julia set J_c admits non-trivial deformations, hence has positive area [Mc, Theorem 4.9]. A theorem of Lyubich and Shishikura then shows that f_c is (infinitely) renormalizable (see [L] or [Mc, Theorem 8.8]). It follows that W is contained in a small copy M_H for some $H \neq H_0$. So, again, $\partial W \subset M_H \subset E$. \square

Proof of Theorem 3. Let $\gamma \subset \mathcal{M}$ be an embedded arc and assume by way of contradiction that γ has positive harmonic measure. Consider the subset $\eta \subset \gamma$ consisting of the parameters which do not belong to the closure of any interior component of \mathcal{M} . By Lemma 7, the harmonic measure of the difference $\gamma \setminus \eta$ is zero, so η has positive harmonic measure. Let $X \subset \partial \mathcal{M}$ be the set given by Theorem 1. Since X has full harmonic measure, the intersection $\eta \cap X$ has positive harmonic measure. Choose three distinct parameters c, c', c'' in $\eta \cap X$, labelled so that c' is between c and c'' on γ . By Theorem 1(iii), each of these parameters forms the (trivial) impression of a unique external ray. But Lemma 6 applied to the segment of γ from c to c'' shows that c' must belong to the impression of at least two rays. This is a contradiction, and the proof is complete. \square

5. THE LAMINATION ASSOCIATED WITH \mathcal{M}

We begin with some definitions. A (geodesic) lamination in \mathbb{D} is a relatively closed set $\Lambda \subset \mathbb{D}$ decomposed into a disjoint union of complete Poincaré geodesics. A lamination Λ has a local product structure: Every point of Λ has a neighborhood U homeomorphic to the unit square $(0, 1) \times (0, 1)$ such that $\Lambda \cap U$ corresponds to $(0, 1) \times \tau$ for some relatively closed set $\tau \subset (0, 1)$, and such that the geodesic




FIGURE 3. Left: The Mandelbrot set \mathcal{M} with its main hyperbolic component H_0 and the satellite component $H_{1/3}$ at whose root the parameter rays at angles $1/7$ and $2/7$ land. The small copy $M_{H_{1/3}}$ is a proper subset of the part of \mathcal{M} cut off by these rays. Right: The lamination $\Lambda_{\mathcal{M}}$ associated with the Mandelbrot set.

segments in $\Lambda \cap U$ correspond to the horizontal segments $(0, 1) \times \{\text{point}\}$ (see for example [B]). We call U a *product neighborhood*. Λ is said to be a *product lamination* if it can be covered by a single product neighborhood.

An equivalence relation \sim on \mathbb{T} is *closed* if its graph $\{(t, s) \in \mathbb{T} \times \mathbb{T} : t \sim s\}$ is a closed subset of $\mathbb{T} \times \mathbb{T}$, and is *unlinked* if any two distinct equivalence classes are contained in a pair of disjoint intervals. Suppose \sim is closed and unlinked. Identify \mathbb{T} with the unit circle $\partial\mathbb{D}$, and form the convex hull (in the Poincaré metric of \mathbb{D}) of each equivalence class of \sim . Take the union of the topological boundaries of all these convex hulls and define Λ as the intersection of this union with the unit disk \mathbb{D} . It is not hard to check that Λ is a geodesic lamination in \mathbb{D} . We refer to it as the *lamination of \sim* .

In the case of the Mandelbrot set, define the equivalence relation $\sim_{\mathbb{Q}}$ on \mathbb{Q}/\mathbb{Z} by declaring $t \sim_{\mathbb{Q}} s$ if and only if the parameter rays $R_{\mathcal{M}}(t)$ and $R_{\mathcal{M}}(s)$ land at the same point. Extend $\sim_{\mathbb{Q}}$ to a relation \sim on the circle \mathbb{T} by taking the closure of the graph of $\sim_{\mathbb{Q}}$, that is, declare $t \sim s$ if and only if there are sequences of rational angles $t_n \rightarrow t$ and $s_n \rightarrow s$ such that $t_n \sim_{\mathbb{Q}} s_n$ for all n . It turns out that \sim is an equivalence relation, is closed and unlinked, and for $t, s \in \mathbb{Q}/\mathbb{Z}$, $t \sim s$ if and only if $t \sim_{\mathbb{Q}} s$ [D2]. The lamination $\Lambda_{\mathcal{M}}$ of this equivalence relation is called the *lamination associated with \mathcal{M}* (see Figure 3 right). Douady and Hubbard have used $\Lambda_{\mathcal{M}}$ to construct a combinatorial model of \mathcal{M} known as the “abstract Mandelbrot set” (see [DH1] or [D2]).

The following lemma will be used in the proof of Theorem 4.

Lemma 8. *Let $\Lambda \subset \mathbb{D}$ be a product lamination. If the closure $\overline{\Lambda}$ meets $\partial\mathbb{D}$ along a set of length zero, then the area of Λ must be zero.*

Proof. Since Λ is a product lamination, it is homeomorphic to $(0, 1) \times \tau$ for some relatively closed set $\tau \subset (0, 1)$. It suffices to prove the lemma when τ is compact; the general case will then follow by writing $\Lambda = \bigcup \Lambda_n$, where Λ_n is the product sub-lamination of Λ which is homeomorphic to $(0, 1) \times (\tau \cap [\frac{1}{n}, 1 - \frac{1}{n}])$. Assuming τ is compact, the closure $\bar{\Lambda}$ is homeomorphic to $[0, 1] \times \tau$, and $\bar{\Lambda} \cap \partial\mathbb{D}$ consists of two disjoint compact parts $\tau_0 \cong \{0\} \times \tau$ and $\tau_1 \cong \{1\} \times \tau$, both having length zero.

Let J_0 and J_1 be disjoint open intervals in $\partial\mathbb{D}$ containing τ_0 and τ_1 , respectively. The holonomy $h : \tau_0 \xrightarrow{\cong} \tau_1$ given by following the geodesics in Λ can be extended to a homeomorphism $h : J_0 \xrightarrow{\cong} J_1$ (for example by interpolating linearly on each component of $J_0 \setminus \tau_0$). Since both τ_0 and τ_1 have zero length, given $\varepsilon > 0$ there is an open neighborhood $V \subset J_0$ of τ_0 such that $\text{length}(V) < \varepsilon$ and $\text{length}(h(V)) < \varepsilon$. Decompose V into the disjoint union of open intervals $\{I_n\}$, so that $h(V)$ is the disjoint union of open intervals $\{h(I_n)\}$. Let $U_n \subset \bar{\mathbb{D}}$ be the convex hull of $I_n \cup h(I_n)$ in the Poincaré metric on the disk. The Euclidean area of the quadrilateral U_n is easily estimated by

$$\text{area}(U_n) \leq C (\text{length}(I_n) + \text{length}(h(I_n))),$$

where $C > 0$ is an absolute constant. Since $\Lambda \subset \bigcup U_n$, it follows that

$$\begin{aligned} \text{area}(\Lambda) &\leq \sum \text{area}(U_n) \leq C \sum (\text{length}(I_n) + \text{length}(h(I_n))) \\ &= C (\text{length}(V) + \text{length}(h(V))) \leq 2C\varepsilon, \end{aligned}$$

which shows $\text{area}(\Lambda) = 0$. □

Proof of Theorem 4. Let \sim be the equivalence relation used in the definition of $\Lambda_{\mathcal{M}}$. If $t \sim s$ and $t \neq s$, there exist sequences of rational angles $t_n \rightarrow t$ and $s_n \rightarrow s$ such that $t_n \sim_{\mathbb{Q}} s_n$ for all n . Let z_n be the common landing point of $R_{\mathcal{M}}(t_n)$ and $R_{\mathcal{M}}(s_n)$. Then, by (2.1), any accumulation point of the sequence $\{z_n\}$ must belong to both impressions $I_{\mathcal{M}}(t)$ and $I_{\mathcal{M}}(s)$; in particular, $I_{\mathcal{M}}(t) \cap I_{\mathcal{M}}(s) \neq \emptyset$. If $X \subset \partial\mathcal{M}$ is the set given by Theorem 1, it follows that every equivalence class of \sim containing more than one element must be disjoint from $\psi_{\mathcal{M}}^{-1}(X)$. This means that the set of endpoints of the geodesics in $\Lambda_{\mathcal{M}}$ is a subset of $\mathbb{T} \setminus \psi_{\mathcal{M}}^{-1}(X)$, which has length zero.

Let U be any product neighborhood for $\Lambda_{\mathcal{M}}$ and let Λ be the union of all geodesics in $\Lambda_{\mathcal{M}}$ which intersect U . Then Λ is a product lamination, and by the preceding paragraph $\bar{\Lambda} \cap \partial\mathbb{D}$ has length zero. By Lemma 8, $\text{area}(\Lambda) = 0$. Since $\Lambda \supset \Lambda_{\mathcal{M}} \cap U$, we conclude that $\text{area}(\Lambda_{\mathcal{M}} \cap U) = 0$. Since $\Lambda_{\mathcal{M}}$ can be covered by countably many such product neighborhoods U , we conclude that $\text{area}(\Lambda_{\mathcal{M}}) = 0$.

Now consider the set $\psi_{\mathcal{M}}^{-1}(\mathbb{R})$ of angles t for which the parameter ray $R_{\mathcal{M}}(t)$ lands on the real line. It is shown in [Z2] that $\psi_{\mathcal{M}}^{-1}(\mathbb{R})$ is a nowhere dense set of Hausdorff dimension 1, and that $t \sim -t$ for every $t \in \psi_{\mathcal{M}}^{-1}(\mathbb{R})$. It easily follows that the union over all $t \in \psi_{\mathcal{M}}^{-1}(\mathbb{R})$ of the Poincaré geodesic in \mathbb{D} joining t to $-t$ has Hausdorff dimension 2. Since this union is contained in $\Lambda_{\mathcal{M}}$, we conclude that the dimension of $\Lambda_{\mathcal{M}}$ must be 2. □

ACKNOWLEDGMENTS

I am indebted to Jacek Graczyk who brought the results of [GS] to my attention, which prompted the writing of this note. I am also grateful to John Milnor for his useful comments on an earlier draft of the paper.

REFERENCES

- [A] L. Ahlfors, *Conformal invariants*, McGraw-Hill, 1973. MR0357743 (50:10211)
- [BS] J. Birman and C. Series, *Geodesics with bounded intersection number on surfaces are sparsely distributed*, *Topology*, **24** (1985) 217-225. MR0793185 (87f:57012)
- [B] F. Bonahon, *Geodesic laminations on surfaces*, in "Laminations and foliations in dynamics, geometry and topology," *Contemp. Math.*, **269**, Amer. Math. Soc. 2001, 1-37. MR1810534 (2001m:57023)
- [BSc] H. Bruin and D. Schleicher, *Symbolic dynamics of quadratic polynomials*, Mittag-Leffler Institute preprints, 2001/2002, No. 07.
- [D1] A. Douady, *Algorithms for computing angles in the Mandelbrot set*, in "Chaotic Dynamics and Fractals," Academic Press, 1986, 155-168. MR0858013
- [D2] A. Douady, *Descriptions of compact sets in \mathbb{C}* , in "Topological methods in modern mathematics," Publish or Perish, 1993, 429-465. MR1215973 (94g:58185)
- [DH1] A. Douady and J. Hubbard, *Etude dynamique de polynomes complexes*, *Publications Math. d'Orsay* **84-02** (1984) (premiere partie) and **85-04** (1985) (deuxieme partie). MR0762431 (87f:58072a) MR0812271 (87f:58072b)
- [DH2] A. Douady and J. Hubbard, *On the dynamics of polynomial-like mappings*, *Ann. Sci. Ec. Norm. Sup.*, **18** (1985) 287-343. MR0816367 (87f:58083)
- [GM] L. Goldberg and J. Milnor, *Fixed points of polynomial maps II*, *Ann. Sci. Ec. Norm. Sup.*, **26** (1993) 51-98. MR1209913 (95d:58107)
- [GS] J. Graczyk and G. Świątek, *Harmonic measure and expansion on the boundary of the connectedness locus*, *Invent. Math.*, **142** (2000) 605-629. MR1804163 (2001k:37066)
- [H] J. Hubbard, *Local connectivity of Julia sets and bifurcation loci*, in "Topological Methods in Modern Mathematics," Publish or Perish, 1993, 457-511. MR1215974 (94c:58172)
- [L] M. Lyubich, *On the Lebesgue measure of the Julia set of a quadratic polynomial*, Stony Brook IMS preprint # 1991/10.
- [Ma] A. Manning, *Logarithmic capacity and renormalizability for landing on the Mandelbrot set*, *Bull. London Math. Soc.*, **28** (1996) 521-526. MR1396155 (97g:58141)
- [Mc] C. McMullen, *Complex Dynamics and Renormalization*, *Annals of Math Studies* **135**, Princeton University Press, 1994. MR1312365 (96b:58097)
- [M1] J. Milnor, *Dynamics in one complex variable: Introductory lectures*, Vieweg, 1999. MR1721240 (2002i:37057)
- [M2] J. Milnor, *Local connectivity of Julia sets: expository lectures*, in "The Mandelbrot set, Theme and Variations," LMS Lecture Note Series 274, Cambr. U. Press 2000, 67-116. MR1765085 (2001b:37073)
- [M3] J. Milnor, *Periodic orbits, external rays and the Mandelbrot set: An expository account*, *Astérisque*, **261** (2000) 277-333. MR1755445 (2002e:37067)
- [M4] J. Milnor, *Pasting together Julia sets: a worked out example of mating*, *Experiment. Math.*, **13** (2004) 55-92. MR2065568 (2005c:37087)
- [P] C. Pommerenke, *Boundary behaviour of conformal maps*, Springer-Verlag, 1992. MR1217706 (95b:30008)
- [R] P. Roesch, *Holomorphic motions and puzzles*, in "The Mandelbrot set, Theme and Variations," LMS Lecture Note Series 274, Cambr. U. Press 2000, 117-131. MR1765086 (2001c:37046)
- [S1] S. Smirnov, *Symbolic dynamics and the Collet-Eckmann condition*, *Internat. Math. Res. Notices*, **7** (2000) 333-351. MR1749741 (2001i:37071)
- [S2] S. Smirnov, *On support of dynamical laminations and biaccessible points in Julia set*, *Colloq. Math.*, **87** (2001) 287-295. MR1814670 (2001m:37092)
- [T1] Tan Lei, *Voisinages connexes des points de Misiurewicz*, *Ann. Inst. Fourier Grenoble*, **42** (1992) 707-735. MR1196091 (94a:58165)
- [T2] Tan Lei, *Local properties of the Mandelbrot set at parabolic points*, in "The Mandelbrot set, Theme and Variations," LMS Lecture Note Series 274, Cambr. U. Press 2000, 133-160. MR1765087 (2001g:37065)
- [Th] W. Thurston, *The geometry and topology of 3-manifolds*, Lecture notes, Princeton University, 1976-1979.
- [Z1] S. Zakeri, *Biaccessibility in quadratic Julia sets*, *Ergod. Th. & Dyn. Sys.*, **20** (2000) 1859-1883. MR1804961 (2001k:37068)

- [Z2] S. Zakeri, *External rays and the real slice of the Mandelbrot set*, *Ergod. Th. & Dyn. Sys.*, **23** (2003) 637-660. MR1972243 (2004a:37057)
- [Zd] A. Zdunik, *On biaccessible points in Julia sets of polynomials*, *Fund. Math.*, **163** (2000) 277-286. MR1758329 (2001f:37058)

INSTITUTE FOR MATHEMATICAL SCIENCES, STONY BROOK UNIVERSITY, STONY BROOK, NEW YORK 11794

Current address: Department of Mathematics, Queens College of CUNY, Flushing, New York 11367

E-mail address: `zakeri@forbin.qc.edu`