

CHARACTERS OF p' -DEGREE WITH CYCLOTOMIC FIELD OF VALUES

GABRIEL NAVARRO AND PHAM HUU TIEP

(Communicated by Jonathan I. Hall)

ABSTRACT. If p is a prime number and G is a finite group, we show that G has an irreducible complex character of degree not divisible by p with values in the cyclotomic field \mathbb{Q}_p .

1. INTRODUCTION

R. Gow conjectured that every finite group of even order has a nontrivial irreducible complex character with odd degree and rational values. This conjecture was finally proven in [7]. In this note we seek an analog of this result which works for every prime p . If G is a finite group and $\chi \in \text{Irr}(G)$ is an irreducible complex character of G , we denote by $\mathbb{Q}(\chi)$ the field of values of χ . Also, we let \mathbb{Q}_n be the cyclotomic field generated by a primitive n th root of unity.

Theorem A. *Let p be a prime and let G be a finite group of order divisible by p . Then there exists a nontrivial $\chi \in \text{Irr}(G)$ of p' -degree such that $\mathbb{Q}(\chi) \subseteq \mathbb{Q}_p$.*

Of course, the $p = 2$ case of Theorem B is Gow's conjecture. We are able, in fact, to prove the following.

Theorem B. *Let p be a prime and let G be a finite group. Then the trivial character is the only p' -degree irreducible character of G with values in \mathbb{Q}_p if and only if G is a group of odd order not divisible by p .*

Once we have that groups of order divisible by p have nontrivial irreducible p' -degree characters with values in \mathbb{Q}_p , it is natural to ask if we can even obtain them having rational values. Of course, some conditions are necessary (since odd p -groups certainly do not possess these characters, nor does $L_2(3^{2a+1})$ for $p = 3$). The following improves on some of the results in [5].

Theorem C. *Let G be a finite group and let p be a prime.*

- (i) *Let G be nonsolvable. Assume that either $p \neq 3$, or that $p = 3$ and G has no composition factor isomorphic to $L_2(3^{2a+1})$ for any $a \geq 1$. Then G has a nontrivial, rational, irreducible character of p' -degree.*

Received by the editors April 22, 2005.

2000 *Mathematics Subject Classification.* Primary 20C15.

The first author was partially supported by the Ministerio de Educación y Ciencia proyecto MTM2004-06067-C02-01.

The second author gratefully acknowledges the support of the NSA (grant H98230-04-0066).

©2006 American Mathematical Society
Reverts to public domain 28 years from publication

- (ii) Let G be solvable. Then G has a nontrivial, rational, irreducible character of p' -degree if and only if $|\mathbf{N}_G(P)|$ is even, where $P \in \text{Syl}_p(G)$.

Part of this work was done while the first author visited the University of Florida, and he would like to thank the Mathematics Department for its hospitality.

2. SIMPLE GROUPS

The proof of our main results depends on the Classification of Finite Simple Groups.

(2.1) Theorem. *Let p be an odd prime and let G be a finite nonabelian simple group.*

- (i) *There exists a nontrivial $\chi \in \text{Irr}(G)$ of p' -degree such that $\mathbb{Q}(\chi) \subseteq \mathbb{Q}_p$.*
(ii) *If $p = 3$, assume in addition that G is not isomorphic to $L_2(3^{2a+1})$ for any $a \geq 1$*

Then there exists a nontrivial rational-valued $\chi \in \text{Irr}(G)$ of p' -degree. Furthermore, if a p -group Q acts on G , then χ may be chosen to be Q -invariant.

Proof. If G is an sporadic simple group, then a check of the ATLAS [2] gives us a desired χ , as Q must act on G only via inner automorphisms. Similarly, if $G = A_n$ is an alternating group with $n \geq 5$, as pointed out in Lemma (6.3) of [7], we have that G has two irreducible rational-valued characters of degree $d > 1$ and $d + 1$. One of these two characters has degree not divisible by p and is Q -invariant, by the same reason as before, and this case follows. Furthermore, if G is a finite group of Lie type in characteristic $\ell \neq p$, then we can take χ to be the Steinberg character. So we may assume that G is a finite group of Lie type in characteristic p .

Assume $G = L_2(q)$. If $q \neq 3^{2a+1}$, then the proof of Lemma (6.4) of [7] shows that G has a rational irreducible character of degree $q \pm 1$ which is stable under field automorphisms of G , and so we are done. On the other hand, if $q = 3^{2a+1}$, then G has two so-called Weil characters of degree $(q - 1)/2$ with character value field equal to \mathbb{Q}_3 . Thus we may assume G is not of type A_1 .

There is a simple simply connected algebraic group \mathcal{G} over the algebraic closure \mathbb{F} of \mathbb{F}_p and a Frobenius map F on \mathcal{G} such that $G = \mathcal{G}^F/\mathbf{Z}(\mathcal{G}^F)$. Let (\mathcal{G}^*, F^*) be dual to (\mathcal{G}, F) . For every conjugacy class $[s]$ of a semisimple element $s \in \mathcal{G}^{F^*}$ such that $\mathbf{C}_{\mathcal{G}^*}(s)$ is connected, there is a so-called semisimple irreducible character $\chi_s \in \text{Irr}(\mathcal{G}^F)$ of degree coprime to p [1], [3], [6]. Furthermore, χ_s belongs to the Lusztig series corresponding to $[s]$. Clearly, \mathcal{G}^F has a unique irreducible character of degree 1 — the trivial character — and it belongs to the Lusztig series corresponding to [1]. Since Lusztig series are disjoint, we see that $\chi_s(1) > 1$ if $s \neq 1$. If s is contained in the derived subgroup of \mathcal{G}^{*F^*} , then $\mathbf{Z}(\mathcal{G}^F) \leq \ker \chi_s$, and so χ_s defines an irreducible character of G . Moreover, by [6], this character is invariant under $\text{Aut}(G) = \text{Aut}(\mathcal{G}^F)$ provided that the class $[s]$ is invariant under $\text{Aut}(\mathcal{G}^{*F^*})$ (and $\mathbf{C}_{\mathcal{G}^*}(s)$ is connected as we assumed above). Finally, by Lemma (6.1) of [7], χ_s is rational-valued if s is rational in \mathcal{G}^{*F^*} . Hence it suffices to find a nontrivial rational semisimple element s in \mathcal{G}^{*F^*} , lying in the derived subgroup of \mathcal{G}^{*F^*} , having connected centralizer in \mathcal{G}^* , and such that its conjugacy class is invariant under $\text{Aut}(\mathcal{G}^{*F^*})$. This element s has indeed been exhibited in the proof of Theorem (4.1) of [5]. \square

3. PROOF OF THEOREM B

If $N \triangleleft G$ and $\theta \in \text{Irr}(N)$, we write $\text{Irr}(G|\theta)$ for the irreducible characters χ of G having θ as an irreducible constituent of χ_N . Recall that if $\chi \in \text{Irr}(G)$ and $\sigma \in \text{Gal}(\mathbb{Q}(\chi)/\mathbb{Q})$, then $\chi^\sigma \in \text{Irr}(G)$ and $\mathbb{Q}(\chi) = \mathbb{Q}(\chi^\sigma)$. We need the following lemma.

(3.1) Lemma. *Suppose that $N \triangleleft G$ and that $\theta \in \text{Irr}(N)$ is G -invariant with $\mathbb{Q}(\theta) \subseteq \mathbb{Q}_{p^e}$, where $p > 2$ is prime and $e \geq 1$ is some integer. If G/N is an odd p' -group, then there exists a unique $\tau \in \text{Irr}(G|\theta)$ such that $\mathbb{Q}(\tau) \subseteq \mathbb{Q}_{p^f}$, where $f \geq 1$ is some integer. In fact, $\tau_N = \theta$.*

Proof. We argue by induction on $|G : N|$. By using the fact that G/N is solvable, we let $N \subseteq M \triangleleft G$ with $|G : M| = q$, where $2 \neq q \neq p$ is a prime. By induction, there exists a unique $\psi \in \text{Irr}(M)$ over θ with values in some cyclotomic field \mathbb{Q}_{p^f} . In fact, $\psi_M = \theta$. Now, if $g \in G$, then ψ^g lies over θ and has values in \mathbb{Q}_{p^f} . By uniqueness, we have that $\psi = \psi^g$. Now, ψ is q -rational, and by Theorem (6.30) of [4] there exists a unique q -rational extension $\tau \in \text{Irr}(G)$ of ψ to G . Now, $\mathbb{Q}(\psi) \subseteq \mathbb{Q}(\tau)$. If $\sigma \in \text{Gal}(\mathbb{Q}(\tau)/\mathbb{Q}(\psi))$, it follows that $\tau^\sigma \in \text{Irr}(G)$ is a q -rational character extending ψ . By uniqueness, $\tau^\sigma = \tau$, and we deduce that $\mathbb{Q}(\tau) = \mathbb{Q}(\psi) \subseteq \mathbb{Q}_{p^f}$. Suppose now that $\rho \in \text{Irr}(G|\theta)$ has its values in \mathbb{Q}_{p^r} for some integer $r \geq 1$. By Gallagher's Corollary (6.17) of [4], we have that $\rho = \xi\tau$, where $\xi \in \text{Irr}(G/N)$. We claim that $\xi = 1$. Since G/N is odd, by Burnside's theorem it suffices to show that ξ is rational valued. Now, let $\alpha \in \text{Gal}(\mathbb{Q}(\xi)/\mathbb{Q})$. Since G/N is a p' -group, we may extend α to some $\beta \in \text{Gal}(\mathbb{Q}_{|G|}/\mathbb{Q})$ that fixes all p -power roots of unity. In particular, θ , ρ and τ are β -fixed. By the uniqueness in Corollary (6.17) of [4], we have that ξ is β -fixed, and therefore α -fixed. This proves the lemma. □

If $K \subseteq \mathbb{C}$ is a field, let us denote by $\text{Irr}_{p',K}(G)$ the set of all irreducible complex characters of G with degree not divisible by p and with values in K . The following is Theorem B of the Introduction.

(3.2) Theorem. *Let p be a prime and let G be a finite group. Then $|\text{Irr}_{p',\mathbb{Q}_p}(G)| = 1$ if and only if G has odd order not divisible by p .*

Proof. Suppose that G has odd order not divisible by p , and let $\chi \in \text{Irr}(G)$ with values in \mathbb{Q}_p . Then χ has its values in $\mathbb{Q}_{|G|} \cap \mathbb{Q}_p = \mathbb{Q}$. Since G has odd order, we conclude that $\chi = 1$ by Burnside's theorem.

Now, assume that $|\text{Irr}_{p',\mathbb{Q}_p}(G)| = 1$. We prove that G is of odd order not divisible by p , by arguing by induction on $|G|$. By Theorem B of [7], we may assume that p is odd and that $|G|$ is divisible by p . Now, let $1 < N$ be a minimal normal subgroup of G . Hence, by induction, we have that G/N is of odd order not divisible by p . Suppose first that N is abelian. If N is of odd order not divisible by p , then we are done. So we may assume that N is an elementary abelian 2-group or p -group. In any case, $(|G : N|, |N|) = 1$. Now, let $1 \neq \lambda \in \text{Irr}(N)$ and let T be the stabilizer of λ in G . By Corollary (8.16) of [4], there exists a unique extension $\hat{\lambda} \in \text{Irr}(T)$ of λ with determinantal order $o(\hat{\lambda}) = o(\lambda)$. Hence $\mathbb{Q}(\hat{\lambda}) = \mathbb{Q}(\lambda) \subseteq \mathbb{Q}_p$ (in both cases). Now, by using the Clifford correspondence, we have that $1 \neq \chi = \hat{\lambda}^G \in \text{Irr}(G)$ has its values in \mathbb{Q}_p and has p' -degree.

So we may assume that N is a direct product of nonabelian isomorphic simple groups of order divisible by p . Now, by Theorem (2.1), there exists $1 \neq \theta \in \text{Irr}(N)$

of p' -degree with values in \mathbb{Q}_p . Let I be the stabilizer of θ in G . By Lemma (3.1), there exists $\rho \in \text{Irr}(I)$ over θ with values in \mathbb{Q}_p . Now $1 \neq \rho^G \in \text{Irr}(G)$ has p' -degree and has its values in \mathbb{Q}_p . \square

4. RATIONAL CHARACTERS OF p' -DEGREE

We need the following lemma.

(4.1) Lemma. *Suppose that $N \triangleleft G$, where G/N is solvable, $p > 2$ is prime, $P/N \in \text{Syl}_p(G/N)$ and $|\mathbf{N}_{G/N}(P/N)|$ is odd. If $\theta \in \text{Irr}(N)$ is rational, P -invariant of p' -degree, then there exists $\chi \in \text{Irr}(G|\theta)$ rational of p' -degree.*

Proof. We argue by induction on $|G : N|$. Let M/N be a chief factor of G . If M/N is a p -group, then θ is M -invariant and has a unique real extension $\eta \in \text{Irr}(M)$ by Lemma (2.1) of [7]. By uniqueness, η is rational and P -invariant. Since $|G : M| < |G : N|$, then the lemma follows by induction. Assume now that M/N is an odd p' -group. By Corollary (2.2) of [7], there is a unique rational $\eta \in \text{Irr}(M|\theta)$. Again by uniqueness, η is P -invariant, and we note that η has p' -degree since M/N is a p' -group (by Corollary (11.29) of [4]). Hence, the lemma again follows by induction. Finally, assume that M/N is a 2-group. By hypothesis, we have that $\mathbf{C}_{M/N}(P) = 1$. By Problem (13.10) of [4], we have that there exists a unique P -invariant $\eta \in \text{Irr}(M|\theta)$. By uniqueness, η is P -invariant, rational, of p' -degree, and we may apply induction once again. \square

Suppose that G is a p -solvable group, and let P be a Sylow p -subgroup of G . In [5], it was proven that $|\text{Irr}_{p',\mathbb{R}}(G)| = 1$ if and only if $|\mathbf{N}_G(P)|$ is odd. Combined with this result, the following theorem yields Theorem C.

(4.2) Theorem. *Let G be a finite group and let $P \in \text{Syl}_p(G)$, where p is prime. If $p = 3$, assume in addition that G has no composition factor isomorphic to $L_2(3^{2a+1})$ for any $a \geq 1$. Suppose that $|\text{Irr}_{p',\mathbb{Q}}(G)| = 1$. Then $|\mathbf{N}_G(P)|$ is odd and G is solvable.*

Proof. If $p = 2$, then by Theorem (5.2) of [7], we see that $|G|$ is odd, and so we are done. We assume, therefore, that $p \neq 2$. We argue by induction on $|G|$.

Let N be minimal normal in G . Then $|\text{Irr}_{p',\mathbb{Q}}(G/N)| = 1$ and by the induction hypothesis, G/N is solvable and $|\mathbf{N}_{G/N}(PN/N)|$ is odd. Since $\mathbf{N}_{G/N}(PN/N) = \mathbf{N}_G(P)N/N$, it follows that $|\mathbf{N}_G(P) : \mathbf{N}_G(P) \cap N|$ is odd. Hence, it suffices to show that $|\mathbf{N}_G(P) \cap N|$ is odd and that N is solvable. In particular, since $p \neq 2$, we may assume that N is not a p -group.

Assume that N is a p' -group. We claim that $|\mathbf{C}_N(P)|$ is odd. Otherwise, $|\mathbf{C}_N(P)|$ is even, and by Theorem (5.2) of [7] $\mathbf{C}_N(P)$ has a nontrivial rational $\eta \in \text{Irr}(\mathbf{C}_N(P))$. Now, let $1 \neq \theta \in \text{Irr}(N)$ be P -invariant such that the Glauberman correspondent of θ is η . Since the Glauberman correspondence commutes with Galois action, it follows that θ is rational, of p' -degree and P -invariant. By Lemma (4.1), there exists $\chi \in \text{Irr}(G|\theta)$ rational of p' -degree, a contradiction. Since $|\mathbf{C}_N(P)|$ is odd, N is solvable by Theorem (3.4) of [5]. Furthermore, $\mathbf{N}_G(P) \cap N = \mathbf{C}_N(P)$, so we are done in this case.

Finally, assume that the minimal normal subgroup N of G is neither a p -group nor a p' -group. Then $N = K_1 \times \cdots \times K_k$, where each K_i is a product of nonabelian simple groups transitively permuted by P . Write $K_1 = T_1 \times \cdots \times T_r$, where the simple groups T_i are transitively permuted by P . Suppose that $Q = \mathbf{N}_P(T_1)$. By

our assumption, $T_1 \not\cong L_2(3^{2a+1})$ for any $a \geq 1$. By Theorem (2.1(ii)), there exists $1 \neq \phi \in \text{Irr}(T_1)$ rational of p' -degree and Q -invariant. For each subscript i with $1 \leq i \leq r$, choose $x_i \in P$ so that $(T_1)^{x_i} = T_i$, and let $\phi_i = \phi^{x_i} \in \text{Irr}(T_i)$. It is straightforward to check that $1 \neq \eta = \phi_1 \times \cdots \times \phi_r$ is P -invariant, rational of p' -degree. Now, consider η as an irreducible character of N (with $K_2 \cdots K_k$ in its kernel) and apply Lemma (4.1) to get a nontrivial rational p' -degree character of G . This final contradiction completes the proof of the theorem. \square

REFERENCES

- [1] R. Carter, '*Finite Groups of Lie type: Conjugacy Classes and Complex Characters*', Wiley, Chichester, 1985. MR0794307 (87d:20060)
- [2] J. H. Conway, R. T. Curtis, S. P. Norton, R. A. Parker, and R. A. Wilson, '*An ATLAS of Finite Groups*', Clarendon Press, Oxford, 1985. MR0827219 (88g:20025)
- [3] F. Digne and J. Michel, '*Representations of Finite Groups of Lie Type*', London Mathematical Society Student Texts 21, Cambridge University Press, 1991. MR1118841 (92g:20063)
- [4] I. M. Isaacs, '*Character Theory of Finite Groups*', Dover, New York, 1994. MR1280461
- [5] I. M. Isaacs, G. Malle, and G. Navarro, Real characters of p' -degree, *J. Algebra* **278** (2004), 611–620. MR2071655 (2005f:20016)
- [6] G. Lusztig, On the representations of reductive groups with disconnected centre, in: '*Orbites Unipotentes et Représentations, I.*', *Astérisque* **168** (1988), 157–166. MR1021495 (90j:20083)
- [7] G. Navarro and Pham Huu Tiep, Rational irreducible characters and rational conjugacy classes in finite groups, submitted.

FACULTAT DE MATEMÀTIQUES, UNIVERSITAT DE VALÈNCIA, BURJASSOT, VALÈNCIA 46100, SPAIN
E-mail address: gabriel@uv.es

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF FLORIDA, GAINESVILLE, FLORIDA 32611
E-mail address: tiep@math.ufl.edu