

NO EMBEDDINGS OF SOLENOIDS INTO SURFACES

BOJU JIANG, SHICHENG WANG, AND HAO ZHENG

(Communicated by Alexander N. Dranishnikov)

ABSTRACT. A quick proof of Bing's theorem indicated by the title is given. Indeed the inverse limit of a sequence of degree > 1 maps between closed oriented m -manifolds can never be embedded into any closed orientable $(m + 1)$ -manifold. The proof also concludes Gumerov's result on the covering of solenoids.

In this paper we always assume that a solenoid is not a manifold.

In his two important papers on solenoids, Bing proved first that no solenoid is planar [B1] and then if a solenoid can be embedded into surfaces, then it must be planar [B2], and therefore no solenoid can be embedded into surfaces. We will give a short proof of this result. Indeed we show that the inverse limit of a sequence of maps of degree > 1 between closed oriented m -manifolds can never be embedded into any closed orientable $(m + 1)$ -manifold. The proof also concludes a recent result of [Gu] on covering degrees of solenoids.

For the sequence $\{\phi_n : X_n \rightarrow X_{n-1}\}_{n \geq 1}$, its *inverse limit* $\varprojlim X_n$ is defined as the subspace

$$\{(x_0, x_1, \dots, x_n, \dots) \mid x_n \in X_n, x_{n-1} = \phi_n(x_n)\}$$

of the product space $\prod_{n=0}^{\infty} X_n$.

Identify S^1 with the abelian Lie group $U_1 = \{z \in \mathbb{C} \mid |z| = 1\}$ and let $\phi_n : S^1 \rightarrow S^1$ be the homomorphism defined by $z \mapsto z^{w_n}$ where $w_n > 1$ is an integer. For the sequence $\{\phi_n : S^1 \rightarrow S^1\}_{n \geq 1}$, its inverse limit, denoted as Θ , is called the *solenoid* of type $\varpi = (w_1, w_2, \dots, w_n, \dots)$. By definition Θ is a connected, closed and hence compact subgroup of the abelian topological group $\prod_{n=0}^{\infty} S^1$ (cf. [Mc]).

Theorem 1. (1) *For a sequence $\{\phi_n : X_n \rightarrow X_{n-1}\}_{n \geq 1}$ of non-zero degree maps of closed oriented m -manifolds having infinitely many degree $\phi_n \neq \pm 1$, there is no embedding of $\varprojlim X_n$ into any closed oriented $(m + 1)$ -manifolds M .*

(2) *No solenoid Θ , can be embedded into a surface.*

Remark 2. In general, we call the inverse limit of maps $\{\phi_n : X_n \rightarrow X_{n-1}\}_{n \geq 1}$ an *m -dimensional solenoid* if all X_n are closed oriented m -manifolds and all ϕ_n are coverings of degree > 1 . By Theorem 1 (1), there is no embedding of m -dimensional solenoids into \mathbb{R}^{m+1} . This fact has been proved for homogeneous solenoids by Prajs [Pr]. Note that solenoids of dimension > 1 need not be homogeneous unless the compositions of all coverings are regular.

Received by the editors November 2, 2006, and, in revised form, August 9, 2007.

2000 *Mathematics Subject Classification.* Primary 54F15, 57N35.

The authors were supported by an NSFC grant.

Theorem 3. *If X is a finite-fold covering of a solenoid Θ , then each component of X is homeomorphic to Θ .*

Remark 4. According to the description in [Gu, p. 2775], it seems Theorem 3 can be derived from several papers of Fox and Moore in the 1970's, and of Grigorian and Gumerov recently. We will give a direct proof of Theorem 3.

Proof of Theorem 1. (1) Suppose there is an embedding $\varprojlim X_n \subset M$ for some closed orientable $(m+1)$ -manifold M . We have to seek a contradiction.

According to the continuity of Čech theory, for any commutative ring R , the Čech cohomology of the inverse limit is the direct limit of the singular cohomology, that is, $\check{H}^*(\varprojlim X_n; R) = \varinjlim H^*(X_n; R)$. See [ES, Chapter X, Theorems 2.2 and 3.1].

Clearly we have $H^m(X_n; R) = R$. Denote the degree of ϕ_n by w_n . Then the Čech cohomology group $\check{H}^m(\varprojlim X_n; R)$ is the direct limit of $R \xrightarrow{w_1} R \xrightarrow{w_2} R \xrightarrow{w_3} R \xrightarrow{w_4} \dots$. Since all $w_n \neq 0$ and infinitely many $w_n \neq \pm 1$, it follows that $\check{H}^m(\varprojlim X_n; \mathbb{Z})$ is an infinitely generated \mathbb{Z} -module, but $\check{H}^m(\varprojlim X_n; \mathbb{Q}) \cong \mathbb{Q}$ is a finitely generated \mathbb{Q} -module.

By the Alexander duality $\check{H}^m(\varprojlim X_n; R) \cong H_1(M, M \setminus \varprojlim X_n; R)$ (cf. [GH, p. 233]), we have the exact sequence

$$\dots \rightarrow H_1(M; R) \rightarrow \check{H}^m(\varprojlim X_n; R) \rightarrow H_0(M \setminus \varprojlim X_n; R) \rightarrow H_0(M; R) \rightarrow \dots$$

Since $H_*(M; R)$ is a finitely generated R -module for any commutative ring R , the above exact sequence implies that $H_0(M \setminus \varprojlim X_n; \mathbb{Z})$ is an infinitely generated \mathbb{Z} -module and $H_0(M \setminus \varprojlim X_n; \mathbb{Q})$ is a finitely generated \mathbb{Q} -module. But on the other hand $H_0(M \setminus \varprojlim X_n; \mathbb{Z})$ is a free abelian group and $H_0(M \setminus \varprojlim X_n; \mathbb{Q}) = H_0(M \setminus \varprojlim X_n; \mathbb{Z}) \otimes_{\mathbb{Z}} \mathbb{Q}$, so $H_0(M \setminus \varprojlim X_n; \mathbb{Z})$ and $H_0(M \setminus \varprojlim X_n; \mathbb{Q})$ should have the same basis, a contradiction.

(2) Suppose there is an embedding $\Theta \subset F$ of a solenoid Θ into a surface F . First, we may assume F is a closed surface. If not, since Θ is compact, Θ is contained in the interior of some connected, compact subsurface $F' \subset F$. Capping a disc on each component of $\partial F'$, we get an embedding of Θ into a closed surface.

According to Theorem 1 (1), F cannot be orientable. If F is not orientable, we consider the orientable double covering $\pi: \tilde{F} \rightarrow F$. Since $\pi^{-1}(\Theta)$ is a double covering of Θ , by Theorem 3 each component of $\pi^{-1}(\Theta)$ is homeomorphic to Θ ; hence we get an embedding $\Theta \subset \tilde{F}$ into a closed, orientable surface, which still contradicts Theorem 1 (1). \square

Proof of Theorem 3. Suppose Θ is a solenoid of type $\varpi = (w_1, w_2, \dots, w_n, \dots)$ and $\pi: X \rightarrow \Theta$ is an r -fold covering.

Let Γ_n be the kernel of the projection $p_n: \Theta \rightarrow S^1$ onto the n th coordinate, i.e.

$$\Gamma_n = \{(x_0, x_1, \dots, x_n, \dots) \in \Theta \mid x_0 = x_1 = \dots = x_n = 1\}.$$

We have the infinite sequence of closed, and hence compact, subgroups (indeed each Γ_n is homeomorphic to the Cantor set; cf. [Mc])

$$\Theta > \Gamma_0 > \Gamma_1 > \Gamma_2 > \dots > \Gamma_n > \dots$$

The right multiplication of Γ_n on Θ makes the projection $p_n : \Theta \rightarrow S^1$ a principal fiber bundle (cf. [Mc]). It follows that Θ is the mapping torus of a left transformation $\psi_n : \Gamma_n \rightarrow \Gamma_n$. We choose the left transformation ψ_n as follows. Fix a closed path $\alpha(t) = e^{2\pi it}$, $t \in [0, 1]$, in S^1 . Then $\alpha_n(t) = \gamma(w_1 \cdots w_n t)$ is a lift of α via the projection p_n , where

$$\gamma(t) = (e^{2\pi it}, e^{2\pi it/w_1}, e^{2\pi it/w_1 w_2}, \dots, e^{2\pi it/w_1 \cdots w_n}, \dots)$$

is the one-parameter subgroup of Θ . Let ψ_n be given by $x \mapsto \alpha_n(1)x$.

Moreover, as the covering space of Θ , X is the mapping torus of a lift $\tilde{\psi}_n$ of ψ_n which is uniquely determined as follows. For each $\tilde{x} \in \pi^{-1}(\Gamma_n)$, let $\tilde{\alpha}$ be the unique lift starting from \tilde{x} of the path $\alpha_n(t)\pi(\tilde{x})$ in Θ via the covering map π ; then $\tilde{\psi}_n(\tilde{x}) = \tilde{\alpha}(1)$. Note that $\psi_n = \psi_0^{w_1 \cdots w_n}|_{\Gamma_n}$ and hence $\tilde{\psi}_n = \tilde{\psi}_0^{w_1 \cdots w_n}|_{\pi^{-1}(\Gamma_n)}$.

Claim 1. Let \mathcal{U} be an open covering of Γ_0 . Then for sufficiently large n each coset of Γ_n in Γ_0 is contained in some element of \mathcal{U} .

Proof. Note that $\mathcal{B} = \{x\Gamma_n \mid n \geq 0, x \in \Gamma\}$ forms a basis for the open sets of Γ_0 . So the covering \mathcal{U} has a refinement \mathcal{U}' which consists of elements of \mathcal{B} . Since Γ_0 is compact we may assume \mathcal{U}' is finite. Therefore, for sufficiently large n each coset of Γ_n in Γ_0 is contained in some element of \mathcal{U}' , hence in some element of \mathcal{U} . \square

Claim 2. We have $\pi^{-1}(\Gamma_0) \cong \Gamma_0 \times \{1, 2, \dots, r\}$ as covering spaces of Γ_0 .

Proof. Let \mathcal{U} be the open covering of Γ_0 which consists of the open fundamental regions of the covering map $\pi|_{\pi^{-1}(\Gamma_0)}$ and let n be sufficiently large as in Claim 1. Then each coset $x\Gamma_n$ in Γ_0 is contained in some element \mathcal{U} ; hence $\pi^{-1}(x\Gamma_n) \cong x\Gamma_n \times \{1, 2, \dots, r\}$ as covering spaces of $x\Gamma_n$. Since the cosets of Γ_n are disjoint and open in Γ_0 , the claim follows. \square

In what follows we fix a homeomorphism from Claim 2 and identify both sets.

Claim 3. For each sufficiently large n there exists a permutation σ_n of $\{1, 2, \dots, r\}$ such that $\tilde{\psi}_n(x, j) = (\psi_n(x), \sigma_n(j))$.

Proof. Applying Claim 1 on the the open covering of Γ_0 which consists of the open (and closed) sets $U_\sigma = \{x \in \Gamma_0 \mid \tilde{\psi}_0(x, j) = (\psi_0(x), \sigma(j))\}$ with σ running over all permutations of $\{1, 2, \dots, r\}$, one notices that for sufficiently large n each coset of Γ_n in Γ_0 is contained in some U_σ . Since $\tilde{\psi}_n = \tilde{\psi}_0^{w_1 \cdots w_n}|_{\pi^{-1}(\Gamma_n)}$, the claim follows. \square

By Claim 3 X is the disjoint union of the mapping tori of $\tilde{\psi}_n|_{\Gamma_n \times J}$ where J runs over all σ_n -orbits. Since Θ is connected, the following two claims eventually establish the theorem. Below we denote by T_f the mapping torus of a self homeomorphism f .

Claim 4. For sufficiently large n the length of each σ_n -orbit is relatively prime to $w_{n'}$ for all $n' > n$.

Proof. Note that $\sigma_{n+1} = \sigma_n^{w_{n+1}}$. Therefore, each $\sigma_{n'}$ -orbit is contained in some σ_n -orbit for $n' > n$ and if the length of a σ_n -orbit J is not relatively prime to $w_{n'}$ for some $n' > n$, then J splits into several $\sigma_{n'}$ -orbits. Since a permutation of $\{1, 2, \dots, r\}$ has at most r orbits, the claim follows. \square

Claim 5. Let n be sufficiently large as in Claim 3. If the length of a σ_n -orbit J is relatively prime to $w_{n'}$ for all $n' > n$, then $T_{\tilde{\psi}_n|_{\Gamma_n \times J}} \cong \Theta$.

Proof. Let $l = |J|$ be the length of J . It is clear that $T_{\tilde{\psi}_n|_{\Gamma_n \times J}} \cong T_{\psi_n^l}$. Note that Γ_n is the inverse limit of the sequence

$$\{\phi_{n+k} : \text{Ker}(\phi_{n+1} \cdots \phi_{n+k}) \rightarrow \text{Ker}(\phi_{n+1} \cdots \phi_{n+k-1})\}_{k \geq 1}.$$

Since l is relatively prime to $w_{n'}$ for all $n' > n$, the homomorphisms

$$\text{Ker}(\phi_{n+1} \cdots \phi_{n+k}) \rightarrow \text{Ker}(\phi_{n+1} \cdots \phi_{n+k})$$

defined by $x \mapsto x^l$ are isomorphic. It follows that the homomorphism $\Gamma_n \rightarrow \Gamma_n$ defined by $x \mapsto x^l$ is isomorphic, via which one notices that ψ_n is topologically conjugate to ψ_n^l . So we have $T_{\psi_n} \cong T_{\psi_n^l}$ and therefore, $T_{\tilde{\psi}_n|_{\Gamma_n \times J}} \cong T_{\psi_n^l} \cong T_{\psi_n} \cong \Theta$. \square

From Claim 4 and Claim 5 we also have

Corollary 5 ([Gu]). *A solenoid of type $\varpi = (w_1, w_2, \dots, w_n, \dots)$ has a connected r -fold covering if and only if r is relatively prime to all but finitely many w_n .*

ACKNOWLEDGEMENTS

The present form of Theorem 1 is influenced by the referee's suggestion to show no embedding of a (non-torus) m -dimensional connected compact Abelian topological group (cf. [KW]) into \mathbb{R}^{m+1} .

REFERENCES

- [B1] Bing, R. H. *A simple closed curve is the only homogeneous bounded plane continuum that contains an arc*, *Canad. J. Math.* **12** (1960), 209–230. MR0111001 (22:1869)
- [B2] Bing, R. H. *Embedding circle-like continua in the plane*, *Canad. J. Math.* **14** (1962), 113–128. MR0131865 (24:A1712)
- [ES] Eilenberg, S.; Steenrod, N. *Foundations of algebraic topology*, Princeton University Press, Princeton, New Jersey, 1952. MR0050886 (14:398b)
- [GH] Greenberg, M. J.; Harper, J. R. *Algebraic topology. A first course*, Mathematics Lecture Note Series 58, Benjamin/Cummings Publishing Co., Inc., Reading, MA, 1981. MR643101 (83b:55001)
- [Gu] Gumerov, R. N. *On finite-sheeted covering mappings onto solenoids*, *Proc. Amer. Math. Soc.* **133** (2005), 2771–2778. MR2146226 (2006d:54024)
- [KW] Keesling, J.; Wilson, D. *Embedding T^n -like continua in Euclidean space*. *Topology Appl.* **21** (1985), no. 3, 241–249. MR812642 (87h:54033)
- [Mc] McCord, M. C. *Inverse limit sequences with covering maps*, *Trans. Amer. Math. Soc.* **114** (1965), no. 1, 197–209. MR0173237 (30:3450)
- [Pr] Prajs, J. *Homogeneous continua in Euclidean $(n + 1)$ -space which contain an n -cube are n -manifolds*. *Trans. Amer. Math. Soc.* **318** (1990), no. 1, 143–148. MR943307 (90f:54055)

DEPARTMENT OF MATHEMATICS, PEKING UNIVERSITY, BEIJING 100871, PEOPLE'S REPUBLIC OF CHINA

E-mail address: bjjiang@math.pku.edu.cn

DEPARTMENT OF MATHEMATICS, PEKING UNIVERSITY, BEIJING 100871, PEOPLE'S REPUBLIC OF CHINA

E-mail address: wangsc@math.pku.edu.cn

DEPARTMENT OF MATHEMATICS, ZHONGSHAN UNIVERSITY, GUANGZHOU 510275, PEOPLE'S REPUBLIC OF CHINA

E-mail address: zhenghao@sysu.edu.cn