

## TENSOR PRODUCTS OF LEAVITT PATH ALGEBRAS

PERE ARA AND GUILLERMO CORTIÑAS

(Communicated by Birge Huisgen-Zimmermann)

ABSTRACT. We compute the Hochschild homology of Leavitt path algebras over a field  $k$ . As an application, we show that  $L_2$  and  $L_2 \otimes L_2$  have different Hochschild homologies, and so they are not Morita equivalent; in particular, they are not isomorphic. Similarly,  $L_\infty$  and  $L_\infty \otimes L_\infty$  are distinguished by their Hochschild homologies, and so they are not Morita equivalent either. By contrast, we show that  $K$ -theory cannot distinguish these algebras; we have  $K_*(L_2) = K_*(L_2 \otimes L_2) = 0$  and  $K_*(L_\infty) = K_*(L_\infty \otimes L_\infty) = K_*(k)$ .

### 1. INTRODUCTION

Elliott's theorem [21] states that  $\mathcal{O}_2 \otimes \mathcal{O}_2 \cong \mathcal{O}_2$  plays an important role in the proof of the celebrated classification theorem of Kirchberg algebras in the UCT class, due to Kirchberg [14] and Phillips [19]. Recall that a Kirchberg algebra is a purely infinite, simple, nuclear and separable  $C^*$ -algebra. The Kirchberg-Phillips theorem states that this class of simple  $C^*$ -algebras is completely classified by its topological  $K$ -theory. The analogous question whether the algebras  $L_2$  and  $L_2 \otimes L_2$  are isomorphic has remained open for some time. Here  $L_2$  is the Leavitt algebra of type  $(1, 2)$  over a field  $k$  (see [17]), that is, the  $k$ -algebra with generators  $x_1, x_2, x_1^*, x_2^*$  and relations given by  $x_i^* x_j = \delta_{i,j}$  and  $\sum_{i=1}^2 x_i x_i^* = 1$ .

In this paper we obtain a negative answer to this question. Indeed, we analyze a much larger class of algebras, namely the tensor products of Leavitt path algebras of finite quivers in terms of their Hochschild homology, and we prove that, for  $1 \leq n < m \leq \infty$ , the tensor products  $E = \bigotimes_{i=1}^n L(E_i)$  and  $F = \bigotimes_{j=1}^m L(F_j)$  of Leavitt path algebras of non-acyclic finite quivers  $E_i, F_j$  are distinguished by their Hochschild homologies (Theorem 5.1). Because Hochschild homology is Morita invariant, we conclude that  $E$  and  $F$  are not Morita equivalent for  $n < m$ . Since  $L_2$  is the Leavitt path algebra of the graph with one vertex and two arrows, we obtain that  $L_2 \otimes L_2$  and  $L_2$  are not Morita equivalent; in particular, they are not isomorphic.

Recall that, by a theorem of Kirchberg [15], a simple, nuclear and separable  $C^*$ -algebra  $A$  is purely infinite if and only if  $A \otimes \mathcal{O}_\infty \cong A$ . We also show that the analogue of Kirchberg's result is not true for Leavitt algebras. We prove in

---

Received by the editors August 1, 2011 and, in revised form, November 9, 2011.

2010 *Mathematics Subject Classification*. Primary 16E40, 16S99; Secondary 19D50.

The first author was partially supported by DGI MICIIN-FEDER MTM2008-06201-C02-01 and by the Comissionat per Universitats i Recerca de la Generalitat de Catalunya.

The second author was supported by CONICET and partially supported by grants PIP 112-200801-00900, UBACyTs X051 and 20020100100386, and MTM2007-64074.

©2013 American Mathematical Society  
Reverts to public domain 28 years from publication

Proposition 5.3 that if  $E$  is a non-acyclic quiver, then  $L_\infty \otimes L(E)$  and  $L(E)$  are not Morita equivalent, and also that  $L_\infty \otimes L_\infty$  and  $L_\infty$  are not Morita equivalent.

Using the results in [5] we prove that the algebras  $L_2$  and  $L_2 \otimes L(F)$ , for  $F$  an arbitrary finite quiver, have trivial  $K$ -theory: all algebraic  $K$ -theory groups  $K_i$ ,  $i \in \mathbb{Z}$ , vanish on them (this follows from Lemma 6.1 and Proposition 6.2). We also compute  $K_*(L(F)) = K_*(L_\infty \otimes L(F))$  and that  $K_*(L_\infty) = K_*(L_\infty \otimes L_\infty) = K_*(k)$  is the  $K$ -theory of the ground field (see Proposition 6.3 and Corollary 6.4). This implies in particular that, in contrast with the analytic situation, no classification result, in terms solely of  $K$ -theory, can be expected for a class of central, simple  $k$ -algebras, containing all purely infinite simple unital Leavitt path algebras and closed under tensor products. It is worth mentioning that an important step towards a  $K$ -theoretic classification of purely infinite simple Leavitt path algebras of finite quivers has been achieved in [2].

We refer the reader to [3], [7] and [20] for the basics on Leavitt algebras, Leavitt path algebras and graph  $C^*$ -algebras, and to [22] for a nice survey on the Kirchberg-Phillips Theorem.

*Notation.* We fix a field  $k$ ; all vector spaces, tensor products and algebras are over  $k$ . If  $R$  and  $S$  are unital  $k$ -algebras, then by an  $(R, S)$ -bimodule we understand a left module over  $R \otimes S^{op}$ . By an  $R$ -bimodule we shall mean an  $(R, R)$  bimodule, that is, a left module over the enveloping algebra  $R^e = R \otimes R^{op}$ . Hochschild homology of  $k$ -algebras is always taken over  $k$ . If  $M$  is an  $R$ -bimodule, we write

$$HH_n(R, M) = \text{Tor}_n^{R^e}(R, M)$$

for the Hochschild homology of  $R$  with coefficients in  $M$  and we abbreviate  $HH_n(R) = HH_n(R, R)$ .

## 2. HOCHSCHILD HOMOLOGY

Let  $k$  be a field,  $R$  a  $k$ -algebra and  $M$  an  $R$ -bimodule. The Hochschild homology  $HH_*(R, M)$  of  $R$  with coefficients in  $M$  was defined in the introduction. It is computed by the *Hochschild complex*  $HH(R, M)$ , which is given in degree  $n$  by

$$HH(R, M)_n = M \otimes R^{\otimes n}.$$

It is equipped with the Hochschild boundary map  $b$  defined by

$$b(a_0 \otimes a_1 \otimes \cdots \otimes a_n) = \sum_{i=0}^{n-1} (-1)^i a_0 \otimes \cdots \otimes a_i a_{i+1} \otimes \cdots \otimes a_n + (-1)^n a_n a_0 \otimes \cdots \otimes a_{n-1}.$$

If  $R$  and  $M$  happen to be  $\mathbb{Z}$ -graded, then  $HH(R, M)$  splits into a direct sum of subcomplexes

$$HH(R, M) = \bigoplus_{m \in \mathbb{Z}} {}_m HH(R, M).$$

The homogeneous component of degree  $m$  of  $HH(R, M)_n$  is the linear subspace of  $HH(R, M)_n$  generated by all elementary tensors  $a_0 \otimes \cdots \otimes a_n$  with  $a_i$  homogeneous and  $\sum_i |a_i| = m$ . One of the first basic properties of the Hochschild complex is that it commutes with filtering colimits. Thus we have

**Lemma 2.1.** *Let  $I$  be a filtered ordered set and let  $\{(R_i, M_i) : i \in I\}$  be a directed system of pairs  $(R_i, M_i)$  consisting of an algebra  $R_i$  and an  $R_i$ -bimodule  $M_i$ , with algebra maps  $R_i \rightarrow R_j$  and  $R_i$ -bimodule maps  $M_i \rightarrow M_j$  for each  $i \leq j$ . Let  $(R, M) = \text{colim}_i (R_i, M_i)$ . Then  $HH_n(R, M) = \text{colim}_i HH_n(R_i, M_i)$  ( $n \geq 0$ ).*

Let  $R_i$  be a  $k$ -algebra and  $M_i$  an  $R_i$ -bimodule ( $i = 1, 2$ ). The Künneth formula establishes a natural isomorphism ([23, 9.4.1])

$$HH_n(R_1 \otimes R_2, M_1 \otimes M_2) \cong \bigoplus_{p=0}^n HH_p(R_1, M_1) \otimes HH_{n-p}(R_2, M_2).$$

Another fundamental fact about Hochschild homology which we shall need is Morita invariance. Let  $R$  and  $S$  be Morita equivalent algebras, and let  $P \in R \otimes S^{op}$ -mod and  $Q \in S \otimes R^{op}$ -mod implement the Morita equivalence. Then ([23, Thm. 9.5.6])

$$(2.2) \quad HH_n(R, M) = HH_n(S, Q \otimes_R M \otimes_R P).$$

**Lemma 2.3.** *Let  $R_1, \dots, R_n$  and  $S_1, \dots, S_m, \dots$  be a finite and an infinite sequence of algebras, and let  $R = \bigotimes_{i=1}^n R_i$ ,  $S_{\leq m} = \bigotimes_{j=1}^m S_j$  and  $S = \bigotimes_{j=1}^\infty S_j$ . Assume:*

- (1)  $HH_q(R_i) \neq 0 \neq HH_q(S_j) \quad (q = 0, 1), (1 \leq i \leq n), (1 \leq j)$ .
- (2)  $HH_p(R_i) = HH_p(S_j) = 0$  for  $p \geq 2, 1 \leq i \leq n, 1 \leq j$ .
- (3)  $n \neq m$ .

Then no two of  $R, S_{\leq m}$  and  $S$  are Morita equivalent.

*Proof.* By the Künneth formula, we have

$$HH_n(R) = \bigotimes_{i=1}^n HH_1(R_i) \neq 0, \quad HH_p(R) = 0, \quad p > n.$$

By the same argument,  $HH_p(S_{\leq m})$  is non-zero for  $p = m$  and zero for  $p > m$ . Hence if  $n \neq m$ ,  $R$  and  $S_{\leq m}$  do not have the same Hochschild homology, and therefore they cannot be Morita equivalent, by (2.2). Similarly, by Lemma 2.1, we have

$$HH_n(S) = \bigoplus_{J \subset \mathbb{N}, |J|=n} \left( \bigotimes_{j \in J} HH_1(S_j) \right) \otimes \left( \bigotimes_{j \notin J} HH_0(S_j) \right)$$

so that  $HH_n(S)$  is non-zero for all  $n \geq 1$ , and thus it cannot be Morita equivalent to either  $R$  or  $S_{\leq m}$ . □

### 3. HOCHSCHILD HOMOLOGY OF CROSSED PRODUCTS

Let  $R$  be a unital algebra and  $G$  a group acting on  $R$  by algebra automorphisms. Form the crossed-product algebra  $S = R \rtimes G$ , and consider the Hochschild complex  $HH(S)$ . For each conjugacy class  $\xi$  of  $G$ , the graded submodule  $HH^\xi(S) \subset HH(S)$  generated in degree  $n$  by the elementary tensors  $a_0 \rtimes g_0 \otimes \dots \otimes a_n \rtimes g_n$  with  $g_0 \dots g_n \in \xi$  is a subcomplex, and we have a direct sum decomposition  $HH(S) = \bigoplus_\xi HH^\xi(S)$ . The following theorem of Lorenz describes the complex  $HH^\xi(S)$  corresponding to the conjugacy class  $\xi = [g]$  of an element  $g \in G$  as hyperhomology over the centralizer subgroup  $Z_g \subset G$ .

**Theorem 3.1** ([16]). *Let  $R$  be a unital  $k$ -algebra,  $G$  a group acting on  $R$  by automorphisms,  $g \in G$  and  $Z_g \subset G$  the centralizer subgroup. Let  $S = R \rtimes G$  be the crossed product algebra and  $HH^{(g)}(S) \subset HH(S)$  be the subcomplex described above. Consider the  $R$ -submodule  $S_g = R \rtimes g \subset S$ . Then there is a quasi-isomorphism*

$$HH^{[g]}(S) \xrightarrow{\sim} \mathbb{H}(Z_g, HH(R, S_g)).$$

In particular, we have a spectral sequence

$$E_{p,q}^2 = H_p(Z_g, HH_q(R, S_g)) \Rightarrow HH_{p+q}^{[g]}(S).$$

*Remark 3.2.* Lorenz formulates his result in terms of the spectral sequence alone, but his proof shows that there is a quasi-isomorphism as stated above. An explicit formula is given for example in the proof of [11, Lemma 7.2].

Let  $A$  be a not necessarily unital  $k$ -algebra and write  $\tilde{A}$  for its unitalization. Recall from [24] that  $A$  is called  $H$ -unital if the groups  $\text{Tor}_n^{\tilde{A}}(k, A)$  vanish for all  $n \geq 0$ . Wodzicki proved in [24] that  $A$  is  $H$ -unital if and only if for every embedding  $A \triangleleft R$  of  $A$  as a two-sided ideal of a unital ring  $R$ , the map

$$HH(A) \rightarrow HH(R : A) = \ker(HH(R) \rightarrow HH(R/A))$$

is a quasi-isomorphism.

**Lemma 3.3.** *Theorem 3.1 still holds if the condition that  $R$  be unital is replaced by the condition that it be  $H$ -unital.*

*Proof.* Follows from Theorem 3.1 and the fact, proved in [11, Prop. A.6.5], that  $R \rtimes G$  is  $H$ -unital if  $R$  is as well.  $\square$

Let  $R$  be a unital algebra and  $\phi : R \rightarrow pRp$  a corner isomorphism. As in [6], we consider the skew Laurent polynomial algebra  $R[t_+, t_-, \phi]$ . This is the  $R$ -algebra generated by elements  $t_+$  and  $t_-$  subject to the following relations:

$$\begin{aligned} t_+a &= \phi(a)t_+ \\ at_- &= t_-\phi(a) \\ t_-t_+ &= 1 \\ t_+t_- &= p \end{aligned}$$

Observe that the algebra  $S = R[t_+, t_-, \phi]$  is  $\mathbb{Z}$ -graded by  $\deg(r) = 0, \deg(t_{\pm}) = \pm 1$ . The homogeneous component of degree  $n$  is given by

$$R[t_+, t_-, \phi]_n = \begin{cases} t_-^{-n}R & n < 0 \\ R & n = 0 \\ Rt_+^n & n > 0 \end{cases}$$

**Proposition 3.4.** *Let  $R$  be a unital ring,  $\phi : R \rightarrow pRp$  a corner isomorphism, and  $S = R[t_+, t_-, \phi]$ . Consider the weight decomposition  $HH(S) = \bigoplus_{m \in \mathbb{Z}} {}^m HH(S)$ . There is a quasi-isomorphism*

$$(3.5) \quad {}^m HH(S) \xrightarrow{\sim} \text{Cone}(1 - \phi : HH(R, S_m) \rightarrow HH(R, S_m)).$$

*Proof.* If  $\phi$  is an automorphism, then  $S = R \rtimes_{\phi} \mathbb{Z}$ , the right hand side of (3.5) computes  $\mathbb{H}(\mathbb{Z}, HH(R, S_m))$ , and the proposition becomes the particular case  $G = \mathbb{Z}$  of Theorem 3.1. In the general case, let  $A$  be the colimit of the inductive system

$$R \xrightarrow{\phi} R \xrightarrow{\phi} R \xrightarrow{\phi} \dots$$

Note that  $\phi$  induces an automorphism  $\hat{\phi} : A \rightarrow A$ . Now  $A$  is  $H$ -unital, since it is a filtering colimit of unital algebras, and thus the assertion of the proposition is true for the pair  $(A, \hat{\phi})$ , by Lemma 3.3. Hence it suffices to show that for  $B = A \rtimes_{\hat{\phi}} \mathbb{Z}$  the maps  $HH(S) \rightarrow HH(B)$  and  $\text{Cone}(1 - \phi : HH(R, S_m) \rightarrow HH(R, S_m)) \rightarrow \text{Cone}(1 - \phi : HH(A, B_m) \rightarrow HH(A, B_m))$  ( $m \in \mathbb{Z}$ ) are quasi-isomorphisms. The analogous property for  $K$ -theory is shown in the course of the third step of the proof of [5, Thm. 3.6]. Since the proof in [5] uses only that  $K$ -theory commutes with filtering colimits and is matrix invariant on those rings for which it satisfies excision, it applies verbatim to Hochschild homology. This concludes the proof.  $\square$

4. HOCHSCHILD HOMOLOGY OF THE LEAVITT PATH ALGEBRA

Let  $E = (E_0, E_1, r, s)$  be a finite quiver and let  $\hat{E} = (E_0, E_1 \sqcup E_1^*, r, s)$  be the double of  $E$ , which is the quiver obtained from  $E$  by adding an arrow  $\alpha^*$  for each arrow  $\alpha \in E_1$ , going in the opposite direction. The *Leavitt path algebra* of  $E$  is the algebra  $L(E)$  with one generator for each arrow  $\alpha \in \hat{E}_1$  and one generator  $p_i$  for each vertex  $i \in E_0$ , subject to the following relations:

$$\begin{aligned} p_i p_j &= \delta_{i,j} p_i && (i, j \in E_0) \\ p_{s(\alpha)} \alpha &= \alpha = \alpha p_{r(\alpha)} && (\alpha \in \hat{E}_1) \\ \alpha^* \beta &= \delta_{\alpha, \beta} p_{r(\alpha)} && (\alpha, \beta \in E_1) \\ p_i &= \sum_{\alpha \in E_1, s(\alpha)=i} \alpha \alpha^* && (i \in E_0 \setminus \text{Sink}(E)) \end{aligned}$$

The algebra  $L = L(E)$  is equipped with a  $\mathbb{Z}$ -grading. The grading is determined by  $|\alpha| = 1, |\alpha^*| = -1$ , for  $\alpha \in E_1$ . Let  $L_{0,n}$  be the linear span of all elements of the form  $\gamma \nu^*$ , where  $\gamma$  and  $\nu$  are paths with  $r(\gamma) = r(\nu)$  and  $|\gamma| = |\nu| = n$ . By [7, proof of Theorem 5.3], we have  $L_0 = \bigcup_{n=0}^\infty L_{0,n}$ . For each  $i$  in  $E_0$  and each  $n \in \mathbb{Z}^+$ , let us denote by  $P(n, i)$  the set of paths  $\gamma$  in  $E$  such that  $|\gamma| = n$  and  $r(\gamma) = i$ . The algebra  $L_{0,0}$  is isomorphic to  $\prod_{i \in E_0} k$ . In general, the algebra  $L_{0,n}$  is isomorphic to

$$(4.1) \quad \left[ \prod_{m=0}^{n-1} \left( \prod_{i \in \text{Sink}(E)} M_{|P(m,i)|}(k) \right) \right] \times \left[ \prod_{i \in E_0} M_{|P(n,i)|}(k) \right].$$

The transition homomorphism  $L_{0,n} \rightarrow L_{0,n+1}$  is the identity on the factors

$$\prod_{i \in \text{Sink}(E)} M_{|P(m,i)|}(k),$$

for  $0 \leq m \leq n - 1$ , and also on the factor

$$\prod_{i \in \text{Sink}(E)} M_{|P(n,i)|}(k)$$

of the last term of the displayed formula. The transition homomorphism

$$\prod_{i \in E_0 \setminus \text{Sink}(E)} M_{|P(n,i)|}(k) \rightarrow \prod_{i \in E_0} M_{|P(n+1,i)|}(k)$$

is a block diagonal map induced by the following identification in  $L(E)_0$ : A matrix unit in a factor  $M_{|P(n,i)|}(k)$ , where  $i \in E_0 \setminus \text{Sink}(E)$ , is a monomial of the form  $\gamma \nu^*$ , where  $\gamma$  and  $\nu$  are paths of length  $n$  with  $r(\gamma) = r(\nu) = i$ . Since  $i$  is not a sink, we can enlarge the paths  $\gamma$  and  $\nu$  using the edges that  $i$  emits, obtaining paths of length  $n + 1$ , and the last relation in the definition of  $L(E)$  gives

$$\gamma \nu^* = \sum_{\{\alpha \in E_1 \mid s(\alpha)=i\}} (\gamma \alpha)(\nu \alpha)^*.$$

Assume  $E$  has no sources. For each  $i \in E_0$ , choose an arrow  $\alpha_i$  such that  $r(\alpha_i) = i$ . Consider the elements

$$t_+ = \sum_{i \in E_0} \alpha_i, \quad t_- = t_+^*.$$

One checks that  $t_-t_+ = 1$ . Thus, since  $|t_{\pm}| = \pm 1$ , the endomorphism

$$(4.2) \quad \phi : L \rightarrow L, \quad \phi(x) = t_+xt_-$$

is homogeneous of degree 0 with respect to the  $\mathbb{Z}$ -grading. In particular, it restricts to an endomorphism of  $L_0$ . By [6, Lemma 2.4], we have

$$(4.3) \quad L = L_0[t_+, t_-, \phi].$$

Consider the matrix  $N'_E = [n_{i,j}] \in M_{e_0}\mathbb{Z}$  given by

$$n_{i,j} = \#\{\alpha \in E_1 : s(\alpha) = i, \quad r(\alpha) = j\}.$$

Let  $e'_0 = |\text{Sink}(E)|$ . We assume that  $E_0$  is ordered so that the first  $e'_0$  elements of  $E_0$  correspond to its sinks. Accordingly, the first  $e'_0$  rows of the matrix  $N'_E$  are 0. Let  $N_E$  be the matrix obtained by deleting these  $e'_0$  rows. The matrix that enters the computation of the Hochschild homology of the Leavitt path algebra is

$$\begin{pmatrix} 0 \\ 1_{e_0-e'_0} \end{pmatrix} - N_E^t : \mathbb{Z}^{e_0-e'_0} \longrightarrow \mathbb{Z}^{e_0}.$$

By a slight abuse of notation, we will write  $1 - N_E^t$  for this matrix. Note that  $1 - N_E^t \in M_{e_0 \times (e_0 - e'_0)}(\mathbb{Z})$ . Of course,  $N_E = N'_E$  in case  $E$  has no sinks.

**Theorem 4.4.** *Let  $E$  be a finite quiver without sources, and let  $N = N_E$ . For each  $i \in E_0 \setminus \text{Sink}(E)$  and  $m \geq 1$ , let  $V_{i,m}$  be the vector space generated by all closed paths  $c$  of length  $m$  with  $s(c) = r(c) = i$ . Let  $\mathbb{Z} = \langle \sigma \rangle$  act on*

$$V_m = \bigoplus_{i \in E_0 \setminus \text{Sink}(E)} V_{i,m}$$

by rotation of closed paths. We have

$${}_mHH_n(L(E)) = \begin{cases} \text{coker}(1 - \sigma : V_{|m|} \rightarrow V_{|m|}) & n = 0, m \neq 0 \\ \text{coker}(1 - N^t) & n = m = 0 \\ \text{ker}(1 - \sigma : V_{|m|} \rightarrow V_{|m|}) & n = 1, m \neq 0 \\ \text{ker}(1 - N^t) & n = 1, m = 0 \\ 0 & n \notin \{0, 1\} \end{cases}$$

*Proof.* Let  $L = L(E)$ ,  $P = P(E) \subset L$  be the path algebras of  $E$  and  $W_m \subset P$  be the subspace generated by all paths of length  $m$ . For each fixed  $n \geq 1$  and  $m \in \mathbb{Z}$ , consider the following  $L_{0,n}$ -bimodule:

$$L_{m,n} = \begin{cases} L_{0,n}W_mL_{0,n} & m > 0 \\ L_{0,n}W_{-m}^*L_{0,n} & m < 0 \end{cases}$$

Write  $L = L(E)$ , and let  ${}_mL$  be the homogeneous part of degree  $m$ ; we have

$${}_mL = \bigcup_{n \geq 1} L_{m,n}.$$

If  $m$  is positive, then there is a basis of  $L_{m,n}$  consisting of the products  $\alpha\theta\beta^*$  where each of  $\alpha$ ,  $\beta$  and  $\theta$  is a path in  $E$ ,  $r(\alpha) = s(\theta)$ ,  $r(\beta) = r(\theta)$ ,  $|\alpha| = |\beta| = n$  and  $|\theta| = m$ . Hence the formula

$$\pi(\alpha\theta\beta^*) = \begin{cases} \theta & \text{if } \alpha = \beta \\ 0 & \text{else} \end{cases}$$

defines a surjective linear map  $L_{m,n} \rightarrow V_m$ . One checks that  $\pi$  induces an isomorphism

$$HH_0(L_{0,n}, L_{m,n}) \cong V_m \quad (m > 0).$$

Similarly, if  $m < 0$ , then

$$HH_0(L_{0,n}, L_{m,n}) = V_{|m|}^* \cong V_{-m}.$$

Next, by (4.1), we have

$$HH_0(L_{0,n}) = k[E \setminus \text{Sink}(E)] \oplus \bigoplus_{i \in \text{Sink}(E)} k^{r(i,n)}.$$

Here

$$r(i, n) = \max\{r \leq n : P(r, i) \neq \emptyset\}.$$

Now note that because  $L_{0,n}$  is a product of matrix algebras, it is separable, and thus  $HH_1(L_{0,n}, M) = 0$  for any bimodule  $M$ . As observed in (4.3), for the automorphism (4.2), we have  $L = L_0[t_+, t_-, \phi]$ . Hence in view of Proposition 3.4 and Lemma 2.1, it only remains to identify the maps  $HH_0(L_{0,n}, L_{m,n}) \rightarrow HH_0(L_{0,n+1}, L_{m,n+1})$  induced by inclusion and by the homomorphism  $\phi$ . One checks that for  $m \neq 0$ , these are respectively the cyclic permutation and the identity  $V_{|m|} \rightarrow V_{|m|}$ . The case  $m = 0$  is dealt with in the same way as in [5, Proof of Theorem 5.10].  $\square$

**Corollary 4.5.** *Let  $E$  be a finite quiver with at least one non-trivial closed path.*

- i)  $HH_n(L(E)) = 0$  for  $n \notin \{0, 1\}$ .
- ii)  ${}_m HH_*(L(E)) \cong {}_{-m} HH_*(L(E))$  ( $m \in \mathbb{Z}$ ).
- iii) *There exist  $m > 0$  such that  ${}_m HH_0(L(E))$  and  ${}_m HH_1(L(E))$  are both non-zero.*

*Proof.* We first reduce to the case where the graph does not have sources. By the proof of [5, Theorem 6.3], there is a finite complete subgraph  $F$  of  $E$  such that  $F$  has no sources,  $F$  contains all the non-trivial closed paths of  $E$ ,  $\text{Sink}(F) = \text{Sink}(E)$ , and  $L(F)$  is a full corner in  $L(E)$  with respect to the homogeneous idempotent  $\sum_{v \in F^0} p_v$ . It follows that  $HH_*(L(E))$  and  $HH_*(L(F))$  are graded-isomorphic. Therefore we can assume that  $E$  has no sources.

The first two assertions are already part of Theorem 4.4. For the last assertion, let  $\alpha$  be a primitive closed path in  $E$ , and let  $m = |\alpha|$ . Let  $\sigma$  be the cyclic permutation; then  $\{\sigma^i \alpha : i = 0, \dots, m - 1\}$  is a linearly independent set. Hence  $N(\alpha) = \sum_{i=0}^{m-1} \sigma^i \alpha$  is a non-zero element of  $V_m^\sigma = {}_m HH_1(L(E))$ . Since on the other hand  $N$  vanishes on the image of  $1 - \sigma : V_m \rightarrow V_m$ , it also follows that the class of  $\alpha$  in  ${}_m HH_0(L(E))$  is non-zero.  $\square$

### 5. APPLICATIONS

**Theorem 5.1.** *Let  $E_1, \dots, E_n$  and  $F_1, \dots, F_m$  be finite quivers. Assume that  $n \neq m$  and that each of the  $E_i$  and the  $F_j$  has at least one non-trivial closed path. Then the algebras  $L(E_1) \otimes \dots \otimes L(E_n)$  and  $L(F_1) \otimes \dots \otimes L(F_m)$  are not Morita equivalent.*

*Proof.* Immediate from Lemma 2.3 and Corollary 4.5(iii).  $\square$

**Example 5.2.** It follows from Theorem 5.1 that  $L_2$  and  $L_2 \otimes_k L_2$  are not Morita equivalent. There is another way of proving this, due to Jason Bell and George Bergman [8]. By Theorem 3.3 of [9],  $\text{l.gl.dim } L_2 \leq 1$ . Using a module-theoretic construction, Bell and Bergman show that  $\text{l.gl.dim}(L_2 \otimes_k L_2) \geq 2$ , which forces  $L_2$  and  $L_2 \otimes_k L_2$  to not be Morita equivalent. Bergman then asked Warren Dicks whether general results were known about global dimensions of tensor products and was pointed to Proposition 10(2) of [12], which is an immediate consequence of

Theorem XI.3.1 of [10] and says that if  $k$  is a field and  $R$  and  $S$  are  $k$ -algebras, then  $\text{l.gl.dim } R + \text{w.gl.dim } S \leq \text{l.gl.dim}(R \otimes_k S)$ . Consequently, if  $\text{l.gl.dim } R < \infty$  and  $\text{w.gl.dim } S > 0$ , then  $\text{l.gl.dim } R < \text{l.gl.dim}(R \otimes_k S)$ ; in particular,  $R$  and  $R \otimes_k S$  are then not Morita equivalent. To see that  $\text{w.gl.dim } L_2 > 0$ , write  $x_1, x_2, x_1^*, x_2^*$  for the usual generators of  $L_2$  and use normal-form arguments to show that  $\{a \in L_2 \mid ax_1 = a + 1\} = \emptyset$  and  $\{b \in L_2 \mid x_1b = b\} = \{0\}$ . Hence, in  $L_2$ ,  $x_1 - 1$  does not have a left inverse and is not a left zerodivisor (or see [4]); thus,  $\text{w.gl.dim } L_2 > 0$ .

We denote by  $L_\infty$  the unital algebra presented by generators  $x_1, x_1^*, x_2, x_2^*, \dots$  and relations  $x_i^*x_j = \delta_{i,j}1$ .

**Proposition 5.3.** *Let  $E$  be any finite quiver having at least one non-trivial closed path. Then  $L_\infty \otimes L(E)$  and  $L(E)$  are not Morita equivalent. Similarly,  $L_\infty \otimes L_\infty$  and  $L_\infty$  are not Morita equivalent.*

*Proof.* Let  $C_n$  be the algebra presented by generators  $x_1, x_1^*, \dots, x_n, x_n^*$  and relations  $x_i^*x_j = \delta_{i,j}1$ , for  $1 \leq i, j \leq n$ . Then

$$(5.4) \quad L_\infty = \varinjlim C_n$$

and  $C_n \cong L(E_n)$ , where  $E_n$  is the graph having two vertices  $v, w$  and  $2n$  arrows  $e_1, \dots, e_n, f_1, \dots, f_n$ , with  $s(e_i) = r(e_i) = v = s(f_i)$  and  $r(f_i) = w$  for  $1 \leq i \leq n$ . (The isomorphism  $C_n \rightarrow L(E_n)$  is obtained by sending  $x_i$  to  $e_i + f_i$  and  $x_i^*$  to  $e_i^* + f_i^*$ .) It follows from Theorem 4.4 and (5.4) that the formulas in Theorem 4.4 for  ${}_mHH_n(L_\infty)$ ,  $m \neq 0$ , hold, taking as  $V_{i,m}$  the vector space generated by all the words in  $x_1, x_2, \dots$  of length  $m$ , and that  ${}_0HH_0(L_\infty) = k$  and  ${}_0HH_n(L_\infty) = 0$  for  $n \geq 1$ . As before, Lemma 2.3 gives the result.  $\square$

**Theorem 5.5.** *Let  $E_1, \dots, E_n$  and  $F_1, \dots, F_m, \dots$  be a finite and an infinite sequence of quivers. Assume that the number of indices  $i$  such that  $F_i$  has at least one non-trivial closed path is infinite. Then the algebras  $L(E_1) \otimes \dots \otimes L(E_n)$  and  $\bigotimes_{i=1}^\infty L(F_i)$  are not Morita equivalent.*

*Proof.* Immediate from Lemma 2.3 and Corollary 4.5(iii).  $\square$

**Example 5.6.** Let  $L^{(\infty)} = \bigotimes_{i=1}^\infty L_2$ , and let  $E$  be any quiver having at least one non-trivial closed path. Then  $L^{(\infty)} \otimes L(E)$  and  $L(E)$  are not Morita equivalent.

It would be interesting to know the answer to the following question:

**Question 5.7.** Is there a unital homomorphism  $\phi: L_2 \otimes L_2 \rightarrow L_2$ ?

Observe that to build a unital homomorphism  $\phi: L_2 \otimes L_2 \rightarrow L_2$ , it is enough to exhibit a *non-zero* homomorphism  $\psi: L_2 \otimes L_2 \rightarrow L_2$  because  $eL_2e \cong L_2$  for every non-zero idempotent  $e$  in  $L_2$ .

## 6. $K$ -THEORY

To conclude the paper we note that algebraic  $K$ -theory cannot distinguish between  $L_2$  and  $L_2 \otimes L_2$  or between  $L_\infty$  and  $L_\infty \otimes L_\infty$ . For this we need a lemma which might be of independent interest. Recall that a unital ring  $R$  is said to be *regular supercoherent* in case all the polynomial rings  $R[t_1, \dots, t_n]$  are regular coherent in the sense of [13].

**Lemma 6.1.** *Let  $E$  be a finite graph. Then  $L(E)$  is regular supercoherent.*

*Proof.* Let  $P(E)$  be the usual path algebra of  $E$ . It was observed in the proof of [3, Lemma 7.4] that the algebra  $P(E)[t]$  is regular coherent. The same proof gives that all the polynomial algebras  $P(E)[t_1, \dots, t_n]$  are regular coherent. This shows that  $P(E)$  is regular supercoherent. By [3, Proposition 4.1], the universal localization  $P(E) \rightarrow L(E) = \Sigma^{-1}P(E)$  is flat on the left. It follows that  $L(E)$  is left regular supercoherent (see [5, page 23]). Since  $L(E) \otimes k[t_1, \dots, t_n]$  admits an involution, it follows that  $L(E)$  is regular supercoherent.  $\square$

**Proposition 6.2.** *Let  $R$  be regular supercoherent. Then the algebraic  $K$ -theories of  $L_2$  and of  $L_2 \otimes R$  are both trivial.*

*Proof.* Let  $E$  be the quiver with one vertex and two arrows. Then  $L_2 \cong L(E)$ , and we have

$$L_2 \otimes R = L_R(E).$$

Applying [5, Theorem 7.6] we obtain that  $K_*(L_R(E)) = K_*(L(E)) = 0$ . The result follows.  $\square$

We finally obtain a  $K$ -absorbing result for Leavitt path algebras of finite graphs, indeed for any regular supercoherent algebra.

**Proposition 6.3.** *Let  $R$  be a regular supercoherent algebra. Then the natural inclusion  $R \rightarrow R \otimes L_\infty$  induces an isomorphism  $K_i(R) \rightarrow K_i(R \otimes L_\infty)$  for all  $i \in \mathbb{Z}$ .*

*Proof.* Adopting the notation used in the proof of Proposition 5.3, we see that it is enough to show that the natural map  $R \rightarrow R \otimes L(E_n)$  induces isomorphisms  $K_i(R) \rightarrow K_i(R \otimes L(E_n))$  for all  $i \in \mathbb{Z}$  and all  $n \geq 1$ . Since  $R$  is regular supercoherent, the  $K$ -theory of  $R \otimes L(E_n) \cong L_R(E_n)$  can be computed by using [5, Theorem 7.6]. By the explicit form of the quiver  $E_n$ , we thus obtain that

$$K_i(R \otimes L(E_n)) \cong (K_i(R) \oplus K_i(R))/(-n, 1 - n)K_i(R).$$

The natural map  $R \rightarrow L_R(E_n)$  factors as

$$R \rightarrow Rv \oplus Rw \rightarrow L_R(E_n).$$

The first map induces the diagonal homomorphism  $K_i(R) \rightarrow K_i(R) \oplus K_i(R)$ , sending  $x$  to  $(x, x)$ . The second map induces the natural surjection

$$K_i(R) \oplus K_i(R) \rightarrow (K_i(R) \oplus K_i(R))/(-n, 1 - n)K_i(R).$$

Therefore the natural homomorphism  $R \rightarrow L_R(E_n)$  induces an isomorphism

$$K_i(R) \xrightarrow{\sim} K_i(L_R(E_n)).$$

This concludes the proof.  $\square$

**Corollary 6.4.** *The natural maps  $k \rightarrow L_\infty \rightarrow L_\infty \otimes L_\infty$  induce  $K$ -theory isomorphisms  $K_*(k) = K_*(L_\infty) = K_*(L_\infty \otimes L_\infty)$ .*

*Proof.* A first application of Proposition 6.3 gives  $K_*(k) = K_*(L_\infty)$ . A second application shows that for  $E_n$  as in the proof above, the inclusion  $L(E_n) \rightarrow L(E_n) \otimes L_\infty$  induces a  $K$ -theory isomorphism; passing to the limit, we obtain the corollary.  $\square$

#### ACKNOWLEDGEMENT

Part of the research for this article was carried out during a visit of the second author to the Centre de Recerca Matemàtica. He is indebted to this institution for its hospitality.

#### REFERENCES

- [1] G. Abrams, G. Aranda Pino. *The Leavitt path algebra of a graph*. J. Algebra **293** (2005) 319–334. MR2172342 (2007b:46085)
- [2] G. Abrams, A. Louly, E. Pardo, C. Smith. *Flow invariants in the classification of Leavitt path algebras*. J. Algebra **333** (2011) 202–231. MR2785945
- [3] P. Ara, M. Brustenga. *Module theory over Leavitt path algebras and  $K$ -theory*. J. Pure Appl. Algebra **214** (2010) 1131–1151. MR2586992 (2011b:16109)
- [4] P. Ara, M. Brustenga. *The regular algebra of a quiver*. J. Algebra **309** (2007) 207–235. MR2301238 (2008a:16019)
- [5] P. Ara, M. Brustenga, G. Cortiñas.  *$K$ -theory of Leavitt path algebras*. Münster Journal of Mathematics **2** (2009) 5–34. MR2545605 (2011d:46145)
- [6] P. Ara, M.A. González-Barroso, K.R. Goodearl, E. Pardo. *Fractional skew monoid rings*. J. Algebra **278** (2004) 104–126. MR2068068 (2005f:16042)
- [7] P. Ara, M. A. Moreno, E. Pardo, *Nonstable  $K$ -theory for graph algebras*. Algebr. Represent. Theory **10** (2007) 157–178. MR2310414 (2008b:46094)
- [8] J. Bell, G. Bergman. Private communication, 2011.
- [9] G. M. Bergman and Warren Dicks, *Universal derivations and universal ring constructions*. Pacific J. Math. **79** (1978) 293–337. MR531320 (81b:16024)
- [10] H. Cartan, S. Eilenberg, *Homological Algebra*. Princeton University Press, Princeton, N. J., 1956. MR0077480 (17:1040e)
- [11] G. Cortiñas, E. Ellis. *Isomorphism conjectures with proper coefficients*. arXiv:1108.5196v3.
- [12] S. Eilenberg, A. Rosenberg, D. Zelinsky, *On the dimension of modules and algebras, VIII. Dimension of tensor products*. Nagoya Math. J. **12** (1957) 71–93. MR0098774 (20:5229)
- [13] S. M. Gersten.  *$K$ -theory of free rings*. Comm. Algebra **1** (1974) 39–64. MR0396671 (53:533)
- [14] E. Kirchberg, *The classification of purely infinite  $C^*$ -algebras using Kasparov theory*. Preprint.
- [15] E. Kirchberg, N.C. Phillips, *Embedding of exact  $C^*$ -algebras in the Cuntz algebra  $\mathcal{O}_2$* . J. Reine Angew. Math. **525** (2000) 17–53. MR1780426 (2001d:46086a)
- [16] M. Lorenz. *On the homology of graded algebras*. Comm. Algebra **20** (1992) 489–507. MR1146311 (93b:19003)
- [17] W. G. Leavitt. *The module type of a ring*. Trans. Amer. Math. Soc. **103** (1962) 113–130. MR0132764 (24:A2600)
- [18] J. L. Loday, *Cyclic homology*. Grund. Math. Wiss., 301. Springer-Verlag, Berlin, Heidelberg, 1998. MR1600246 (98h:16014)
- [19] N. C. Phillips, *A classification theorem for nuclear purely infinite simple  $C^*$ -algebras*. Doc. Math. **5** (2000) 49–114. MR1745197 (2001d:46086b)
- [20] I. Raeburn. *Graph algebras*. CBMS Regional Conference Series in Mathematics, 103. Published for the Conference Board of the Mathematical Sciences, Washington, DC, by the American Mathematical Society, Providence, RI, 2005. MR2135030 (2005k:46141)
- [21] M. Rørdam, *A short proof of Elliott’s theorem*. C. R. Math. Rep. Acad. Sci. Canada **16** (1994) 31–36. MR1276341 (95d:46064)
- [22] M. Rørdam, *Classification of nuclear, simple  $C^*$ -algebras*. Classification of nuclear  $C^*$ -algebras. Entropy in operator algebras, Encyclopaedia Math. Sci., 126, 1–145, Springer, Berlin, 2002. MR1878882 (2003i:46060)

- [23] C. Weibel, *An introduction to homological algebra*, Cambridge Univ. Press, 1994. MR1269324 (95f:18001)
- [24] M. Wodzicki, *Excision in cyclic homology and in rational algebraic K-theory*. Ann. of Math. **129** (1989) 591–639. MR997314 (91h:19008)

DEPARTAMENT DE MATEMÀTIQUES, UNIVERSITAT AUTÒNOMA DE BARCELONA, 08193 BELLA-TERRA (BARCELONA), SPAIN

*E-mail address:* `para@mat.uab.cat`

DEPARTAMENTO DE MATEMÁTICA AND INSTITUTO SANTALÓ, CIUDAD UNIVERSITARIA PAB 1, 1428 BUENOS AIRES, ARGENTINA

*E-mail address:* `gcorti@dm.uba.ar`

*URL:* `http://mate.dm.uba.ar/~gcorti`