

SPECTRAL ISOMETRIES ON NON-SIMPLE C^* -ALGEBRAS

MARTIN MATHIEU AND AHMED R. SOUROUR

(Communicated by Thomas Schlumprecht)

ABSTRACT. We prove that unital surjective spectral isometries on certain non-simple unital C^* -algebras are Jordan isomorphisms. Along the way, we establish several general facts in the setting of semisimple Banach algebras.

1. INTRODUCTION

Let A and B be unital C^* -algebras. A linear mapping $T: A \rightarrow B$ preserving the spectral radius of every element, that is, $r(Ta) = r(a)$ for every $a \in A$, is called a *spectral isometry*. This is a seemingly very weak property, and one wonders how much more of the structure of a C^* -algebra such a mapping may be able to preserve. It is known that if A is commutative and T is unital, that is, $T1_A = 1_B$, and surjective, then T is an algebra isomorphism [12, Proposition 2.2]. It is also known that if A is finite-dimensional and T is a unital surjective spectral isometry, then T preserves squares, that is, $T(a^2) = (Ta)^2$ for every $a \in A$ [17, Corollary 5]; see also [6, Corollary 1.4]. The latter property of T implies that T preserves the Jordan structure; that is, it is a *Jordan homomorphism*: $T(ab + ba) = (Ta)(Tb) + (Tb)(Ta)$ for all $a, b \in A$. In the non-commutative setting, this is the most one can hope for.

Examples of spectral isometries are spectrum-preserving mappings, that is, $\sigma_B(Ta) = \sigma_A(a)$ for every $a \in A$. These have been studied intensively over the past decades. Aupetit showed in [4] that every surjective spectrum-preserving linear mapping between von Neumann algebras is a bijective Jordan homomorphism (i.e., a *Jordan isomorphism*). It is not known (yet) whether this extends to general unital C^* -algebras. Nevertheless, in [15], the first-named author conjectured that this conclusion should hold for all unital surjective spectral isometries. Many results confirming this conjecture under additional hypotheses have appeared over the last few years; see [5], [7], [8], [12], [13], [14] and [17], for example. Sometimes, the conclusion that T preserves the Jordan product is even valid under the weaker assumption that T is *spectrally bounded*, i.e., $r(Ta) \leq M r(a)$ for all $a \in A$ and some constant $M > 0$; cf. [10], [11], [16]. However, in the presence of traces, one cannot expect such a general statement; for a detailed discussion of this, see Section 5 in [13].

Received by the editors July 25, 2011 and, in revised form, February 16, 2012.

2010 *Mathematics Subject Classification*. Primary 47A65, 47A10, 47B48, 46H10, 46L05.

Key words and phrases. Spectral isometries, spectrally bounded operators, Jordan isomorphisms, C^* -algebras.

This paper was written during a visit of the first-named author to the University of Victoria in May 2011, supported by a Research in Pairs grant of the London Mathematical Society.

The second-named author's research was supported by an NSERC Discovery Grant.

Some of the methods used to obtain the conclusion that a spectral isometry is a Jordan isomorphism are global in nature; see, for instance, [7], [10] or [16]. Others are reduction techniques using suitable quotients of the algebras in question. In this paper, we shall follow this second approach by combining ideas from [8], [13], [14] and [17]. Our main result, Theorem 3.6, states that a unital spectral isometry from a unital C^* -algebra with real rank zero and without tracial states that contains enough maximal ideals onto a unital C^* -algebra must be a Jordan isomorphism. This is obtained by first working out a reduction to simple quotients which is valid in general semisimple Banach algebras in Section 2 and then applying results known for simple C^* -algebras, in particular [11]. The same approach works for separable unital C^* -algebras with Hausdorff spectrum (Theorem 3.5), thus generalising results obtained in [14].

We conclude this introduction with a few comments on the general setting of our results. Suppose $T: A \rightarrow B$ is a spectral isometry between two unital Banach algebras A and B . The surjectivity assumption is inevitable, as is well known. If the domain of T is semisimple, then T is injective [15, Proposition 4.2] and the inverse T^{-1} is a bijective spectral isometry as well. In this case, the codomain is semisimple too (Lemma 2.1 below), and it follows that T is bounded [1, Corollary 5.46]. The open mapping theorem entails that T^{-1} is bounded as well. When T is non-unital and A and B are C^* -algebras, then $T1$ is a central unitary. Thus, replacing T by \tilde{T} defined by $\tilde{T}x = (T1)^{-1}Tx$, $x \in A$, we can reduce the general to the unital case (see the proof of [10, Corollary 2.6]).

2. PREPARATIONS

In this section we collect several auxiliary results needed for the main results in the next section. These are in fact valid in the framework of (semisimple) Banach algebras. When A is a Banach algebra, $\text{rad}(A)$ will denote its Jacobson radical (see, e.g., [1, 5.3] or [2, 3.1.3]) and $Z(A)$ its centre.

The first lemma is easily deduced from known properties of spectral isometries (see, e.g., [15]), but we include a proof for completeness.

Lemma 2.1. *Let A and B be unital Banach algebras. Let $T: A \rightarrow B$ be a surjective spectral isometry. Then $T \text{rad}(A) = \text{rad}(B)$.*

Proof. Take $a \in \text{rad}(A)$ and $y \in B$ such that $r(y) = 0$. Choose $x \in A$ with $y = Tx$; then, by hypothesis, $r(x) = r(y) = 0$. It follows that

$$r(Ta + y) = r(T(a + x)) = r(a + x) = 0$$

so that $Ta \in \text{rad}(B)$, by Zemánek's characterisation of the radical [1, Theorem 5.40].

Conversely, take $b \in \text{rad}(B)$ and let $a \in A$ be such that $b = Ta$. Let $x \in A$ be quasinilpotent. Then

$$r(a + x) = r(T(a + x)) = r(b + Tx) = 0$$

since Tx is quasinilpotent. As before, it follows that $a \in \text{rad}(A)$; therefore $b \in T \text{rad}(A)$. We conclude that $T \text{rad}(A) = \text{rad}(B)$. \square

In particular, the image of a semisimple Banach algebra under a surjective spectral isometry is semisimple.

The next result, too, is an immediate consequence of known results together with Lemma 2.1. Compare this with [12, Proposition 2.2].

Lemma 2.2. *Let $T: A \rightarrow B$ be a unital surjective spectral isometry from the semisimple unital Banach algebra A onto the unital Banach algebra B . If A is commutative, then B is commutative.*

Proof. By Lemma 2.1, B is semisimple, and by [15, Corollary 4.4], $B = TA = TZ(A) = Z(B)$ so that B is commutative. \square

The following observation allows us to pass conveniently to the quotient of a spectral isometry.

Lemma 2.3. *Let $S: B \rightarrow B$ be a unital surjective spectral isometry on a unital Banach algebra B . Let I be a closed ideal of B such that the quotient algebra B/I is semisimple. If $r(Sy + I) = r(y + I)$ for all $y \in B$, then S induces a unital surjective spectral isometry $S^I: B/I \rightarrow B/I$ such that $S^I(y + I) = Sy + I$ for all $y \in B$.*

Proof. Clearly, all we need to show is that $SI \subseteq I$; in this case, $S^I(y + I) = Sy + I$, $y \in B$, defines a unital surjective mapping on B/I which is a spectral isometry by assumption.

Let $x \in I$ and take $y \in B$. Choose $x' \in B$ such that $y = Sx'$. For each $\lambda \in \mathbb{C}$, we have

$$\begin{aligned} r(\lambda(Sx + I) + y + I) &= r(\lambda Sx + y + I) \\ &= r(S(\lambda x + x') + I) \\ &= r(\lambda x + x' + I) = r(x' + I). \end{aligned}$$

Applying Liouville's theorem [1, Corollary 5.43] to the subharmonic function $\lambda \mapsto r(\lambda(Sx + I) + y + I)$ we find that

$$r(Sx + I + y + I) = r(y + I) \quad (y \in B).$$

Zemánek's characterisation of the radical implies that $Sx + I \in \text{rad}(B/I) = 0$, which yields that $Sx \in I$. \square

We now establish a key result for our main theorem; the method of proof is inspired by [8]. Here, and in the following, X^c denotes the commutant

$$X^c = \{y \in A \mid yx = xy \text{ for all } x \in X\}$$

of a subset $X \subseteq A$ in A .

Lemma 2.4. *Let A and B be unital semisimple Banach algebras, and let $T: A \rightarrow B$ be a unital surjective spectral isometry. Let I be a closed ideal of B such that B/I is semisimple and each unital surjective spectral isometry $S: B/I \rightarrow B/I$ is multiplicative or anti-multiplicative. Let $a \in A$ and put $A_0 = \{a\}^{cc}$. Let $B_0 = TA_0$. For all $b_1, b_2 \in B_0$ and $x \in \{a\}^c$, we have*

$$(2.1) \quad T^{-1}(b_1 b_2) x + I = x T^{-1}(b_1 b_2) + I.$$

Proof. Let $x \in \{a\}^c$ and $\lambda \in \mathbb{C}$. Using the inverse spectral isometry $T^{-1}: B \rightarrow A$ we define $S_{\lambda,x}: B \rightarrow B$ by $S_{\lambda,x}(y) = T(e^{-\lambda x} T^{-1}(y) e^{\lambda x})$, $y \in B$. Then $S_{\lambda,x}$ is a unital surjective spectral isometry of B with $S_{0,x} = \text{id}_B$ and $S_{\lambda,x}|_{B_0} = \text{id}_{B_0}$. We have

$$r(S_{\lambda,x}(y) + I) \leq r(T(e^{-\lambda x} T^{-1}(y) e^{\lambda x})) = r(y)$$

independent of x and λ . Therefore the subharmonic function $\lambda \mapsto r(S_{\lambda,x}(y) + I)$ is constant by Liouville's theorem and, taking $\lambda = 0$, we obtain

$$(2.2) \quad r(S_{\lambda,x}(y) + I) = r(y + I) \quad (y \in B, x \in \{a\}^c, \lambda \in \mathbb{C}).$$

By Lemma 2.3, $S_{\lambda,x}I \subseteq I$ and we obtain an induced unital surjective spectral isometry $S^I = S_{\lambda,x}^I$ on B/I . By hypothesis, S^I is either multiplicative or anti-multiplicative. Evidently, $S_{0,x}^I$ is the identity on B/I .

Suppose B/I is commutative. Then, by Nagasawa's theorem, S^I is multiplicative. Hence, for the purpose of the remaining argument, we can assume that B/I is non-commutative. For fixed x , the set $\{\lambda \in \mathbb{C} \mid S_{\lambda,x}^I \text{ is multiplicative}\}$ contains 0 and is closed since $\lambda \mapsto S_{\lambda,x}^I$ is continuous. Suppose that the above set is not open and take a point λ_0 in it which is an adherence point of the complement. Using the continuity of $\lambda \mapsto S_{\lambda,x}^I$ again, we conclude that

$$S_{\lambda_0,x}^I(b_1 + I) S_{\lambda_0,x}^I(b_2 + I) = S_{\lambda_0,x}^I(b_2 + I) S_{\lambda_0,x}^I(b_1 + I) \quad (b_1, b_2 \in B),$$

as $S_{\lambda_0,x}^I$ is both multiplicative and anti-multiplicative. However, as $S_{\lambda_0,x}^I$ is surjective, it follows that B/I is commutative, a contradiction. We thus conclude that $S_{\lambda,x}^I$ is an algebra isomorphism for all $\lambda \in \mathbb{C}$ and every $x \in \{a\}^c$.

Let $b_1, b_2 \in B_0$. As $S_{\lambda,x}(b_i) = b_i, i = 1, 2$, we obtain

$$S_{\lambda,x}^I(b_1 b_2 + I) = S_{\lambda,x}^I(b_1 + I) S_{\lambda,x}^I(b_2 + I) = (b_1 + I)(b_2 + I)$$

or, equivalently,

$$(2.3) \quad T(e^{-\lambda x} T^{-1}(b_1 b_2) e^{\lambda x}) + I = (b_1 + I)(b_2 + I) \quad (\lambda \in \mathbb{C}).$$

Taking derivatives in (2.3), evaluating at $\lambda = 0$ and using the fact that T is injective, we find that

$$(2.4) \quad T^{-1}(b_1 b_2) x - x T^{-1}(b_1 b_2) + I = 0,$$

as claimed. □

3. RESULTS

Our first result in this section explains the significance of commutative subalgebras for the general conjecture. For an element a in a unital C^* -algebra A , $C^*(a)$ stands for the unital C^* -subalgebra of A generated by a .

Proposition 3.1. *Let A be a unital C^* -algebra and let B be a unital Banach algebra. The following conditions on a unital surjective spectral isometry $T: A \rightarrow B$ are equivalent:*

- (a) T is a Jordan isomorphism;
- (b) TA_0 is a subalgebra of B for every commutative unital subalgebra A_0 of A ;
- (c) $T(\{a\}^{cc})$ is a subalgebra of B for every element $a \in A_{sa}$;
- (d) $TC^*(a)$ is a subalgebra of B for every element $a \in A_{sa}$.

Proof. By Lemma 2.1, B is automatically semisimple.

(a) \Rightarrow (b) It is well known that every Jordan isomorphism between semisimple algebras preserves commutativity (see, e.g., [9]) and therefore restricts to a multiplicative mapping on each commutative subalgebra. As a result, TA_0 is a subalgebra of B for every commutative unital subalgebra A_0 of A .

(b) \Rightarrow (c) and (b) \Rightarrow (d) are trivial.

We finally show that (c) and (d) individually imply (a). To this end, set $A_0 = \{a\}^{cc}$ or $A_0 = C^*(a)$, respectively, for an arbitrary fixed element $a \in A_{sa}$. Note that $\{a\}^{cc}$ is a C^* -subalgebra by the Fuglede–Putnam theorem. Since T is a topological isomorphism between A and B , the subspace $B_0 = TA_0$ is closed in B . By hypothesis, B_0 is thus a unital Banach algebra. Let $T_0: A_0 \rightarrow B_0$ denote the restriction of T to A_0 which is a unital surjective spectral isometry. By Lemma 2.1, B_0 is semisimple, and by Lemma 2.2, B_0 is commutative. Nagasawa’s theorem [2, Theorem 4.1.17] entails that T_0 is multiplicative; therefore $T(a^2) = T_0(a^2) = (T_0a)^2 = (Ta)^2$. It follows that T is a Jordan isomorphism. \square

Remark 3.2. If in the above proposition A is merely assumed to be a unital semisimple Banach algebra instead of a unital C^* -algebra, the condition that $T(\{a\}^{cc})$ is a subalgebra of B for every element $a \in A$ still yields the fact that T is invertibility preserving. This can be seen by using Lemma 2.1 to obtain $T_0 \operatorname{rad}(A_0) = \operatorname{rad}(B_0)$, with $A_0 = \{a\}^{cc}$ and $B_0 = TA_0$, and therefore a unital surjective spectral isometry $T_1: A_1 = A_0/\operatorname{rad}(A_0) \rightarrow B_1 = B_0/\operatorname{rad}(B_0)$, which must be an isomorphism by Nagasawa’s theorem. It follows that

$$\sigma_A(a) = \sigma_{A_0}(a) = \sigma_{A_1}(a + \operatorname{rad}(A_0)) = \sigma_{B_1}(T_0a + \operatorname{rad}(B_0)) = \sigma_{B_0}(T_0a) \supseteq \sigma_B(Ta).$$

We will now combine Proposition 3.1 with the results in Section 2.

Proposition 3.3. *Let A and B be unital semisimple Banach algebras, and let $T: A \rightarrow B$ be a unital surjective spectral isometry. Suppose that B has a separating family \mathcal{S} of closed ideals I such that B/I is semisimple and each unital surjective spectral isometry $S^I: B/I \rightarrow B/I$ is multiplicative or anti-multiplicative. Then T preserves invertibility. If, moreover, A is a C^* -algebra, then T is a Jordan isomorphism.*

Proof. Let $a \in A$ and put $A_0 = \{a\}^{cc}$. Let $B_0 = TA_0$. Take $b_1, b_2 \in B_0$ and $x \in \{a\}^c$. By Lemma 2.4, $T^{-1}(b_1b_2)x + I = xT^{-1}(b_1b_2) + I$ for every $I \in \mathcal{S}$. As \mathcal{S} is separating, this yields $T^{-1}(b_1b_2)x = xT^{-1}(b_1b_2)$, which entails that $T^{-1}(b_1b_2) \in A_0$; equivalently, $b_1b_2 \in B_0$. We conclude that B_0 is a (closed, unital) subalgebra of B , and Remark 3.2 gives the first assertion.

If, moreover, A is a C^* -algebra, the above argument shows that $T(\{a\}^{cc})$ is a subalgebra of B for every element $a \in A_{sa}$. Now Proposition 3.1 completes the proof. \square

Applying the above general principles in various concrete situations enables us to obtain new incidences in which every unital surjective spectral isometry is a Jordan isomorphism.

The next two results were obtained in [14] under the additional assumption that the spectrum of the domain algebra is totally disconnected.

Theorem 3.4. *Let $T: A \rightarrow B$ be a unital surjective spectral isometry from a unital type I C^* -algebra with Hausdorff primitive ideal space onto a unital C^* -algebra B . Then T is a Jordan isomorphism.*

Proof. By replacing T with T^{-1} , we can interchange the roles of A and B . If the primitive ideal space $\operatorname{Prim}(B)$ is Hausdorff, every primitive ideal is maximal, and thus every irreducible representation π maps onto a simple unital C^* -algebra. As we assume B to be of type I, every irreducible image $\pi(B)$ contains the compact operators and, consequently, is a finite-dimensional simple unital C^* -algebra, that

is, is of the form $M_n(\mathbb{C})$. Every unital spectral isometry on $M_n(\mathbb{C})$ is multiplicative or anti-multiplicative ([3, Proposition 2]; for an alternative argument, see [13, Example 5.4]). Therefore, the hypotheses of Proposition 3.3 are satisfied, and thus T is a Jordan isomorphism. \square

Theorem 3.5. *Let $T: A \rightarrow B$ be a unital surjective spectral isometry from a separable unital C^* -algebra with Hausdorff spectrum onto a unital C^* -algebra B . Then T is a Jordan isomorphism.*

Proof. If the spectrum of A , that is, the space of all equivalence classes of irreducible representations, is Hausdorff then it is homeomorphic to $\text{Prim}(A)$; hence all irreducible images are unital and simple. Since they are separable too, Glimm's theorem [18, Theorem 6.8.7] entails that they are of the form $M_n(\mathbb{C})$ (i.e., A is of type I); compare [14, Corollary 12]. Now the proof is completed as above. \square

The class of type I C^* -algebras with Hausdorff spectrum is quite large; it encompasses in particular all continuous-trace C^* -algebras. For a more detailed discussion, see [14]. On the other hand, our methods also apply to very different non-simple C^* -algebras which are far from type I.

Theorem 3.6. *Let A and B be unital C^* -algebras, and let $T: A \rightarrow B$ be a unital surjective spectral isometry. Suppose that A has real rank zero and no tracial states and that $\text{Prim}(A)$ contains a dense subset of closed points. Then T is a Jordan isomorphism.*

Proof. As before we swap the roles of A and B to obtain the additional assumptions on B . By hypothesis, there exists a separating family \mathcal{S} of maximal ideals of B . Let $I \in \mathcal{S}$. Then B/I has real rank zero and no tracial states (and, of course, is unital simple). By [11, Theorem 3.1], every unital spectral isometry from B/I onto itself is a Jordan isomorphism, and it is well known that Jordan isomorphisms of simple algebras are either multiplicative or anti-multiplicative. Consequently, we can apply Proposition 3.3 to obtain the result. \square

Remark 3.7. The reduction theory employed in this paper is closely related to the reduction via Glimm ideals used in [17], [14] and [7]. For a unital C^* -algebra A , $\text{Prim}(A)$ is Hausdorff if and only if every Glimm ideal of A is maximal [14, Lemma 9]. Moreover, the condition that $T(\{a\}^{cc})$ is an algebra for every selfadjoint $a \in A$ implies that T is Z -multiplicative, i.e., $T(xz) = (Tx)(Tz)$ for all $x \in A$ and all $z \in Z(A)$. This in turn entails invariance of Glimm ideals, which is the essential tool in the Glimm reduction procedure.

REFERENCES

- [1] Graham R. Allan, *Introduction to Banach spaces and algebras*, Oxford Graduate Texts in Mathematics, vol. 20, Oxford University Press, Oxford, 2011. Prepared for publication and with a preface by H. Garth Dales. MR2761146 (2012j:46001)
- [2] Bernard Aupetit, *A primer on spectral theory*, Universitext, Springer-Verlag, New York, 1991. MR1083349 (92c:46001)
- [3] Bernard Aupetit, *Spectral characterization of the radical in Banach and Jordan-Banach algebras*, Math. Proc. Cambridge Philos. Soc. **114** (1993), no. 1, 31–35, DOI 10.1017/S0305004100071371. MR1219911 (94e:46081)
- [4] Bernard Aupetit, *Spectrum-preserving linear mappings between Banach algebras or Jordan-Banach algebras*, J. London Math. Soc. (2) **62** (2000), no. 3, 917–924, DOI 10.1112/S0024610700001514. MR1794294 (2001h:46078)

- [5] M. Bendaoud, A. Bourhim, and M. Sarih, *Linear maps preserving the essential spectral radius*, Linear Algebra Appl. **428** (2008), no. 4, 1041–1045, DOI 10.1016/j.laa.2007.09.010. MR2382111 (2008m:47051)
- [6] Constantin Costara, *A Cartan type theorem for finite-dimensional algebras*, Linear Algebra Appl. **426** (2007), no. 2-3, 299–304, DOI 10.1016/j.laa.2007.04.022. MR2347842 (2008g:46068)
- [7] C. Costara, *Commuting holomorphic maps on the spectral unit ball*, Bull. Lond. Math. Soc. **41** (2009), no. 1, 57–62, DOI 10.1112/blms/bdn104. MR2481989 (2010c:47093)
- [8] Constantin Costara and Dušan Repovš, *Spectral isometries onto algebras having a separating family of finite-dimensional irreducible representations*, J. Math. Anal. Appl. **365** (2010), no. 2, 605–608, DOI 10.1016/j.jmaa.2009.11.040. MR2587062 (2011f:46053)
- [9] N. Jacobson and C. E. Rickart, *Jordan homomorphisms of rings*, Trans. Amer. Math. Soc. **69** (1950), 479–502. MR0038335 (12,387h)
- [10] Ying-Fen Lin and Martin Mathieu, *Jordan isomorphism of purely infinite C^* -algebras*, Q. J. Math. **58** (2007), no. 2, 249–253, DOI 10.1093/qmath/hal024. MR2334865 (2008k:46189)
- [11] Martin Mathieu, *Spectrally bounded operators on simple C^* -algebras. II*, Irish Math. Soc. Bull. **54** (2004), 33–40. MR2138429 (2005m:47074)
- [12] Martin Mathieu, *Towards a non-selfadjoint version of Kadison’s theorem*, Ann. Math. Inform. **32** (2005), 87–94. MR2264870 (2008a:46057)
- [13] Martin Mathieu, *A collection of problems on spectrally bounded operators*, Asian-Eur. J. Math. **2** (2009), no. 3, 487–501, DOI 10.1142/S1793557109000418. MR2568008 (2011a:47079)
- [14] Martin Mathieu and Conal Ruddy, *Spectral isometries. II*, Function spaces, Contemp. Math., vol. 435, Amer. Math. Soc., Providence, RI, 2007, pp. 301–309, DOI 10.1090/conm/435/08385. MR2359437 (2008i:47033)
- [15] Martin Mathieu and Gerhard J. Schick, *First results on spectrally bounded operators*, Studia Math. **152** (2002), no. 2, 187–199, DOI 10.4064/sm152-2-6. MR1916549 (2003e:47066)
- [16] Martin Mathieu and Gerhard J. Schick, *Spectrally bounded operators from von Neumann algebras*, J. Operator Theory **49** (2003), no. 2, 285–293. MR1991740 (2004c:47079)
- [17] Martin Mathieu and Ahmed Ramzi Sourour, *Hereditary properties of spectral isometries*, Arch. Math. (Basel) **82** (2004), no. 3, 222–229, DOI 10.1007/s00013-003-0595-5. MR2053626 (2004m:47073)
- [18] Gert K. Pedersen, *C^* -algebras and their automorphism groups*, London Mathematical Society Monographs, vol. 14, Academic Press Inc. [Harcourt Brace Jovanovich Publishers], London, 1979. MR548006 (81e:46037)

DEPARTMENT OF PURE MATHEMATICS, QUEEN’S UNIVERSITY BELFAST, BELFAST BT7 1NN,
NORTHERN IRELAND

E-mail address: m.m@qub.ac.uk

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF VICTORIA, VICTORIA, BC,
CANADA V8W 3R4

E-mail address: sourour@math.uvic.ca