

TRACE ASYMPTOTICS FOR C^* -ALGEBRAS FROM SMALE SPACES

D. B. KILLOUGH AND I. F. PUTNAM

(Communicated by Varghese Mathai)

ABSTRACT. We consider C^* -algebras associated with stable and unstable equivalence in hyperbolic dynamical systems known as Smale spaces. These systems include shifts of finite type, in which case these C^* -algebras are both AF-algebras. These algebras have fundamental representations on a single Hilbert space (subject to a choice of periodic points) which have a number of special properties. In particular, the product between any element of the first algebra with one from the second is compact. In addition, there is a single unitary operator which implements actions on both. Here, under the hypothesis that the system is mixing, we show that the (semi-finite) traces on these algebras may be obtained through a limiting process and the usual operator trace.

1. INTRODUCTION

1.1. Smale spaces. A Smale space, as defined by David Ruelle [14], is a compact metric space, X , together with a homeomorphism, φ , which is hyperbolic. These include the basic sets of Smale's Axiom A systems [17]. Another special case of great interest are the shifts of finite type [1, 8].

Informally, the structure of (X, φ) is such that, for each point x in X and each $\epsilon > 0$ and sufficiently small, there are sets $X^s(x, \epsilon)$ and $X^u(x, \epsilon)$, called the local stable and unstable sets respectively. Their Cartesian product is homeomorphic to a neighbourhood of x (in a canonical and φ -invariant way) and the map φ is (uniformly) contracting on the former while φ^{-1} is contracting on the latter.

The precise axiom for a Smale space is the existence of a map defined on pairs (x, y) in $X \times X$ which are sufficiently close. We have a constant $\epsilon_X > 0$ and for x, y in X with $d(x, y) \leq \epsilon_X$, the image of (x, y) is denoted $[x, y]$. This map satisfies a number of identities and we refer the reader to [14] or [9] for a complete discussion. From the map $[,]$, we then define $X^s(x, \epsilon) = \{y \in X \mid d(x, y) < \epsilon, [y, x] = x\}$ and $X^u(x, \epsilon) = \{y \in X \mid d(x, y) < \epsilon, [x, y] = x\}$, for any $0 < \epsilon \leq \epsilon_X$. The contracting and expanding properties of the map φ on $X^s(x, \epsilon)$ and $X^u(x, \epsilon)$ are that there is a constant $0 < k_e < 1$ such that

$$\begin{aligned}d(\varphi(y), \varphi(z)) &\leq k_e d(y, z), y, z \in X^s(x, \epsilon), \\d(\varphi^{-1}(y), \varphi^{-1}(z)) &\leq k_e d(y, z), y, z \in X^u(x, \epsilon),\end{aligned}$$

which are axioms.

Received by the editors August 24, 2012 and, in revised form, April 8, 2013.
2010 *Mathematics Subject Classification*. Primary 37D20; Secondary 46L55, 46L51.
The second author was supported in part by an NSERC Discovery Grant.

There is also a notion of a global stable (unstable) set for a point x , which we denote $X^s(x)$ ($X^u(x)$). This is simply the set of all points $y \in X$ such that $d(\varphi^n(x), \varphi^n(y)) \rightarrow 0$ as $n \rightarrow +\infty$ ($-\infty$). The collection of sets $\{X^s(y, \delta) \mid y \in X^s(x), \delta > 0\}$ forms a neighbourhood base for a topology on $X^s(x)$ in which it is locally compact and Hausdorff. This is the topology that we use on $X^s(x)$ (not the relative topology from X). There is an analogous topology on $X^u(x)$. Briefly, we have two equivalence relations on X which are transverse in some sense. It is this transversality which lies at the core of the results in [3] and our main result below.

We say that (X, φ) is irreducible if, for every (ordered) pair of non-empty open sets U and V , there exists an $N \geq 1$ such that $\varphi^N(U) \cap V \neq \emptyset$. Also, we say that (X, φ) is mixing if, for every (ordered) pair of non-empty open sets U and V , there exists an $N_0 \geq 1$ such that $\varphi^N(U) \cap V \neq \emptyset$, for all $N \geq N_0$.

An important feature of an irreducible Smale space (X, φ) is the existence and uniqueness of a φ -invariant probability measure maximizing the entropy of φ , see [4, 16]. We call this the Bowen measure and denote it by μ_X , or when the space is obvious, simply μ . It was shown in [16] (but also see [5] for further discussion) that, as our space is locally a product space, the Bowen measure is locally a product measure. Specifically, for each x in X , we have measures $\mu^{s,x}$ and $\mu^{u,x}$ defined on $X^s(x)$ and $X^u(x)$, respectively. Secondly, the measure $\mu^{s,x}$ depends only on the stable equivalence class of x ; that is, if y is in $X^s(x)$, then $\mu^{s,x} = \mu^{s,y}$. (Put another way, we should be writing $\mu^{s, X^s(x)}$, but that notation is rather too clumsy.) A similar statement holds for $\mu^{u,x}$. It is worth noting that these measures are not finite, but are regular Borel measures. Moreover, these satisfy the following conditions (see [16] or [5] for details).

Theorem 1.1. (1) For all x in X , $\epsilon > 0$ and Borel sets $B \subset X^u(x, \epsilon)$ and $C \subset X^s(x, \epsilon)$, we have

$$\mu([B, C]) = \mu^{u,x}(B)\mu^{s,x}(C)$$

whenever ϵ is sufficiently small so that $[B, C]$ is defined.

(2) For x, y in X , $\epsilon > 0$ and a Borel set $B \subset X^u(x, \epsilon)$, we have

$$\mu^{u,y}([B, y]) = \mu^{u,x}(B),$$

whenever $d(x, y)$ and ϵ are sufficiently small so that $[B, y]$ is defined.

(3) For x, y in X , $\epsilon > 0$ and a Borel set $C \subset X^s(x, \epsilon)$, we have

$$\mu^{s,y}([y, C]) = \mu^{s,x}(C),$$

whenever $d(x, y)$ and ϵ are sufficiently small so that $[y, C]$ is defined.

(4) $\mu^{s, \varphi(x)} \circ \varphi = \lambda^{-1} \mu^{s,x}$.

(5) $\mu^{u, \varphi(x)} \circ \varphi = \lambda \mu^{u,x}$.

Here $\log(\lambda)$ is the topological entropy of (X, φ) .

In the future, it will cause no confusion to drop the superscript x .

1.2. C^* -algebras. Several C^* -algebras can be constructed from a given Smale space. The construction is originally due to Ruelle [15]. In the case of a shift of finite type, these algebras were defined and studied earlier by Krieger, Cuntz and Krieger in [2, 7]. In the more general Smale space setting we refer the reader to [9–11]. The algebras that we will be concerned with for a Smale space (X, φ) are known as the stable and unstable algebras.

We consider the groupoid C^* -algebras associated with stable and unstable equivalence. Here, it is most convenient to consider stable equivalence on unstable sets, since these function as abstract transversals. Also, it is convenient to have such sets which are φ -invariant. So we fix a finite φ -invariant set, P , (the periodic points in an irreducible Smale space are always dense, so there is a good supply of such sets) and let

$$X^s(P) = \bigcup_{p \in P} X^s(p), \quad X^u(P) = \bigcup_{p \in P} X^u(p).$$

We then define

$$\begin{aligned} G^s(X, \varphi, P) &= \{(x, y) \mid x \stackrel{s}{\sim} y, x, y \in X^u(P)\}, \\ G^u(X, \varphi, P) &= \{(x, y) \mid x \stackrel{u}{\sim} y, x, y \in X^s(P)\}. \end{aligned}$$

These groupoids have topologies, which will be described in detail in section 2, in which they are étale. We define $S(X, \varphi, P) = C^*(G^s(X, \varphi, P))$ and $U(X, \varphi, P) = C^*(G^u(X, \varphi, P))$.

We summarize some of the properties of $S(X, \varphi, P)$ and $U(X, \varphi, P)$. Both are simple if and only if (X, φ) is mixing. They are both amenable, stable and finite. In the case of a shift of finite type, each is an AF-algebra. We also point out that the homeomorphism φ yields an automorphism, α , of each of these C^* -algebras, which is induced by $\alpha(a)(x, y) = a(\varphi^{-1}(x), \varphi^{-1}(y))$.

Our system of measures $\mu^{u,x}$ and $\mu^{s,x}$ provide faithful, semi-finite traces on $S(X, \varphi, P)$ and $U(X, \varphi, P)$ as follows: $\sum_{x \in P} \mu^{u,x}$ defines a measure on the unit space of $G^s(X, \varphi, P)$ and the invariance property 2 of Theorem 1.1 implies that this can be used to define a trace. Again, we will denote this measure simply by μ^u . We summarize the following result. A version first appears in [9]. Our definition of $S(X, \varphi, Q)$, which differs from that in [9], can be found in [6, 10, 12], however, the proofs used in [9] apply with little alteration.

Theorem 1.2. *Let (X, φ) be an irreducible Smale space and let P, Q be finite φ -invariant subsets of X . For $a \in C_c(G^s(X, \varphi, Q))$ and $b \in C_c(G^u(X, \varphi, P))$, define*

$$\begin{aligned} \tau^s(a) &= \int_{X^u(Q)} a(x, x) d\mu^u, \text{ and} \\ \tau^u(b) &= \int_{X^s(P)} b(x, x) d\mu^s. \end{aligned}$$

Then τ^s and τ^u extend to semi-finite traces on $S(X, \varphi, Q)$ and $U(X, \varphi, P)$.

1.3. Fundamental representation. Here, we choose two finite φ -invariant sets, P and Q . We will consider the C^* -algebras $S(X, \varphi, Q)$ and $U(X, \varphi, P)$. Define

$$X^h(P, Q) = X^s(P) \cap X^u(Q).$$

(The ‘h’ stands for heteroclinic.) This is a subset of $X^u(Q)$, the unit space of $G^s(X, \varphi, Q)$ and consists of some stable equivalence classes in that set. Hence, the Hilbert space $l^2(X^h(P, Q))$ admits a natural representation of $S(X, \varphi, Q)$ which we will describe in detail in the next section. In an analogous way, it also admits a representation of $U(X, \varphi, P)$. We suppress these representations in our notation and simply assume that these C^* -algebras are acting on this Hilbert space. We also note that $X^h(P, Q)$ is countable and that $\delta_x, x \in X^h(P, Q)$ denotes the usual orthonormal basis. We note that $u\xi = \xi \circ \varphi^{-1}$ is a unitary operator on the Hilbert space and implements the automorphisms α of $S(X, \varphi, Q)$ and $U(X, \varphi, P)$.

This pair of C^* -algebras acting on this Hilbert space possesses a number of interesting features. We summarize the following results (6.1, 6.2 and 6.3 from [3]).

Theorem 1.3. *Let (X, φ) be a mixing Smale space and P, Q be finite φ -invariant sets.*

- (1) *If a is in $C_c(G^s(X, \varphi, Q))$ and b is in $C_c(G^u(X, \varphi, P))$, then ab and ba are both finite rank operators on $l^2(X^h(P, Q))$.*
- (2) *If a is in $S(X, \varphi, Q)$ and b is in $U(X, \varphi, P)$, then ab and ba are both compact operators on $l^2(X^h(P, Q))$.*
- (3) *Assuming that $P \cap Q = \emptyset$, if a is in $S(X, \varphi, Q)$ and b is in $U(X, \varphi, P)$, then*

$$\lim_{n \rightarrow +\infty} \alpha^{-n}(a)b = \lim_{n \rightarrow +\infty} b\alpha^{-n}(a) = 0.$$

- (4) *If a is in $S(X, \varphi, Q)$ and b is in $U(X, \varphi, P)$, then*

$$\lim_{n \rightarrow +\infty} \|\alpha^n(a)\alpha^{-n}(b) - \alpha^{-n}(b)\alpha^n(a)\| = 0.$$

Our main objective in this paper is to study the asymptotic behavior of $Tr(\alpha^n(a)\alpha^{-n}(b))$, for a in $C_c(G^s(X, \varphi, Q))$ and b in $C_c(G^u(X, \varphi, P))$ and relate it to $\tau^s(a)$ and $\tau^u(b)$.

2. MAIN RESULT

We consider the case that (X, φ) is a *mixing* Smale space and show how the above traces on $S(X, \varphi, Q)$ and $U(X, \varphi, P)$ are related to asymptotics of the usual trace on $\mathfrak{B}(l^2(X^h(P, Q)))$. That is, the formula

$$Tr(A) = \sum_{x \in X^h(P, Q)} \langle A\delta_x, \delta_x \rangle$$

defines a trace for trace class operators and can be extended to positive operators by allowing the value ∞ .

Theorem 2.1. *Let (X, φ) be a mixing Smale space with topological entropy $\log(\lambda)$, and let $a \in S(X, \varphi, Q)$, $b \in U(X, \varphi, P)$. If either*

- (1) *$a \in C_c(G^s(X, \varphi, Q))$ and $b \in C_c(G^u(X, \varphi, P))$, or*
- (2) *a and b are both positive,*

then

$$\lim_{k \rightarrow +\infty} \lambda^{-2k} Tr(\alpha^k(a)\alpha^{-k}(b)) = \tau^s(a)\tau^u(b).$$

In the second case, both sides of the equality may be $+\infty$.

To begin the proof, we must describe the topologies on $G^s(X, \varphi, Q)$ and $G^u(X, \varphi, P)$. We take the following from [9] or [18]. To consider the former, suppose that (x, y) is in $G^s(X, \varphi, Q)$. It follows that for some positive integer n , $\varphi^n(y) \in X^s(\varphi^n(x), \epsilon_X/2)$. Then the function

$$h^s(z) = \varphi^{-n}[\varphi^n(z), \varphi^n(x)]$$

sends y to x . It is also defined on $X^u(y, \delta)$ for some $\delta > 0$ and is a homeomorphism to its image in $X^u(x, \epsilon_X)$, which is open. Finally, we have that $h^s(z)$ is stably equivalent to z for all z in the domain. We define

$$X^s(x, y, \delta) = \{(h^s(z), z) \mid z \in X^u(y, \delta), h^s(z) \in X^u(x, \delta)\}.$$

We have $(x, y) \in X^s(x, y, \delta) \subset G^s(X, \varphi, Q)$. Such sets (allowing x, y, δ to vary) form a neighbourhood base for the topology of $G^s(X, \varphi, Q)$. The topology on $G^u(X, \varphi, P)$ is defined in an analogous way via functions denoted h^u and sets $X^u(x, y, \delta)$, respectively.

The representation of $S(X, \varphi, Q)$ is as follows, the representation of $U(X, \varphi, P)$ is completely analogous. Let a be a function with compact support in $X^s(x, y, \delta)$, so that a is in $C_c(G^s(X, \varphi, Q))$. In fact, every element of $C_c(G^s(X, \varphi, Q))$ is a sum of such functions. For $w \in X^h(P, Q)$, we will define the representations by

$$\begin{aligned} a\delta_w(x) &= \sum_{(x,y) \in G^s(X, \varphi, Q)} a(x, y)\delta_w(y) \\ &= \begin{cases} a(h^s(w), w)\delta_{h^s(w)} & \text{if } w \in X^u(y, \delta), h^s(w) \in X^u(x, \delta), \\ 0 & \text{if } w \notin X^u(y, \delta). \end{cases} \end{aligned}$$

For convenience, we now extend the class of functions that we consider on $G^s(X, \varphi, Q)$, $G^u(X, \varphi, P)$. As in Chapter II of [13] we consider the set of bounded Borel functions with compact support on these groupoids, denoted $B(G^s(X, \varphi, Q))$ ($B(G^u(X, \varphi, P))$ respectively). With convolution and involution defined as for the continuous functions, $B(G^s(X, \varphi, Q))$ is a $*$ -algebra. Furthermore, every representation of $C_c(G^s(X, \varphi, Q))$ extends to a representation of $B(G^s(X, \varphi, Q))$. This allows us to consider $Tr(a)$ where $a \in B(G^s(X, \varphi, Q))$. Moreover, since $B(G^s(X, \varphi, Q))$ consists of integrable functions, we can make the following definition.

Definition 2.2. For $a \in B(G^s(X, \varphi, Q))$ and $b \in B(G^u(X, \varphi, P))$, define

$$\int_{X^u(Q)} a(x, x)d\mu^u, \text{ and} \\ I^u(b) = \int_{X^s(P)} b(x, x)d\mu^s.$$

We start with the following lemma.

Lemma 2.3. *Let (X, φ) be a mixing Smale space. Let $a \in B(G^s(X, \varphi, Q))$ be supported on a set of the form $X^s(x_a, y_a, \delta_a)$, and $b \in B(G^u(X, \varphi, P))$ supported on $X^u(x_b, y_b, \delta_b)$. We use h_a^s to denote the function used in the definition of $X^s(x_a, y_a, \delta_a)$, and h_b^u to denote the function used in the definition of $X^u(x_b, y_b, \delta_b)$ (see the discussion after the statement of Theorem 2.1). Let $X_a = r(X(x_a, y_a, \delta_a))$, $X_b = s(X(x_b, y_b, \delta_b))$, where r, s are the range and source maps from $G^s(X, \varphi, Q)$ to $X^u(Q)$. If $h_a^s = id$ and $h_b^u = id$, then for each $k \in \mathbb{N}$*

$$Tr(\alpha^k(a)\alpha^{-k}(b)) = \sum_{w \in \varphi^k(X_a) \cap \varphi^{-k}(X_b)} a(\varphi^{-k}(w), \varphi^{-k}(w))b(\varphi^k(w), \varphi^k(w));$$

otherwise

$$\lim_{k \rightarrow +\infty} Tr(\alpha^k(a)\alpha^{-k}(b)) = 0.$$

Proof. If $x \neq h_a^s(y)$, $a(x, y) = 0$. To simplify notation, let $a(y) = a(h_a^s(y), y)$. Similarly, let $b(y) = b(h_b^u(y), y)$. Now

$$\begin{aligned} \text{Tr}(\alpha^k(a)\alpha^{-k}(b)) &= \sum_{w \in X^h(P)} \langle \alpha^k(a)\alpha^{-k}(b)\delta_w, \delta_w \rangle \\ &= \sum_{w \in X^h(P)} a(\varphi^{-2k}h_b^u\varphi^k(w))b(\varphi^k(w))\langle \delta_{\varphi^k h_a^s \varphi^{-2k} h_b^u \varphi^k(w)}, \delta_w \rangle. \end{aligned}$$

Suppose $h_a^s = h_b^u = id$. Then we have

$$\text{Tr}(\alpha^k(a)\alpha^{-k}(b)) = \sum_{w \in X^h(P)} a(\varphi^{-k}(w))b(\varphi^k(w))\langle \delta_w, \delta_w \rangle.$$

Moreover, $a(\varphi^{-k}(w)) \neq 0$ only if $w \in \varphi^k(X_a)$, and $b(\varphi^k(w)) \neq 0$ only if $w \in \varphi^{-k}(X_b)$, so

$$\text{Tr}(\alpha^k(a)\alpha^{-k}(b)) = \sum_{w \in \varphi^k(X_a) \cap \varphi^{-k}(X_b)} a(\varphi^{-k}(w))b(\varphi^k(w)).$$

Now suppose $h_a^s \neq id$.

$$\text{Tr}(\alpha^k(a)\alpha^{-k}(b)) = \sum_{w \in E_k} a(\varphi^{-2k}h_b^u\varphi^k(w))b(\varphi^k(w))\langle \delta_{\varphi^k h_a^s \varphi^{-2k} h_b^u \varphi^k(w)}, \delta_w \rangle,$$

where

$$E_k = \{w \in X^h(P) \mid \varphi^k(w) \in X_b, \varphi^{-2k}h_b^u\varphi^k(w) \in X_a, \varphi^k h_a^s \varphi^{-2k} h_b^u \varphi^k(w) = w\}.$$

We show that for large enough k , the set E_k is empty. We can find $\delta > 0$ such that

$$d(z, h_a^s(z)) > \delta$$

for all $z \in X_a$. Now, we can find K sufficiently large so that for all $k > K$ we have

$$d(\varphi^{-2k}(y), \varphi^{-2k}h_b^u(y)) < \delta$$

for all $y \in X_b$. So for $w \in E_k$, $\varphi^k(w) \in X_b$ and thus

$$d(\varphi^{-k}(w), \varphi^{-2k}h_b^u(\varphi^k(w))) < \delta$$

but $\varphi^{-2k}h_b^u(\varphi^k(w)) \in X_a$ so

$$d(\varphi^{-2k}h_b^u(\varphi^k(w)), h_a^s\varphi^{-2k}h_b^u(\varphi^k(w))) > \delta$$

so $\varphi^{-k}(w) \neq h_a^s\varphi^{-2k}h_b^u(\varphi^k(w))$, contradicting $w \in E_k$. Hence for large k , E_k is empty and

$$\text{Tr}(\alpha^k(a)\alpha^{-k}(b)) = 0.$$

A similar argument gives the result in the case $h_b^u \neq id$. □

Lemma 2.4. *Let (X, φ) be a mixing Smale space. Let $a \in B(G^s(X, \varphi, Q))$ and $b \in B(G^u(X, \varphi, P))$ be step functions. Then*

$$\lim_{k \rightarrow \infty} \lambda^{-2k} \text{Tr}(\alpha^k(a)\alpha^{-k}(b)) = I^s(a)I^u(b).$$

Proof. By Lemma 2.3 we can assume, without loss of generality, that $a(x, y) = 0$ whenever $x \neq y$, and $b(x, y) = 0$ if $x \neq y$. We therefore consider $a(x, x)$ to be supported on $A \subset X^s(x_a)$, and similarly with b .

Let $\{A_j\}_{j=1}^\infty$ be precompact sets on which a is constant, and a_j the value on A_j . Similarly $\{B_j\}_{j=1}^\infty$ and b_j . Then for each k we have

$$\begin{aligned} I^s(a) &= \sum_{j=1}^{\infty} \mu^u(A_j) a_j, \\ I^u(b) &= \sum_{j=1}^{\infty} \mu^u(B_j) b_j, \\ \text{Tr}(\alpha^k(a)\alpha^{-k}(b)) &= \sum_j \sum_l (\#h_{j,l}^k) a_j b_l, \end{aligned}$$

where $h_{j,l}^k = \varphi^k(A_j) \cap \varphi^{-k}(B_l)$. From Theorem 2.4 of [5] we know that

$$\lim_{k \rightarrow \infty} \lambda^{-2k} h_{j,l}^k = \mu^u(A_j) \mu^s(B_l).$$

So

$$\begin{aligned} \lim_{k \rightarrow \infty} \lambda^{-2k} \text{Tr}(\alpha^k(a)\alpha^{-k}(b)) &= \sum_j \sum_l \lim_{k \rightarrow \infty} \lambda^{-2k} \#h_{j,l}^k a_j b_l \\ &= \sum_j \sum_l \mu^u(A_j) \mu^s(B_l) a_j b_l \\ &= \left(\sum_j \mu^u(A_j) a_j \right) \left(\sum_l \mu^s(B_l) b_l \right) \\ &= I^s(a) I^u(b). \end{aligned}$$

□

Proof of Theorem 2.1. We start with the special case that $a \in C_c(G^s(X, \varphi, Q))$ and $b \in C_c(G^u(X, \varphi, P))$ are both positive. Lemma 2.3 allows us to assume, without loss of generality, that $a(x, y) = 0$ whenever $x \neq y$, and $b(x, y) = 0$ if $x \neq y$.

We find bounded Borel step functions with compact support a_l, a_u and b_l, b_u such that

$$a_l \leq a \leq a_u, \quad b_l \leq b \leq b_u, \\ I^s(a_l) + \epsilon > \tau^s(a) > I^s(a_u) - \epsilon, \quad \text{and} \quad I^u(b_l) + \epsilon > \tau^u(b) > I^u(b_u) - \epsilon.$$

For each k ,

$$\lambda^{-2k} \text{Tr}(\alpha^k(a_l)\alpha^{-k}(b_l)) \leq \lambda^{-2k} \text{Tr}(\alpha^k(a)\alpha^{-k}(b)) \leq \lambda^{-2k} \text{Tr}(\alpha^k(a_u)\alpha^{-k}(b_u)).$$

So, by Lemma 2.4, in the limit as $k \rightarrow \infty$ we have

$$I^s(a_l) I^u(b_l) \leq \lim_{k \rightarrow \infty} \lambda^{-2k} \text{Tr}(\alpha^k(a)\alpha^{-k}(b)) \leq I^s(a_u) I^u(b_u).$$

Therefore, for all $\epsilon > 0$, we have

$$(\tau^s(a) - \epsilon)(\tau^u(b) - \epsilon) < \lim_{k \rightarrow \infty} \lambda^{-2k} \text{Tr}(\alpha^k(a)\alpha^{-k}(b)) < (\tau^s(a) + \epsilon)(\tau^u(b) + \epsilon).$$

Hence, we conclude

$$\lim_{k \rightarrow \infty} \lambda^{-2k} \text{Tr}(\alpha^k(a)\alpha^{-k}(b)) = \tau^s(a) \tau^u(b).$$

To prove case (1) of the theorem (the C_c case) we simply apply the above argument separately to the positive/negative parts of the real/imaginary components of a, b .

Now suppose a, b are positive. If $\tau^s(a)$ and $\tau^u(b)$ are finite, then as above we can approximate above and below with (integrable) step functions, and by Lemma 2.4, the theorem holds. Now suppose one of the traces is not finite. Without loss of generality, assume $\tau^s(a) = +\infty$. We can then find $\{a_i\}$ such that $a_i \leq a$, a_i increasing, $a_i \in C_c(G^s(X, \varphi, Q))$ positive, and $\tau^s(a_i) \geq i$. Similarly choose $\{b_i\}$ (if $b \in C_c(G^u(X, \varphi, P))$), choose $b_i = b$ for all i). Then we have

$$\begin{aligned} \lim_{k \rightarrow \infty} \lambda^{-2k} \text{Tr}(\alpha^k(a)\alpha^{-k}(b)) &\geq \lim_{k \rightarrow \infty} \lambda^{-2k} \text{Tr}(\alpha^k(a_i)\alpha^{-k}(b_i)) \\ &= \tau^s(a_i)\tau^u(b_i) \\ &\geq i\tau^u(b_1). \end{aligned}$$

As this holds for any $i \in \mathbb{Z}$, we have

$$\lim_{k \rightarrow \infty} \lambda^{-2k} \text{Tr}(\alpha^k(a)\alpha^{-k}(b)) = \infty = \tau^s(a)\tau^u(b).$$

□

3. CONCLUDING REMARKS

The C^* -algebras $S(X, \varphi, Q)$ and $U(X, \varphi, P)$, with their fundamental representations, their natural automorphisms and the associated crossed products, have a number of remarkable features. Some of the most important are summarized in Theorem 1.3. In [3], this is used to prove a duality theorem which lives at the level of K-theory. Here, we move in a different direction by including the usual operator trace. A natural goal would be to establish a type of duality closer to that for vector spaces, in which one algebra is seen as linear functionals over the other. Our main result is a step in this direction.

ACKNOWLEDGMENT

The authors would like to thank the referee for many helpful suggestions.

REFERENCES

- [1] Rufus Bowen, *On Axiom A diffeomorphisms*, Regional Conference Series in Mathematics, No. 35, American Mathematical Society, Providence, R.I., 1978. MR0482842 (58 #2888)
- [2] Joachim Cuntz and Wolfgang Krieger, *A class of C^* -algebras and topological Markov chains*, Invent. Math. **56** (1980), no. 3, 251–268, DOI 10.1007/BF01390048. MR561974 (82f:46073a)
- [3] J. Kaminker, I. F. Putnam, and M. F. Whittaker, *K-theoretic duality for hyperbolic dynamical systems*. math. arxiv:1009.4999, 2010.
- [4] Anatole Katok and Boris Hasselblatt, with a supplementary chapter by Katok and Leonardo Mendoza, *Introduction to the modern theory of dynamical systems*, Encyclopedia of Mathematics and its Applications, vol. 54, Cambridge University Press, Cambridge, 1995. MR1326374 (96c:58055)
- [5] D. B. Killough and I. F. Putnam, *Bowen measure from heteroclinic points*, Canad. J. Math. **64** (2012), no. 6, 1341–1358, DOI 10.4153/CJM-2011-083-0. MR2994668
- [6] D. B. Killough and I. F. Putnam, *Ring and module structures on dimension groups associated with a shift of finite type*, Ergodic Theory Dynam. Systems **32** (2012), no. 4, 1370–1399, DOI 10.1017/S0143385712000272. MR2955318
- [7] Wolfgang Krieger, *On dimension functions and topological Markov chains*, Invent. Math. **56** (1980), no. 3, 239–250, DOI 10.1007/BF01390047. MR561973 (81m:28018)

- [8] Douglas Lind and Brian Marcus, *An introduction to symbolic dynamics and coding*, Cambridge University Press, Cambridge, 1995. MR1369092 (97a:58050)
- [9] Ian F. Putnam, *C^* -algebras from Smale spaces*, *Canad. J. Math.* **48** (1996), no. 1, 175–195, DOI 10.4153/CJM-1996-008-2. MR1382481 (97m:46106)
- [10] Ian F. Putnam, *Functoriality of the C^* -algebras associated with hyperbolic dynamical systems*, *J. London Math. Soc.* (2) **62** (2000), no. 3, 873–884, DOI 10.1112/S002461070000140X. MR1794291 (2002e:46067)
- [11] Ian F. Putnam, *Lifting factor maps to resolving maps*, *Israel J. Math.* **146** (2005), 253–280, DOI 10.1007/BF02773536. MR2151603 (2007i:37020)
- [12] Ian F. Putnam and Jack Spielberg, *The structure of C^* -algebras associated with hyperbolic dynamical systems*, *J. Funct. Anal.* **163** (1999), no. 2, 279–299, DOI 10.1006/jfan.1998.3379. MR1680475 (2000g:46098)
- [13] Jean Renault, *A groupoid approach to C^* -algebras*, *Lecture Notes in Mathematics*, vol. 793, Springer, Berlin, 1980. MR584266 (82h:46075)
- [14] David Ruelle, *Thermodynamic formalism*, *The mathematical structures of classical equilibrium statistical mechanics*, with a foreword by Giovanni Gallavotti and Gian-Carlo Rota, *Encyclopedia of Mathematics and its Applications*, vol. 5, Addison-Wesley Publishing Co., Reading, Mass., 1978. MR511655 (80g:82017)
- [15] David Ruelle, *Noncommutative algebras for hyperbolic diffeomorphisms*, *Invent. Math.* **93** (1988), no. 1, 1–13, DOI 10.1007/BF01393685. MR943921 (89k:58233)
- [16] David Ruelle and Dennis Sullivan, *Currents, flows and diffeomorphisms*, *Topology* **14** (1975), no. 4, 319–327. MR0415679 (54 #3759)
- [17] S. Smale, *Differentiable dynamical systems*, *Bull. Amer. Math. Soc.* **73** (1967), 747–817. MR0228014 (37 #3598)
- [18] Michael Whittaker, *Spectral triples for hyperbolic dynamical systems*, *J. Noncommut. Geom.* **7** (2013), no. 2, 563–582, DOI 10.4171/JNCG/127. MR3054307

DEPARTMENT OF MATHEMATICS, PHYSICS, AND ENGINEERING, MOUNT ROYAL UNIVERSITY,
CALGARY, ALBERTA, CANADA T3E 6K6
E-mail address: bkillough@mtroyal.ca

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF VICTORIA, VICTORIA, BRITISH
COLUMBIA, CANADA V8W 3R4
E-mail address: ifputnam@uvic.ca