

ON TOEPLITZ OPERATORS AND LOCALIZATION OPERATORS

LUÍS DANIEL ABREU AND NELSON FAUSTINO

(Communicated by Pamela B. Gorkin)

ABSTRACT. This note is a contribution to a problem of Lewis Coburn concerning the relation between Toeplitz operators and Gabor-Daubechies localization operators. We will show that, for any *localization operator* with a *general window* $w \in \mathcal{F}_2(\mathbb{C})$ (the Fock space of analytic functions square-integrable on the complex plane), there exists a differential operator of infinite order D , with constant coefficients explicitly determined by w , such that *the localization operator with symbol f coincides with the Toeplitz operator with symbol Df* . This extends results of Coburn, Lo and Engliš, who obtained similar results in the case where w is a polynomial window. Our technique of proof combines their methods with a direct sum decomposition in true polyanalytic Fock spaces. Thus, *polyanalytic functions are used as a tool to prove a theorem about analytic functions*.

1. INTRODUCTION

Let $\mathcal{L}_p(\mathbb{C})$ ($1 \leq p < \infty$) be the weighted L^p -space of functions with norm

$$\|u\|_{\mathcal{L}_p(\mathbb{C})} = \left(\int_{\mathbb{C}} |u(z)|^p e^{-\frac{\pi}{2}p|z|^2} dz \right)^{\frac{1}{p}},$$

dz being the area measure on \mathbb{C} and $\mathcal{L}_{\infty}(\mathbb{C})$ the space of measurable functions on \mathbb{C} such that $|u(z)|e^{-\frac{\pi}{2}|z|^2}$ is bounded, endowed with the norm

$$\|u\|_{\mathcal{L}_{\infty}(\mathbb{C})} = \sup_{z \in \mathbb{C}} |u(z)|e^{-\frac{\pi}{2}|z|^2}.$$

Let H be a Hilbert space contained in $\mathcal{L}_2(\mathbb{C})$, with reproducing kernel $K(\zeta, z)$. For $f \in L^{\infty}(\mathbb{C})$, the *Toeplitz operator* Toep_f with symbol $f(z)$ is the operator acting on H defined by

$$(\text{Toep}_f g)(\zeta) = \int_{\mathbb{C}} f(z)g(z)K(\zeta, z)e^{-\pi|z|^2} dz, \forall g \in H.$$

Received by the editors December 26, 2012 and, in revised form, March 20, 2013 and August 7, 2013.

2010 *Mathematics Subject Classification*. Primary 47B32, 30H20; Secondary 81R30, 81S30.

Key words and phrases. Localization operators, polyanalytic Fock spaces, Toeplitz operators.

Both authors were supported by CMUC and FCT (Portugal), through European program COMPETE/FEDER and by FCT project PTDC/MAT/114394/2009. The first author was also supported by Austrian Science Foundation (FWF) project “Frames and Harmonic Analysis” and START-project FLAME (“Frames and Linear Operators for Acoustical Modeling and Parameter Estimation”, Y 551-N13). The second author was also supported by São Paulo Research Foundation (FAPESP) through the grant 13/07590-8.

Denote by $\mathcal{F}_p(\mathbb{C})$ the space of entire functions with membership in $\mathcal{L}_p(\mathbb{C})$. The Weyl operator

$$(1.1) \quad W_z w(\zeta) = e^{\pi \bar{z} \zeta - \frac{\pi}{2} |z|^2} w(\zeta - z)$$

acts unitarily on $\mathcal{L}_2(\mathbb{C})$. If $w \in \mathcal{L}_2(\mathbb{C})$ and $f \in L^\infty(\mathbb{C})$, the *Gabor-Daubechies localization operator* $L_f^{(w)}$, with “window” w and “symbol” f , is the operator acting on $\mathcal{F}_2(\mathbb{C})$ defined, in the weak sense, by

$$\langle L_f^{(w)} u, v \rangle = \int_{\mathbb{C}} f(z) \langle u, W_z w \rangle_{d\mu} \langle W_z w, v \rangle_{d\mu} dz, \forall u, v \in \mathcal{F}_2(\mathbb{C}).$$

Considerable research activity on localization operators has been motivated by the article of Daubechies [6]. (See also [10], [5].) An interesting connection between localization and Toeplitz operators has been observed by Coburn [4].

In this note we will investigate further the relation between localization and Toeplitz operators. Our main contribution is simply outlined in the following paragraph.

Denote by $\mathcal{F}_p(\mathbb{C})$ the space of entire functions with membership in $\mathcal{L}_p(\mathbb{C})$. We will show that it is possible to write a localization operator with a window $w \in \mathcal{F}_2(\mathbb{C})$ as

$$L_f^{(w)} = \text{Toep}_{D(w)f}$$

where, in the right-hand side, we have a Toeplitz operator whose symbol is obtained from the symbol of the localization operator by the action of a differential operator $D(w)$, whose coefficients are constants explicitly determined by w .

Our methods of proof build on previous works by Coburn [4], Lo [13] and Engliš [7], but contain a technical innovation, based on *polyanalytic functions*.

A polyanalytic function of order n is a polynomial of order $n - 1$ in \bar{z} with analytic functions $\{\varphi_k(z)\}_{k=0}^{n-1}$ as coefficients:

$$(1.2) \quad F(z) = \sum_{k=0}^{n-1} \bar{z}^k \varphi_k(z),$$

so that an analytic function is a polyanalytic function of order 1. Alternatively, F satisfies the generalized Cauchy-Riemann equations

$$(1.3) \quad (\partial_{\bar{z}})^n F(z) = \frac{1}{2^n} \left(\frac{\partial}{\partial x} + i \frac{\partial}{\partial \xi} \right)^n F(x + i\xi) = 0.$$

A simple example of a polyanalytic function of order 2 is

$$F(z) = 1 - |z|^2 = 1 - z\bar{z}.$$

Observe that

$$\partial_{\bar{z}} F(z) = -z, \text{ while } (\partial_{\bar{z}})^2 F(z) = 0.$$

Polyanalytic functions have applications in signal analysis [1], [3], [2] and determinantal point processes [12].

We would like to emphasize that, in our proofs, *polyanalytic functions are used as a tool to prove a theorem about analytic functions*. Indeed, to extend the Coburn-Lo-Engliš theorem from the polynomial case to a general window, we will use an idea of Vasilevski [15], who has shown that arbitrary $\mathcal{L}_2(\mathbb{C})$ functions can be decomposed using a sequence of orthogonal spaces which are “slices” of Fock spaces of polyanalytic functions. Vasilevski coined the expression *true polyanalytic Fock spaces* to quote such slices. We will require a slightly more general version of his

result, since an essential step in the proof requires the decomposition of a non-analytic function of $\mathcal{L}_1(\mathbb{C})$ into polyanalytic basis functions. For this purpose we will use the direct sum decomposition of $\mathcal{L}_p(\mathbb{C})$ [3] into L^p -counterparts of *true* polyanalytic Fock spaces. Polyanalytic functions were already implicit in the papers of Coburn [4], Lo [13] and Engliš [7]. There is a time-frequency interpretation of this fact: if one takes a Hermite function as a window in a Gabor transform, the result is, up to a weight, a polyanalytic version of the Bargmann transform [1], [3].

The paper is organized in the following way: section 2 describes the main result and gives a brief account of its forerunners. In section 3 we collect the preliminary material about polyanalytic expansions and Engliš preparatory results which we will use in section 4, where the main result is proved.

2. MAIN RESULT

For an element $z = x + i\omega$ in \mathbb{C} , consider the Cauchy-Riemann operator and its conjugate,

$$\partial_{\bar{z}} = \frac{d}{d\bar{z}} = \frac{1}{2} \left(\frac{\partial}{\partial x} + i \frac{\partial}{\partial \omega} \right), \quad \partial_z = \frac{d}{dz} = \frac{1}{2} \left(\frac{\partial}{\partial x} - i \frac{\partial}{\partial \omega} \right).$$

It is clear from (1.1) that $W_z 1 = e^{\pi\bar{z}\zeta - \frac{\pi}{2}|z|^2}$ and $\langle F, W_z 1 \rangle_{d\mu} = e^{-\frac{\pi}{2}|z|^2} F(z)$, since $e^{\pi\bar{z}\zeta}$ is the reproducing kernel of $\mathcal{F}_2(\mathbb{C})$. This gives, for $w(z) = 1$ and $f \in L^\infty(\mathbb{C})$, the following relation between Toeplitz and localization operators in the *weak sense* for any $g, h \in \mathcal{F}_2(\mathbb{C})$:

$$\langle \text{Toep}_f g, h \rangle_{d\mu} = \langle L_f^{(1)} g, h \rangle_{d\mu}.$$

This is equivalent to $L_f^{(1)} g = \text{Toep}_f g$ for any $g \in \mathcal{F}_2(\mathbb{C})$. In case the constant polynomial $e_0(z) = 1$ is replaced by $e_1(z) = \sqrt{\pi}z$ or $e_2(z) = \frac{\pi}{2}z^2$, Coburn's result (cf. [4]) under the change of variable $z \mapsto \sqrt{\pi}z$ gives

$$L_f^{(e_1)} = \text{Toep}_{f + \frac{1}{\pi}\partial_z \partial_{\bar{z}} f}, L_f^{(e_2)} = \text{Toep}_{f + \frac{1}{\pi}\partial_z \partial_{\bar{z}} f + 2(\frac{1}{\pi}\partial_z \partial_{\bar{z}})^2 f}.$$

The above relations are satisfied by every symbol $f(z)$ which is either a polynomial in z and \bar{z} or belongs to the algebra $B_a(\mathbb{C})$ of Fourier-Stieltjes transforms with compactly supported measures. Coburn has also shown in [4] that for any *polynomial* $w \in \mathcal{F}_2(\mathbb{C})$ and polynomial f in z and \bar{z} , there exists a unique polynomial differential operator D depending on $\partial_z, \partial_{\bar{z}}$ and w such that

$$(2.1) \quad L_f^{(w)} = \text{Toep}_{D(w)f}.$$

Coburn conjectured that the coefficients of the polynomial $D(w)$ were constant and that (2.1) holds for any $f \in B_a(\mathbb{C})$. This conjecture was proved by Lo in [13] when Toep_f acts on analytic polynomials on \mathbb{C} . Moreover, Lo extended relation (2.1) to the following class of symbols $E(\mathbb{C})$ which includes $\mathbb{C}[z, \bar{z}]$ and $B_a(\mathbb{C})$:

$$E(\mathbb{C}) = \left\{ f \in C^\infty(\mathbb{C}) : \forall k \in \mathbb{N}_0 \exists C, \alpha > 0 \text{ s.t. } |D^k f(z)| \leq C e^{\alpha|z|}, z \in \mathbb{C} \right\}.$$

An alternative proof of (2.1) has been obtained by Engliš [7]. Engliš' approach works for the whole $\mathcal{F}_2(\mathbb{C})$ and analytic polynomial windows w . It also provides an explicit formula for D . Our main result is the extension of the Coburn-Lo-Engliš theorem to the case of a general window $w \in \mathcal{F}_2(\mathbb{C})$. If the window is not

a polynomial, then the differential operator is of infinite order. Recall that the monomials

$$e_k(z) = \left(\frac{\pi^k}{k!}\right)^{\frac{1}{2}} z^k$$

span $\mathcal{F}_p(\mathbb{C})$. Consider also the sequence of polyanalytic orthogonal polynomials defined as

$$(2.2) \quad e_{j,k}(z) = \left(\frac{\pi^j}{j!}\right)^{\frac{1}{2}} e^{\pi|z|^2} (\partial_z)^j \left[e^{-\pi|z|^2} e_k(z) \right].$$

Let $BC^\infty(\mathbb{C})$ be the space of all $C^\infty(\mathbb{C})$ functions whose partial derivatives are bounded. With this notation, our result is the following statement:

Theorem 1. *For any $w \in \mathcal{F}_2(\mathbb{C})$ and for each symbol $f \in BC^\infty(\mathbb{C})$, the operator $D(w) := D(-\partial_{\bar{z}}, -\partial_z)$ satisfying*

$$L_f^{(w)} = \text{Toep}_{D(w)f}$$

is uniquely determined by

$$D(w)f = \sum_{j,k=0}^{\infty} h_{j,k} (-\partial_{\bar{z}})^j (-\partial_z)^k f,$$

where $h_{j,k} = (\pi^{j+k} j!k!)^{-\frac{1}{2}} \langle |w|^2, e_{j,k} \rangle_{L^2(\mathbb{C}, d\mu)}$. Moreover, $D(w)f \in L^\infty(\mathbb{C})$ only if $f \in BC^\infty(\mathbb{C})$.

We first present the results from [7], [15] and [3] that will be used in the proof of Theorem 1.

3. PRELIMINARY RESULTS

3.1. Direct sum decomposition of $\mathcal{L}_p(\mathbb{C})$. We say that a function F belongs to the *true polyanalytic Fock space* $\mathcal{F}_p^{j+1}(\mathbb{C})$ if $\|F\|_{\mathcal{L}_p(\mathbb{C})} < \infty$ and there exists an entire function H such that

$$F(z) = \left(\frac{\pi^j}{j!}\right)^{\frac{1}{2}} e^{\pi|z|^2} (\partial_z)^j \left[e^{-\pi|z|^2} H(z) \right].$$

The following \mathcal{L}_p -extension of Vasilevski's orthogonal decomposition [15] has been proved in [3, Corollary 1]:

$$(3.1) \quad \mathcal{L}_p(\mathbb{C}) = \bigoplus_{j=0}^{\infty} \mathcal{F}_p^j(\mathbb{C}).$$

It has been shown in [3, Proposition 1] that, for fixed j the linear span of the sequence of polynomials $\{e_{j,k} : k \geq 0\}$ is dense in $\mathcal{F}_p^{j+1}(\mathbb{C})$. Combining this statement with (3.1) it follows that every $F \in \mathcal{L}_p(\mathbb{C})$ can be written in the form

$$F(z) = \sum_{j,k=0}^{\infty} h_{j,k} e_{j,k}(z),$$

where $h_{j,k} = \langle F, e_{j,k} \rangle_{d\mu}$ represents the Fourier-Hermite coefficients.

Similar decompositions appeared in the study of middle Hankel operators on Bergman spaces [14].

3.2. **Berezin symbols.** Let T be a bounded linear operator on $\mathcal{F}_2(\mathbb{C})$ and set

$$K_\zeta(z) = K(\zeta, \zeta)^{-\frac{1}{2}} K(z, \zeta).$$

We define the Berezin symbol $\widetilde{T(\zeta)}$ as $\widetilde{T(\zeta)} = \langle T K_\zeta, K_\zeta \rangle_{d\mu}$. In the case of $z = \zeta$, $K(\zeta, \zeta) = e^{\pi|\zeta|^2}$. This gives

$$(3.2) \quad K_\zeta(z) = W_\zeta e_0(z).$$

On the other hand, since $W_z^* = W_{-z}$ is the adjoint of W_z on $\mathcal{L}_2(\mathbb{C})$, it is easy to check the above relations:

$$W_z^* K_\zeta = e^{i\pi\Im(\bar{\zeta}z)} K_{\zeta-z}.$$

The following relations from [7, (17), p. 6] will be used in the proof of the main result.

Proposition 1. *Let T a bounded linear operator on $\mathcal{F}_2(\mathbb{C})$. Then the following statements hold:*

- (1) $|\widetilde{T(\zeta)}| \leq \|T\|$ for each $\zeta \in \mathbb{C}$.
- (2) $\langle TF, K_\zeta \rangle_{d\mu} = e^{-\frac{\pi}{2}|\zeta|^2} (T F)(\zeta)$ for any $F \in \mathcal{F}_2(\mathbb{C})$.
- (3) T is uniquely determined by $\widetilde{T(\zeta)}$.
- (4) For any $z \in \mathbb{C}$ we have $[W_z^* T W_z](\zeta) = \widetilde{T(\zeta + z)}$.

The boundedness of the operator $L_f^{(w)}$ follows from the following proposition.

Proposition 2. *For any $w \in \mathcal{F}_2(\mathbb{C})$ and $f \in L^\infty(\mathbb{C})$, the operator $L_f^{(w)}$ satisfies the boundedness condition:*

$$\|L_f^{(w)}\| \leq \|f\|_{L^\infty(\mathbb{C})} \|w\|_{\mathcal{L}_2(\mathbb{C})}^2.$$

4. PROOF OF THE MAIN RESULT

Let $w \in \mathcal{F}_2(\mathbb{C})$ and consider

$$k(z) = |w(z)|^2 e^{-\pi|z|^2}$$

and

$$h(z) = e^{-\pi|z|^2}.$$

Using the same arguments as in [7, pp. 6-7], in terms of the standard convolution on \mathbb{C} :

$$(g * f)(\zeta) = \int_{\mathbb{C}} g(z) f(\zeta - z) dz,$$

we can see that

$$(4.1) \quad \widetilde{L_f^{(w)}}(\zeta) = (k * f)(\zeta),$$

$$(4.2) \quad \widetilde{\text{Toep}_f}(\zeta) = (h * f)(\zeta).$$

Now, if $w(z) \in \mathcal{F}_2(\mathbb{C})$, then $|w|^2 \in \mathcal{L}_1(\mathbb{C})$. Thus, we can use the results in section 3.1 with $p = 1$ and find Fourier-Hermite coefficients $h_{j,k} = \langle |w|^2, e_{j,k} \rangle_{d\mu}$ such that

$$(4.3) \quad |w(z)|^2 = \sum_{j,k=0}^{\infty} h_{j,k} e_{j,k}(z) = H(z, \bar{z}).$$

Now let $D(\bar{z}, z)$ be the (uniformly convergent on compact sets) series in z and \bar{z} :

$$(4.4) \quad D(\bar{z}, z) = \sum_{j,k=0}^{\infty} (\pi^{j+k} j!k!)^{-\frac{1}{2}} h_{j,k} \bar{z}^j z^k.$$

From Proposition 2 and statements (1), (2) and (4) of Proposition 1, for each $f \in L^\infty(\mathbb{C})$ the Berezin symbols $\widetilde{L_f^{(w)}}(\zeta)$ and moreover, $\widetilde{\text{Toep}_f}(\zeta)$, are invariant under translations. Then, for each $w \in \mathcal{F}_2(\mathbb{C})$, the function $k(z) = |w(z)|^2 e^{-\pi|z|^2}$ in (4.1) is uniquely determined. Thus, from Proposition 1 (3), the proof of $L_f^{(w)} = \text{Toep}_{D(w)f}$ amounts to showing that

$$\widetilde{L_f^{(w)}}(\zeta) = \widetilde{\text{Toep}_{D(w)f}}(\zeta).$$

By (4.1) and (4.2) this consists of verifying, for $|w(z)|^2 = H(z, \bar{z})$, the following identity:

$$H(\cdot, \cdot) e^{-\pi|\cdot|^2} * f(\cdot) = e^{-\pi|\cdot|^2} * D(w)f,$$

or, using the notation (4.4),

$$(4.5) \quad \int_{\mathbb{C}} D(-\partial_{\bar{z}}, -\partial_z) f(\zeta - z) e^{-\pi|z|^2} dz = \int_{\mathbb{C}} f(\zeta - z) H(z, \bar{z}) e^{-\pi|z|^2} dz.$$

Let us prove (4.5). First, a direct computation shows that

$$(-\partial_{\bar{z}})^j (-\partial_z)^k \left(e^{-\pi|z|^2} \right) = (\pi^{j+k} j!k!)^{\frac{1}{2}} e^{-\pi|z|^2} e_{j,k}(z).$$

Then, we make repeated use of Green’s formula on the disk $\mathbf{D}_r = \{z \in \mathbb{C} : |z| \leq r\}$, valid for $f, g \in C^1(\mathbf{D}_r)$:

$$(4.6) \quad \int_{\mathbf{D}_r} f(z) \partial_z \overline{g(z)} dz = - \int_{\mathbf{D}_r} \partial_z f(z) \overline{g(z)} dz + \frac{1}{i} \int_{\delta \mathbf{D}_r} f(z) \overline{g(z)} dz,$$

$$(4.7) \quad \int_{\mathbf{D}_r} f(z) \partial_{\bar{z}} \overline{g(z)} dz = - \int_{\mathbf{D}_r} \partial_{\bar{z}} f(z) \overline{g(z)} dz + \frac{1}{i} \int_{\delta \mathbf{D}_r} f(z) \overline{g(z)} dz,$$

where the line integral over the circle $\delta \mathbf{D}_r$ is oriented counterclockwise. The result is

$$(4.8) \quad \int_{\mathbf{D}_r} (-\partial_{\bar{z}})^k (-\partial_z)^j f(\zeta - z) e^{-\pi|z|^2} dz = (\pi^{j+k} j!k!)^{\frac{1}{2}} \int_{\mathbf{D}_r} f(\zeta - z) e_{j,k}(z) e^{-\pi|z|^2} dz,$$

since the assumption $f \in BC^\infty(\mathbb{C})$ assures that all the line integrals appearing after the application of Green’s formula are zero. Since $D(\bar{z}, z)$ converges uniformly on $|z| \leq r$, we can use (4.4) and (4.8), interchange the sum with the integral, and use (4.3) to get

$$\int_{\mathbf{D}_r} D(-\partial_{\bar{z}}, -\partial_z) f(\zeta - z) e^{-\pi|z|^2} dz = \int_{\mathbf{D}_r} f(\zeta - z) H(z, \bar{z}) e^{-\pi|z|^2} dz.$$

Letting $r \rightarrow \infty$ we obtain (4.5).

Remark 1. Our methods of proof apply also for windows $w(z) \in \mathcal{F}_p(\mathbb{C})$, $p \geq 2$. However, we believe that the investigation of the case $1 \leq p \leq 2$ may be an interesting problem, since it includes the Feichtinger algebra [8] (corresponding to $p = 1$) which is of particular interest in several applications [9].

REFERENCES

- [1] Luís Daniel Abreu, *Sampling and interpolation in Bargmann-Fock spaces of poly-analytic functions*, Appl. Comput. Harmon. Anal. **29** (2010), no. 3, 287–302, DOI 10.1016/j.acha.2009.11.004. MR2672228 (2011d:42076)
- [2] Luis Daniel Abreu and Hans G. Feichtinger, *Function spaces of polyanalytic functions*, Harmonic and complex analysis and its applications, Trends Math., Birkhäuser/Springer, Cham, 2014, pp. 1–38, DOI 10.1007/978-3-319-01806-5_1. MR3203099
- [3] Luís Daniel Abreu and Karlheinz Gröchenig, *Banach Gabor frames with Hermite functions: polyanalytic spaces from the Heisenberg group*, Appl. Anal. **91** (2012), no. 11, 1981–1997, DOI 10.1080/00036811.2011.584186. MR2983996
- [4] L. A. Coburn, *The Bargmann isometry and Gabor-Daubechies wavelet localization operators*, (Bordeaux, 2000), Oper. Theory Adv. Appl., vol. 129, Birkhäuser, Basel, 2001, pp. 169–178. MR1882695 (2003a:47054)
- [5] Elena Cordero and Karlheinz Gröchenig, *Time-frequency analysis of localization operators*, J. Funct. Anal. **205** (2003), no. 1, 107–131, DOI 10.1016/S0022-1236(03)00166-6. MR2020210 (2004j:47100)
- [6] Ingrid Daubechies, *Time-frequency localization operators: a geometric phase space approach*, IEEE Trans. Inform. Theory **34** (1988), no. 4, 605–612, DOI 10.1109/18.9761. MR966733
- [7] Miroslav Engliš, *Toeplitz operators and localization operators*, Trans. Amer. Math. Soc. **361** (2009), no. 2, 1039–1052, DOI 10.1090/S0002-9947-08-04547-9. MR2452833 (2010a:47056)
- [8] Hans G. Feichtinger, *On a new Segal algebra*, Monatsh. Math. **92** (1981), no. 4, 269–289, DOI 10.1007/BF01320058. MR643206 (83a:43002)
- [9] H. G. Feichtinger, G. A. Zimmermann, *A Banach space of test functions for Gabor analysis*. Gabor analysis and algorithms, 123–170, Appl. Numer. Harmon. Anal., Birkhäuser Boston, Boston, MA, 1998.
- [10] H. G. Feichtinger and K. Nowak, *A Szegő-type theorem for Gabor-Toeplitz localization operators*, Michigan Math. J. **49** (2001), no. 1, 13–21, DOI 10.1307/mmj/1008719032. MR1827072 (2003b:47049)
- [11] Gerald B. Folland, *Harmonic analysis in phase space*, Annals of Mathematics Studies, vol. 122, Princeton University Press, Princeton, NJ, 1989. MR983366 (92k:22017)
- [12] A. Haimi, H. Hendenmalm, *The polyanalytic Ginibre ensembles*. preprint arXiv:1106.2975.
- [13] Min-Lin Lo, *The Bargmann transform and windowed Fourier localization*, Integral Equations Operator Theory **57** (2007), no. 3, 397–412, DOI 10.1007/s00020-006-1462-0. MR2307818 (2008b:47049)
- [14] Li Zhong Peng, Richard Rochberg, and Zhi Jian Wu, *Orthogonal polynomials and middle Hankel operators on Bergman spaces*, Studia Math. **102** (1992), no. 1, 57–75. MR1164633 (93b:47054)
- [15] N. L. Vasilevski, *Poly-Fock spaces*, Differential operators and related topics, Vol. I (Odessa, 1997), Oper. Theory Adv. Appl., vol. 117, Birkhäuser, Basel, 2000, pp. 371–386. MR1764974 (2002a:46026)

ACOUSTIC RESEARCH INSTITUTE, AUSTRIAN ACADEMY OF SCIENCES, WOHLLEBENGASSE 12-14, 1040, VIENNA, AUSTRIA – AND – CMUC, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF COIMBRA, COIMBRA, PORTUGAL

E-mail address: daniel@mat.uc.pt

CMUC, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF COIMBRA, COIMBRA, PORTUGAL
Current address: Departamento de Matemática Aplicada, IMECC-Unicamp, CEP 13083-859, Campinas, SP, Brasil

E-mail address: faustino@ime.unicamp.br