TOPOLOGY, INTERSECTIONS AND FLAT MODULES

CARMELO A. FINOCCHIARO AND DARIO SPIRITO

(Communicated by Irena Peeva)

ABSTRACT. It is well known that, in general, multiplication by an ideal I does not commute with the intersection of a family of ideals, but that this fact holds if I is flat and the family is finite. We generalize this result by showing that finite families of ideals can be replaced by compact subspaces of a natural topological space, and that ideals can be replaced by submodules of an epimorphic extension of a base ring. As a particular case, we give a new proof of a conjecture by Glaz and Vasconcelos.

1. INTRODUCTION

Let D be an integral domain with quotient field K . An *overring* of D is a ring between D and K . The set of all overrings of D is denoted by $\text{Over}(D)$, and can be endowed with a natural topology (called the *Zariski topology*) whose basis of open sets consists of the sets of the form

$$\mathcal{B}(x_1, \dots, x_n) := \{T \in \text{Over}(D) : x_1, \dots, x_n \in T\},$$

as $x_1, \dots, x_n$ vary in K . Under this topology, $\text{Over}(D)$ is a compact T_0 space with a unique closed point (D itself) and a generic point (the quotient field K). One of the clues that this topology is the most natural to be put on $\text{Over}(D)$ is that it makes the localization map

$$\begin{aligned} \lambda: \text{Spec}(D) &\longrightarrow \text{Over}(D) \\ P &\longmapsto D_P \end{aligned}$$

a topological inclusion [4, Lemma 2.4].

This topology, whose origins can be traced back to Zariski's study of the space $\text{Zar}(D)$ of the valuation overrings of an integral domain D [25, Chapter 6, §17] (what is now called the *Zariski space* or the *Riemann-Zariski space* of D), has recently been studied in greater detail (see for example [7, 8, 20, 21]). For example, it has been proved that $\text{Over}(D)$ is a *spectral space*, meaning that there is a ring R such that $\text{Spec}(R)$ is homeomorphic to $\text{Over}(D)$ [5, Proposition 3.5]; the same can be proved of several distinguished subspaces of $\text{Over}(D)$, like for example local overrings [7, Corollary 2.14] or integrally closed overrings [5, Proposition 3.6].

The aim of this paper is to prove a simple and very general result (Theorem 3, in the form of Corollary 5) which intertwines the Zariski topology on $\text{Over}(D)$ with the algebraic properties of the overrings, namely the possibility to commute

Received by the editors October 1, 2015 and, in revised form, December 10, 2015.

2010 *Mathematics Subject Classification.* Primary 13A15, 13A18, 13C11.

Key words and phrases. Zariski topology, overrings, flat ideals.

intersections and products in the case of compact spaces of overrings. In this way, we generalize [13, Lemma 1.1] (which deals with locally finite intersections) and [21, Theorem 3.5] (which proves the same for Noetherian collections of integrally closed overrings). As a consequence, we obtain a new proof of the Graz-Vasconcelos conjecture [12, page 340], independent from the one obtained in [23]. Since it poses no additional challenge, we also work in a more general setting, substituting to the extension $D \subseteq K$ any ring extension that is also an epimorphism, and using modules instead of only overrings.

2. RESULTS

Let $A \subseteq B$ be a ring extension; we denote by $\mathcal{F}(B|A)$ the collection of all the A -submodules of B . The set $\mathcal{F}(B|A)$ becomes a T_0 topological space by declaring, as a basis of open sets, the family of the sets of the form $\mathcal{B}(x_1, \dots, x_n) := \{G \in \mathcal{F}(B|A) : x_1, \dots, x_n \in G\}$, for $x_1, \dots, x_n$ varying in B . Note that, since $\mathcal{B}(x_1, \dots, x_n) = \mathcal{B}(x_1) \cap \dots \cap \mathcal{B}(x_n)$, a convenient subbasis for this topology is $\{\mathcal{B}(x) : x \in B\}$. We call this topology the *Zariski topology*, as it generalizes the Zariski topology on $\text{Over}(D)$ defined in the Introduction. Note that, in particular, the set $\mathcal{I}(A)$ of all the integral ideals of A becomes then a subspace of $\mathcal{F}(B|A)$. On the set $\text{Spec}(A)$ of the prime ideals of A , this topology does *not* coincide with the classical Zariski topology, but rather with the so-called *inverse topology* (see [14] and the discussion before Example 2.2 of [22]). This should, however, not cause any confusion; the only place where we will consider $\text{Spec}(R)$ will be Proposition 11.

If X is any topological space and $Y \subseteq X$, we will denote by $\overline{Y}$ the closure of Y in X .

Remark 1. Let $A \subseteq B$ be a ring extension and let $\mathcal{F}(B|A)$ be endowed with the Zariski topology. The following properties hold:

- (1) For any $F, G \in \mathcal{F}(B|A)$, we have $F \in \overline{\{G\}}$ if and only if $F \subseteq G$.
- (2) Any compact nonempty subspace C of $\mathcal{F}(B|A)$ has minimal elements, with respect to the inclusion $\subseteq$. As a matter of fact, by Zorn's lemma it is enough to show that any chain (under inclusion) $\Sigma \subseteq C$ has a lower bound. By (1), the collection of sets $\mathcal{F} := \{\overline{\{F\}} \cap C : F \in \Sigma\}$ is a chain. Thus, in particular, given any finite subset $F_1, \dots, F_n \in \Sigma$, if G is contained in all F_i , then $G \in \bigcap_{i=1}^n \overline{\{F_i\}} \cap C$. This proves that $\mathcal{F}$ is a collection of closed sets of C with the finite intersection property. By compactness, there exists a submodule $F^* \in \overline{\{F\}} \cap C$, for any $F \in \Sigma$, and applying again (1) we see that F^* is a lower bound of Σ in C .

Now let $\phi : A \longrightarrow B$ be a ring homomorphism. Then, ϕ is an *epimorphism* in the category of rings if, for every $\psi_1, \psi_2 : B \longrightarrow C$, the equality $\psi_1 \circ \phi = \psi_2 \circ \phi$ implies that $\psi_1 = \psi_2$. If the inclusion map $A \hookrightarrow B$ is an epimorphism, we will call the ring extension $A \subseteq B$ an *epimorphic extension*.

Examples of epimorphisms are surjective maps and localizations; more generally, a map $\phi : A \longrightarrow B$ such that the induced homomorphism $\phi_{\mathfrak{p}} : A_{\mathfrak{p}} \longrightarrow B_{\mathfrak{p}}$ is surjective for every $\mathfrak{p} \in \text{Spec}(A)$ such that $\phi(\mathfrak{p})B \neq B$ is an epimorphism (maps with this property are called *weakly surjective* [16, Chapter 1, §3]; on extensions, being an epimorphism and being weakly surjective are equivalent conditions [16, Theorem 4.4]). In particular, if D is an integral domain and K is its quotient field, the

ring extension $D \subseteq K$ is an epimorphic extension. On the other hand, if X is an indeterminate over A , then the extension $A \subseteq A[X]$ is not epimorphic: indeed, for every $\alpha \in A$, we can build a ring homomorphism $\psi_\alpha : A[X] \rightarrow A$ by defining $\psi_\alpha(a) := a$ if $a \in A$ and $\psi_\alpha(X) = \alpha$. In this case, we have $\psi_\alpha \neq \psi_\beta$ if $\alpha \neq \beta$, but every $i \circ \psi_\alpha$ is the identity on A .

The first step of our way is the following fact, which is a generalization of [1, Theorem 2].

Proposition 2. *Let $A \subseteq B$ be an epimorphic extension. Let I be a flat A -submodule of B , and let $G_1, \dots, G_n \in \mathcal{F}(B|A)$. Then,*

$$I(G_1 \cap \dots \cap G_n) = IG_1 \cap \dots \cap IG_n.$$

Proof. With a small abuse of notation, for any $F, G \in \mathcal{F}(B|A)$, we will denote by $F \otimes G$ the submodule of $B \otimes B$ generated by the elements $f \otimes g$, as f varies in F and g varies in G . By induction, it suffices to show the statement for $n = 2$. Consider the map

$$\begin{aligned} \lambda : B \otimes_A B &\longrightarrow B \\ b_1 \otimes b_2 &\longmapsto b_1 b_2. \end{aligned}$$

Clearly, if $I, G \in \mathcal{F}(B|A)$, then $\lambda(I \otimes G) = IG$; therefore, by [18, Theorem 7.4]

$$I(G_1 \cap G_2) = \lambda(I \otimes (G_1 \cap G_2)) = \lambda((I \otimes G_1) \cap (I \otimes G_2)).$$

Since $A \subseteq B$ is an epimorphic extension, λ is an isomorphism (indeed, this property actually characterizes epimorphisms [17, Lemma 1.0]); in particular, λ is a bijection, and thus

$$\lambda((I \otimes G_1) \cap (I \otimes G_2)) = \lambda(I \otimes G_1) \cap \lambda(I \otimes G_2) = IG_1 \cap IG_2.$$

This completes the proof. $\square$

Note that this proposition does not hold if $A \subseteq B$ is not an epimorphism: for example, if X is an indeterminate over A , $B = A[X] = I$, $G_1 = A$, $G_2 = XA[X]$, then $G_1 \cap G_2 = (0)$ and so $I(G_1 \cap G_2) = (0)$, while $IG_1 \cap IG_2 = A[X] \cap XA[X] = XA[X]$.

Theorem 3. *Let $A \subseteq B$ be an epimorphic extension, let I be a flat A -submodule of B and let Y be a (nonempty) compact subspace of $\mathcal{F}(B|A)$. Then, the following equality holds:*

$$I \left(\bigcap_{J \in Y} J \right) = \bigcap_{J \in Y} IJ.$$

Proof. The $(\subseteq)$ containment is obvious. Now take an element $x \in \bigcap \{IJ : J \in Y\}$. For any $J \in Y$, by definition, there exist a positive integer n_J and elements $i_1^{(J)}, \dots, i_{n_J}^{(J)} \in I$, $t_1^{(J)}, \dots, t_{n_J}^{(J)} \in J$ such that

$$x = i_1^{(J)} t_1^{(J)} + \dots + i_{n_J}^{(J)} t_{n_J}^{(J)} = \sum_{h=1}^{n_J} i_h^{(J)} t_h^{(J)}.$$

Consider the open neighborhood $\Omega_J := \mathcal{B}(\{t_1^{(J)}, \dots, t_{n_J}^{(J)}\})$ of J . Then the collection of sets $\mathcal{A} := \{\Omega_J : J \in Y\}$ is an open cover of Y . By compactness, $\mathcal{A}$ admits a finite

subcover, say $\{\Omega_{J_1}, \dots, \Omega_{J_r}\}$, for suitable $J_1, \dots, J_r \in Y$. For any $l = 1, \dots, r$, set $Y_l := \Omega_{J_l} \cap Y$. By Proposition 2, we have

$$I \left(\bigcap_{J \in Y} J \right) = I \left(\bigcap_{J \in Y_1} J \cap \dots \cap \bigcap_{J \in Y_r} J \right) = I \left(\bigcap_{J \in Y_1} J \right) \cap \dots \cap I \left(\bigcap_{J \in Y_r} J \right),$$

and thus it suffices to show that $x \in I \left(\bigcap_{J \in Y_l} J \right)$, for each $l = 1, \dots, r$. However, the elements $t_1^{(J_l)}, \dots, t_{n_{J_l}}^{(J_l)}$ belong to J for every $J \in Y_l$, and thus they belong to the intersection $\bigcap \{J : J \in Y_l\}$; hence, the representation $x = \sum_{h=1}^{n_{J_l}} i_h^{(J_l)} t_h^{(J_l)}$ shows that $x \in I \left(\bigcap_{J \in Y_l} J \right)$. $\square$

Before giving some corollaries of independent interest, we state the following useful lemma.

Lemma 4. *Let $A \subseteq B$ be a ring extension and let $\mathcal{F}(B|A)$ be endowed with the Zariski topology. Fix a submodule $I \in \mathcal{F}(B|A)$. Then, the maps*

$$\begin{array}{ccc} s_I: \mathcal{F}(B|A) \longrightarrow \mathcal{F}(B|A) & & m_I: \mathcal{F}(B|A) \longrightarrow \mathcal{F}(B|A) \\ J \longmapsto I + J & \text{and} & I \longmapsto IJ \end{array}$$

are continuous.

Proof. Let $\mathcal{B}(x)$ be a subbasic open set of $\mathcal{F}(B|A)$, with $x \in B$. If $J_0 \in s_I^{-1}(\mathcal{B}(x))$, then $x = i + j$ for some $i \in I, j \in J_0$; therefore, $\mathcal{B}(j)$ is an open neighborhood of J_0 contained in $s_I^{-1}(\mathcal{B}(x))$, and thus s_I is continuous. Similarly, if $J_0 \in m_I^{-1}(\mathcal{B}(x))$, then $x = i_1 j_1 + \dots + i_n j_n$ for some $j_1, \dots, j_n \in J_0$ and $i_1, \dots, i_n \in I$. Then, $J_0 \in \mathcal{B}(j_1, \dots, j_n) \subseteq m_I^{-1}(\mathcal{B}(x))$, and this shows that m_I is continuous. $\square$

Note that the continuity of m_I makes it possible to shorten the proof of [21, Lemma 3.7].

Corollary 5. *Let D be an integral domain, let I and T be D -submodules of the quotient field K of D , and let Δ be a compact subset of $\text{Over}(D)$, with respect to the Zariski topology. If T is flat over D , then*

$$\left(\bigcap_{U \in \Delta} IU \right) T = \bigcap_{U \in \Delta} (IUT).$$

Proof. By Lemma 4, the collection $\{IU : U \in \Delta\}$ is compact, since it is the continuous image of Δ via m_I . The conclusion is now an immediate consequence of Theorem 3. $\square$

As a particular case of the main results, we now provide a new topological proof of the Glaz-Vasconcelos conjecture.

Corollary 6 ([23, Theorem 1.7]). *Let D be an integrally closed integral domain, and let I be a D -submodule of its quotient field K . If I is flat over D , then $I = \bigcap \{IV : V \in \text{Zar}(D)\}$.*

Proof. The space $\text{Zar}(D)$ is compact in the Zariski topology [25, Chapter 6, Theorem 40]; moreover, since D is integrally closed, $D = \bigcap \{V : V \in \text{Zar}(D)\}$ [3, Corollary 5.22]. Hence, by Theorem 3,

$$I = ID = I \left(\bigcap_{V \in \text{Zar}(D)} V \right) = \bigcap_{V \in \text{Zar}(R)} IV,$$

as claimed. $\square$

Another immediate consequence of the main results deals with intersections of localizations of integral domains.

Corollary 7. *Let D be an integral domain, let Y be a compact nonempty subspace of $\text{Over}(D)$ such that $D = \bigcap \{R : R \in Y\}$, and let S be a multiplicative subset of D . Then, $S^{-1}D = \bigcap \{S^{-1}R : R \in Y\}$.*

Proof. It suffices to use Theorem 3, keeping in mind that $S^{-1}D$ is a flat D -module. $\square$

Corollary 8 ([11, Proposition 43.5]). *Let D be an integral domain, let Y be a locally finite subspace of $\text{Over}(D)$ (i.e., any nonzero element of D is noninvertible only in finitely many members of Y) such that $D = \bigcap \{R : R \in Y\}$, and let S be a multiplicative subset of D . Then, $S^{-1}D = \bigcap \{S^{-1}R : R \in Y\}$.*

Proof. By Corollary 7, it is enough to show that a locally finite collection of overrings of D is compact, with respect to the Zariski topology of $\text{Over}(D)$.

Let $\mathcal{A}$ be an open cover of Y . By Alexander's Subbasis Theorem (see e.g. [15, Chapter 5, Theorem 6, page 139]), we can assume, without loss of generality, that $\mathcal{A}$ consists of subbasic open sets of $\text{Over}(D)$, say $\mathcal{A} = \left\{ \mathcal{B} \left(\frac{a_i}{b_i} \right) : i \in I \right\}$, where $a_i, b_i \in D, b_i \neq 0$, for any $i \in I$. Now fix an index $i' \in I$ and note that, by assumption, the set $Y' := \{R \in Y : b_{i'}^{-1} \notin R\}$ is finite, say $Y' = \{R_1, \dots, R_n\}$. Thus, any member of $Y - Y'$ belongs to $\mathcal{B} \left(\frac{a_{i'}}{b_{i'}} \right)$ and any $R_j \in Y'$ belongs to some $\mathcal{B} \left(\frac{a_{i_j}}{b_{i_j}} \right)$. The proof is now complete. $\square$

Note that the main part of the proof of the previous corollary is also a consequence of [8, Proposition 2.9], where it was proved in the more general context of semistar operations; we inserted the proof here for the reader's convenience. Moreover, the proof of the previous corollary also extends [9, Remark 4.7], where the authors proved that any locally finite family of localizations is compact.

Corollary 9. *Let D be a Prüfer domain with quotient field K , let $\mathfrak{a}$ be an ideal of D , and let $Y \subseteq \mathcal{I}(D)$ be compact. Then,*

$$\mathfrak{a} + \bigcap_{\mathfrak{b} \in Y} \mathfrak{b} = \bigcap_{\mathfrak{b} \in Y} (\mathfrak{a} + \mathfrak{b}).$$

Proof. It suffices to prove that, for every prime ideal $\mathfrak{p}$, the equality

$$\mathfrak{a}D_{\mathfrak{p}} + \left(\bigcap_{\mathfrak{b} \in Y} \mathfrak{b} \right) D_{\mathfrak{p}} = \left(\bigcap_{\mathfrak{b} \in Y} (\mathfrak{a} + \mathfrak{b}) \right) D_{\mathfrak{p}}$$

holds. Fix thus a prime ideal $\mathfrak{p}$, and let $V := D_{\mathfrak{p}}$; since D is a Prüfer domain, V is a valuation domain.

Since V is flat over D and $\{\mathfrak{a} + \mathfrak{b} : \mathfrak{b} \in Y\}$ is compact (Lemma 4), we have, by Theorem 3,

$$(1) \quad \left(\bigcap_{\mathfrak{b} \in Y} (\mathfrak{a} + \mathfrak{b}) \right) V = \bigcap_{\mathfrak{b} \in Y} ((\mathfrak{a} + \mathfrak{b})V) = \bigcap_{\mathfrak{b} \in Y} (\mathfrak{a}V + \mathfrak{b}V).$$

Now observe that, since V is a valuation domain, the collection of ideals $Y' := \{\mathfrak{b}V : \mathfrak{b} \in Y\}$ is totally ordered and compact, by Lemma 4. Thus, since by compactness Y' has minimal elements under inclusion (Remark 1), it follows that Y' has a minimum. Then, there is an ideal $\mathfrak{b}_0 \in Y$ such that $\mathfrak{b}_0V \subseteq \mathfrak{b}V$, for any $\mathfrak{b} \in Y$. It follows that the last member of the equality (1) becomes

$$\bigcap_{\mathfrak{b} \in Y} (\mathfrak{a}V + \mathfrak{b}V) = \mathfrak{a}V + \mathfrak{b}_0V = \mathfrak{a}V + \left(\bigcap_{\mathfrak{b} \in Y} \mathfrak{b}V \right) = \mathfrak{a}V + \left(\bigcap_{\mathfrak{b} \in Y} \mathfrak{b} \right) V,$$

where the last equality is again a consequence of Theorem 3. The proof is now complete. $\square$

Remark 10. The previous corollary is closely related to the dual AB-5* of Grothendieck AB-5 (see, for example, [2]). Precisely, if D is a Prüfer domain and any filter base of ideals of D is compact, with respect to the Zariski topology of $\mathcal{I}(D)$, then D is AB-5* (as D -module).

In the case of Prüfer domains, we can also prove a partial converse of Theorem 3. Recall that a prime ideal P of a Prüfer domain D is *branched* if the set of prime ideals of D properly contained in P has a maximum (see e.g. [11, Theorem 17.3]). If the dimension of D is finite, every prime ideal is branched.

Proposition 11. *Let D be a Prüfer domain with quotient field K , and let $\Delta \subseteq \text{Spec}(D)$ be a nonempty set.*

(a) *Δ is compact (in the “classical” Zariski topology of $\text{Spec}(D)$) if and only*

$$\text{if, for every flat } D\text{-submodule } I \text{ of } K, \bigcap_{\mathfrak{p} \in \Delta} ID_{\mathfrak{p}} = I \left(\bigcap_{\mathfrak{p} \in \Delta} D_{\mathfrak{p}} \right).$$

(b) *Suppose that every prime ideal of D is branched. Then, Δ is compact (in the “classical” Zariski topology of $\text{Spec}(R)$), if and only if $\bigcap_{\mathfrak{p} \in \Delta} D_{\mathfrak{q}}D_{\mathfrak{p}} =$*

$$D_{\mathfrak{q}} \left(\bigcap_{\mathfrak{p} \in \Delta} D_{\mathfrak{p}} \right) \text{ for every } \mathfrak{q} \in \text{Spec}(D).$$

Proof. In both points, one implication follows from Corollary 7 and the fact that the map $\lambda : \text{Spec}(D) \rightarrow \text{Over}(D)$, $P \mapsto D_P$, is a topological inclusion. Suppose Δ is not compact, and let $T := \bigcap_{\mathfrak{p} \in \Delta} D_{\mathfrak{p}}$; note that, without loss of generality, we can suppose that $\Delta = \Delta^{\downarrow} = \{\mathfrak{q} \in \text{Spec}(D) : \mathfrak{q} \subseteq \mathfrak{p} \text{ for some } \mathfrak{p} \in \Delta\}$, since Δ is compact if and only if $\Delta^{\downarrow}$ is compact.

The set of prime ideals $\mathfrak{p}$ such that $\mathfrak{p}T \neq T$ is the image of $\text{Spec}(T)$ under the canonical map $\text{Spec}(T) \rightarrow \text{Spec}(D)$; since it contains Δ , and Δ is not compact,

it must also contain a prime ideal $\mathfrak{q} \notin \Delta$. Since D is a Prüfer domain, $\mathfrak{q}$ is a flat D -module; however,

$$\bigcap_{\mathfrak{p} \in \Delta} \mathfrak{q} D_{\mathfrak{p}} = \bigcap_{\mathfrak{p} \in \Delta} D_{\mathfrak{p}} = T \neq \mathfrak{q} T = \mathfrak{q} \left(\bigcap_{\mathfrak{p} \in \Delta} D_{\mathfrak{p}} \right),$$

against the hypothesis. Therefore, part (a) is proved.

If every prime ideal of D is branched, so is $\mathfrak{q}$; therefore, there is a prime ideal $\mathfrak{q}_0$ directly below $\mathfrak{q}$. No ideal $\mathfrak{p} \in \Delta$ contains $\mathfrak{q}$; therefore, $D_{\mathfrak{p}} D_{\mathfrak{q}} \supsetneq D_{\mathfrak{q}}$, and in particular $D_{\mathfrak{q}_0} \subseteq D_{\mathfrak{p}} D_{\mathfrak{q}}$. Hence,

$$\bigcap_{\mathfrak{p} \in \Delta} D_{\mathfrak{q}} D_{\mathfrak{p}} \supseteq D_{\mathfrak{q}_0} \supsetneq D_{\mathfrak{q}} = D_{\mathfrak{q}} T = D_{\mathfrak{q}} \left(\bigcap_{\mathfrak{p} \in \Delta} D_{\mathfrak{p}} \right),$$

against the hypothesis. Part (b) is proved. $\square$

Note that part (b) of the previous proposition does not hold without the hypothesis that the prime ideals are branched: indeed, if V is a valuation domain with maximal ideal $\mathfrak{m}$ unbranched, and $\Delta := \operatorname{Spec}(V) \setminus \{\mathfrak{m}\}$, then

$$V_{\mathfrak{m}} V = V = \bigcap_{\mathfrak{p} \in \Delta} V_{\mathfrak{p}} = \bigcap_{\mathfrak{p} \in \Delta} V_{\mathfrak{p}} V_{\mathfrak{m}},$$

despite Δ not being compact.

Another question arising from Theorem 3 is if the equality $I(\bigcap_{J \in Y} J) = \bigcap_{J \in Y} IJ$, for all compact families Y of submodules of an epimorphic extension $A \subseteq B$, implies that I is flat. This is true if the base ring A is a domain, but fails in general (see [1, Theorem 2] and the subsequent discussion).

Remark 12. While Theorem 3 is quite general, it may be in general hard, or at least not easy, to find examples of compact subspaces to which it can be applied, or to prove that a given family is actually compact.

Some examples can be constructed using the fact that, under the Zariski topology, $\mathcal{F}(B|A)$ is a spectral space, i.e., it is homeomorphic to the prime spectrum of a ring [22, Example 2.2(2)]. For example, it follows from Remark 1(1) and either [24, Proposition 2.3] or [19, Proposition 2.2] that a subset Y of $\mathcal{F}(B|A)$ is compact if and only if every element of the closure of Y , with respect to the constructible topology, contains a point of Y , where the *constructible topology* on $\mathcal{F}(B|A)$ is the coarsest topology on $\mathcal{F}(B|A)$ for which any open and compact subspace of $\mathcal{F}(B|A)$ is both open and closed.

Another class of examples comes from the domination map $d : \operatorname{Zar}(A) \rightarrow \operatorname{Spec}(A)$ of the Zariski space of a domain A (i.e., the set of valuation overrings of A). For example, if S is a compact subspace of $\operatorname{Spec}(A)$, then $d^{-1}(S)$ is compact, by [19, Proposition 2.2 and Lemma 2.7(3)].

ACKNOWLEDGMENT

The authors would like to thank the referee for helpful comments and suggestions.

REFERENCES

- [1] D. D. Anderson, *On the ideal equation $I(B \cap C) = IB \cap IC$* , Canad. Math. Bull. **26** (1983), no. 3, 331–332, DOI 10.4153/CMB-1983-054-3. MR703406
- [2] Pham Ngoc Anh, Dolores Herbera, and Claudia Menini, *AB-5* and linear compactness*, J. Algebra **200** (1998), no. 1, 99–117, DOI 10.1006/jabr.1997.7216. MR1603265
- [3] M. F. Atiyah and I. G. Macdonald, *Introduction to commutative algebra*, Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills, Ont., 1969. MR0242802
- [4] David E. Dobbs, Richard Fedder, and Marco Fontana, *Abstract Riemann surfaces of integral domains and spectral spaces* (English, with Italian summary), Ann. Mat. Pura Appl. (4) **148** (1987), 101–115, DOI 10.1007/BF01774285. MR932760
- [5] Carmelo Antonio Finocchiaro, *Spectral spaces and ultrafilters*, Comm. Algebra **42** (2014), no. 4, 1496–1508, DOI 10.1080/00927872.2012.741875. MR3169645
- [6] Carmelo A. Finocchiaro, Marco Fontana, and K. Alan Loper, *The constructible topology on spaces of valuation domains*, Trans. Amer. Math. Soc. **365** (2013), no. 12, 6199–6216, DOI 10.1090/S0002-9947-2013-05741-8. MR3105748
- [7] Carmelo Antonio Finocchiaro, Marco Fontana, and Dario Spirito, *New distinguished classes of spectral spaces: A survey*, Proceedings of the Graz conference, to appear.
- [8] Carmelo Antonio Finocchiaro and Dario Spirito, *Some topological considerations on semistar operations*, J. Algebra **409** (2014), 199–218, DOI 10.1016/j.jalgebra.2014.04.002. MR3198840
- [9] Marco Fontana and James A. Huckaba, *Localizing systems and semistar operations*, Non-Noetherian commutative ring theory, Math. Appl., vol. 520, Kluwer Acad. Publ., Dordrecht, 2000, pp. 169–197. MR1858162
- [10] Marco Fontana, James A. Huckaba, and Ira J. Papick, *Prüfer domains*, Monographs and Textbooks in Pure and Applied Mathematics, vol. 203, Marcel Dekker, Inc., New York, 1997. MR1413297
- [11] Robert Gilmer, *Multiplicative ideal theory*, Queen’s Papers in Pure and Applied Mathematics, vol. 90, Queen’s University, Kingston, ON, 1992. Corrected reprint of the 1972 edition. MR1204267
- [12] Sarah Glaz and Wolmer V. Vasconcelos, *Flat ideals. II*, Manuscripta Math. **22** (1977), no. 4, 325–341. MR0472797
- [13] William Heinzer and Jack Ohm, *Noetherian intersections of integral domains*, Trans. Amer. Math. Soc. **167** (1972), 291–308. MR0296095
- [14] M. Hochster, *Prime ideal structure in commutative rings*, Trans. Amer. Math. Soc. **142** (1969), 43–60. MR0251026
- [15] John L. Kelley, *General topology*, Springer-Verlag, New York-Berlin, 1975. Reprint of the 1955 edition [Van Nostrand, Toronto, Ont.]; Graduate Texts in Mathematics, No. 27. MR0370454
- [16] Manfred Knebusch and Digen Zhang, *Manis valuations and Prüfer extensions. I*, Lecture Notes in Mathematics, vol. 1791, Springer-Verlag, Berlin, 2002. A new chapter in commutative algebra. MR1937245
- [17] Daniel Lazard, *Autour de la platitude* (French), Bull. Soc. Math. France **97** (1969), 81–128. MR0254100
- [18] Hideyuki Matsumura, *Commutative ring theory*, Cambridge Studies in Advanced Mathematics, vol. 8, Cambridge University Press, Cambridge, 1986. Translated from the Japanese by M. Reid. MR879273
- [19] Bruce Olberding, *Affine schemes and topological closures in the Zariski-Riemann space of valuation rings*, J. Pure Appl. Algebra **219** (2015), no. 5, 1720–1741, DOI 10.1016/j.jpaa.2014.07.009. MR3299704
- [20] Bruce Olberding, *Overrings of two-dimensional Noetherian domains representable by Noetherian spaces of valuation rings*, J. Pure Appl. Algebra **212** (2008), no. 7, 1797–1821, DOI 10.1016/j.jpaa.2007.11.008. MR2400744
- [21] Bruce Olberding, *Noetherian spaces of integrally closed rings with an application to intersections of valuation rings*, Comm. Algebra **38** (2010), no. 9, 3318–3332, DOI 10.1080/00927870903114979. MR2724221
- [22] Bruce Olberding, *Topological aspects of irredundant intersections of ideals and valuation rings*, Proceedings of the Graz conference, to appear.
- [23] Giampaolo Picozza and Francesca Tartarone, *Flat ideals and stability in integral domains*, J. Algebra **324** (2010), no. 8, 1790–1802, DOI 10.1016/j.jalgebra.2010.07.021. MR2678822

- [24] Niels Schwartz and Marcus Tressl, *Elementary properties of minimal and maximal points in Zariski spectra*, J. Algebra **323** (2010), no. 3, 698–728, DOI 10.1016/j.jalgebra.2009.11.003. MR2574858
- [25] Oscar Zariski and Pierre Samuel, *Commutative algebra. Vol. II*, Springer-Verlag, New York-Heidelberg, 1975. Reprint of the 1960 edition; Graduate Texts in Mathematics, Vol. 29. MR0389876

DIPARTIMENTO DI MATEMATICA E FISICA, UNIVERSITÀ DEGLI STUDI “ROMA TRE”, LARGO SAN LEONARDO MURIALDO, 1, 00146 ROMA, ITALY

E-mail address: `carmelo@mat.uniroma3.it`

Current address: Institute of Analysis and Number Theory, Graz University of Technology, 8010 Graz, Steyrergasse 31/II, Austria

E-mail address: `finocchiaro@math.tugraz.at`

DIPARTIMENTO DI MATEMATICA E FISICA, UNIVERSITÀ DEGLI STUDI “ROMA TRE”, LARGO SAN LEONARDO MURIALDO, 1, 00146 ROMA, ITALY

E-mail address: `spirito@mat.uniroma3.it`