

RELATIVE ENERGY GAP FOR HARMONIC MAPS OF RIEMANN SURFACES INTO REAL ANALYTIC RIEMANNIAN MANIFOLDS

PAUL M. N. FEEHAN

(Communicated by Lei Ni)

ABSTRACT. We extend the well-known Sacks–Uhlenbeck energy gap result for harmonic maps from closed Riemann surfaces into closed Riemannian manifolds from the case of maps with small energy (thus near a constant map), to the case of harmonic maps with high absolute energy but small energy relative to a reference harmonic map.

1. INTRODUCTION

Let (M, g) and (N, h) be a pair of smooth Riemannian manifolds, with M orientable. One defines the *harmonic map energy function* by

$$(1.1) \quad \mathcal{E}_{g,h}(f) := \frac{1}{2} \int_M |df|_{g,h}^2 d\text{vol}_g,$$

for smooth maps, $f : M \rightarrow N$, where $df : TM \rightarrow TN$ is the differential map. A map $f \in C^\infty(M; N)$ is (*weakly*) *harmonic* if it is a *critical point* of $\mathcal{E}_{g,h}$, so $\mathcal{E}'_{g,h}(f)(u) = 0$ for all $u \in C^\infty(M; f^*TN)$, where

$$\mathcal{E}'_{g,h}(f)(u) = \int_M \langle df, u \rangle_{g,h} d\text{vol}_g.$$

The purpose of this article is to prove

Theorem 1 (Relative energy gap for harmonic maps of Riemann surfaces into real analytic Riemannian manifolds). *Let (M, g) be a closed Riemann surface and let (N, h) be a closed, real analytic Riemannian manifold equipped with a real analytic isometric embedding into a Euclidean space, \mathbb{R}^n . If $f_\infty \in C^\infty(M; N)$ is a harmonic map, then there is a constant $\epsilon = \epsilon(f_\infty, g, h) \in (0, 1]$ with the following significance. If $f \in C^\infty(M; N)$ is a harmonic map obeying*

$$(1.2) \quad \|d(f - f_\infty)\|_{L^2(M; \mathbb{R}^n)} + \|f - f_\infty\|_{L^2(M; \mathbb{R}^n)} < \epsilon,$$

then $\mathcal{E}_{g,h}(f) = \mathcal{E}_{g,h}(f_\infty)$.

Received by the editors September 12, 2017.

2010 *Mathematics Subject Classification*. Primary 58E20; Secondary 37D15.

Key words and phrases. Harmonic maps, Łojasiewicz–Simon gradient inequality.

The author was partially supported by National Science Foundation grant DMS-1510064 and the Oswald Veblen Fund and Fund for Mathematics (Institute for Advanced Study, Princeton) during the preparation of this article.

Remark 1.1 (Generalizations to the case of harmonic maps with potentials). In physics, harmonic maps arise in the context of non-linear sigma models and with such applications in mind, Theorem 1 should admit generalizations to allow, for example, the addition to $\mathcal{E}_{g,h}$ of a real analytic potential function, $V : C^\infty(M; N) \rightarrow \mathbb{R}$, in the definition (1.1) of the energy, as explored by Branding [6].

Naturally, Theorem 1 continues to hold if the condition (1.2) is replaced by the stronger (and conformally invariant) hypothesis,

$$\|d(f - f_\infty)\|_{L^2(M; \mathbb{R}^n)} + \|f - f_\infty\|_{L^\infty(M; \mathbb{R}^n)} < \epsilon.$$

Thus, if φ is any conformal diffeomorphism of (M, g) , then the preceding condition on f, f_∞ holds if and only if the harmonic maps $f \circ \varphi, f_\infty \circ \varphi$ obey

$$\|d(f \circ \varphi - f_\infty \circ \varphi)\|_{L^2(M; \mathbb{R}^n)} + \|f \circ \varphi - f_\infty \circ \varphi\|_{L^\infty(M; \mathbb{R}^n)} < \epsilon.$$

Hence, the constants, Z, σ, θ in Theorem 1 are in this sense independent of the action of the conformal group of (M, g) on harmonic maps from M to N .

Theorem 1 may be viewed, in part, as a generalization of the following energy gap result due to Sacks and Uhlenbeck and who do not require that the target manifold be real analytic.

Theorem 1.2 (Energy gap near the constant map). [34, Theorem 3.3] *Let (M, g) be a closed Riemann surface and (N, h) be a closed, smooth Riemannian manifold. Then there is a constant, $\epsilon > 0$, such that if $f \in C^\infty(M; N)$ is harmonic and $\mathcal{E}_{g,h}(f) < \epsilon$, then f is a constant map and $\mathcal{E}_{g,h}(f) = 0$.*

The Sacks–Uhlenbeck Energy Gap Theorem 1.2 has been generalized by Branding [5, Lemma 4.9] and by Jost and his collaborators [9, Proposition 4.2], [25, Proposition 5.2] to the case of Dirac-harmonic pairs. Theorem 1.2 ensures positivity of the constant \hbar in the

Definition 1.3 (Dirac–Planck constant). Let (N, h) be a closed, smooth Riemannian manifold. Then \hbar denotes the *least energy* of a non-constant C^∞ map from (S^2, g_{round}) into (N, h) , where g_{round} is the standard round metric of radius one on S^2 .

The energy gap near the ‘ground state’ characterized by the constant maps from (S^2, g_{round}) to (N, h) appears to be unusual in light of the following counter-example due to Li and Wang [27] when (N, h) is only C^∞ rather than real analytic.

Example 1.4 (Non-discreteness of the energy spectrum for harmonic maps from S^2 into a smooth Riemannian manifold with boundary). (See [27, Section 4].) There exists a smooth Riemannian metric h on $N = S^2 \times (-1, 1)$ such that the energies of harmonic maps from (S^2, g_{round}) to (N, h) have an accumulation point at the energy level 4π , where, g_{round} denotes the standard round metric of radius one.

Thus we would *not* expect Theorem 1 to hold when the hypothesis that (N, h) is real analytic is omitted, except for the case where f_∞ is a constant map. On the other hand, when (N, h) is real analytic, one has the following conjecture due to Lin [28].

Conjecture 1.5 (Discreteness for energies of harmonic maps from closed Riemann surfaces into analytic closed Riemannian manifolds). (*Lin* [28, Conjecture 5.7].) *Assume the hypotheses of Theorem 1 and that (M, g) is the two-sphere, S^2 , with its*

standard, round metric. Then the subset of critical values of the energy function, $\mathcal{E}_{g,h} : C^\infty(S^2; N) \rightarrow [0, \infty)$, is closed and discrete.

One may therefore view Theorem 1 as supporting evidence of the validity of Conjecture 1.5. In the special case that N is the Lie group $U(n)$ with $n \geq 2$ and its standard Riemannian metric, Valli [48, Corollary 8] has shown (using ideas of Uhlenbeck [46]) that the energies of harmonic maps from (S^2, g_{round}) into $U(n)$ are integral multiples of 8π . If (N, h) has non-positive curvature sectional curvature, then Adachi and Sunada [1, Theorem 1] have shown that Conjecture 1.5 holds when (M, g) is any closed Riemann surface.

1.1. Outline of the article. In Section 2, we review the Lojasiewicz–Simon gradient inequality for the harmonic map energy function based on results of the author and Maridakis [12] and Simon [38, 39]. In Section 3, we prove certain a priori estimates for the difference of two harmonic maps and, with the aid of the Lojasiewicz–Simon gradient inequality, complete the proof of Theorem 1.

2. LOJASIEWICZ–SIMON GRADIENT INEQUALITY FOR THE HARMONIC MAP ENERGY FUNCTION

In this section, we closely follow our treatment of the Lojasiewicz–Simon gradient inequality for abstract and harmonic map energy functions provided by the author and Maridakis in [12, Sections 1.1, 1.2, and 1.4]. Useful references for harmonic maps include Eells and Lemaire [10, 11], Hamilton [16], Hélein [17], Hélein and Wood [18], Jost [20–24], Moser [30], Parker [33], Sacks and Uhlenbeck [34, 35], Schoen [36], Simon [40], Struwe [42], Urakawa [47], and Xin [49], and the citations contained therein.

When clear from the context, we omit explicit mention of the Riemannian metrics g on M and h on N and write $\mathcal{E} = \mathcal{E}_{g,h}$. Although initially defined for smooth maps, the energy function \mathcal{E} in (1.1), extends to the case of Sobolev maps of class $W^{1,2}$. To define the gradient, $\mathcal{M} = \mathcal{M}_{g,h}$, of the energy function \mathcal{E} in (1.1) with respect to the L^2 metric on $C^\infty(M; N)$, we first choose an isometric embedding, $(N, h) \subset \mathbb{R}^n$ for a sufficiently large n (courtesy of the isometric embedding theorem due to Nash [31]), and recall that¹ by [40, Equations (2.2)(i) and (ii)]

$$\begin{aligned} (u, \mathcal{M}(f))_{L^2(M,g)} &:= \mathcal{E}'(f)(u) = \left. \frac{d}{dt} \mathcal{E}(\pi(f + tu)) \right|_{t=0} \\ &= (u, \Delta_g f)_{L^2(M,g)} \\ &= (u, d\pi_h(f) \Delta_g f)_{L^2(M,g)} \end{aligned}$$

for all $u \in C^\infty(M; f^*TN)$, where π_h is the nearest point projection onto N from a normal tubular neighborhood and $d\pi_h(y) : \mathbb{R}^n \rightarrow T_y N$ is an orthogonal projection for all $y \in N$. By [17, Lemma 1.2.4], we have

$$(2.1) \quad \mathcal{M}(f) = d\pi_h(f) \Delta_g f = \Delta_g f - A_h(f)(df, df),$$

as in [40, Equations (2.2)(iii) and (iv)]. Here, A_h denotes the second fundamental form of the isometric embedding, $(N, h) \subset \mathbb{R}^n$, and

$$(2.2) \quad \Delta_g := -\operatorname{div}_g \operatorname{grad}_g = d^{*,g} d = -\frac{1}{\sqrt{\det g}} \frac{\partial}{\partial x^\beta} \left(\sqrt{\det g} \frac{\partial f}{\partial x^\alpha} \right)$$

¹Compare [24, Equations (8.1.10) and (8.1.13)], where Jost uses variations of f of the form $\exp_f(tu)$.

denotes the Laplace–Beltrami operator for (M, g) (with the opposite sign convention to that of [8, Equations (1.14) and (1.33)]) acting on the scalar components f^i of $f = (f^1, \dots, f^n)$ and the $\{x^\alpha\}$ denote local coordinates on M . As usual, the gradient vector field, $\text{grad}_g f^i \in C^\infty(TM)$, is defined by $\langle \text{grad}_g f^i, \xi \rangle_g := df^i(\xi)$ for all $\xi \in C^\infty(TM)$ and $1 \leq i \leq n$ and the divergence function, $\text{div}_g \xi \in C^\infty(M; \mathbb{R})$, by the pointwise trace, $\text{div}_g \xi := \text{tr}(\eta \mapsto \nabla_\xi^g \eta)$, for all $\eta \in C^\infty(TM)$.

Given a smooth map, $f : M \rightarrow N$, an isometric embedding, $(N, h) \subset \mathbb{R}^n$, a non-negative integer k , and constant $p \in [1, \infty)$, we define the Sobolev norms,

$$\|f\|_{W^{k,p}(M; \mathbb{R}^n)} := \left(\sum_{i=1}^n \|f^i\|_{W^{k,p}(M; \mathbb{R}^n)}^p \right)^{1/p},$$

with

$$\|f^i\|_{W^{k,p}(M; \mathbb{R}^n)} := \left(\sum_{j=0}^k \int_M |(\nabla^g)^j f^i|^p d\text{vol}_g \right)^{1/p},$$

where ∇^g denotes the Levi–Civita connection on TM and all associated bundles (that is, T^*M and their tensor products), and if $p = \infty$, we define

$$\|f\|_{W^{k,\infty}(M; \mathbb{R}^n)} := \sum_{i=1}^n \sum_{j=0}^k \text{ess sup}_M |(\nabla^g)^j f^i|.$$

If $k = 0$, then we denote $\|f\|_{W^{0,p}(M; \mathbb{R}^n)} = \|f\|_{L^p(M; \mathbb{R}^n)}$. For $p \in [1, \infty)$ and non-negative integers k , we use [2, Theorem 3.12] (applied to $W^{k,p}(M; \mathbb{R}^n)$ and noting that M is a closed manifold) and Banach space duality to define

$$W^{-k,p'}(M; \mathbb{R}^n) := (W^{k,p}(M; \mathbb{R}^n))^*,$$

where $p' \in (1, \infty]$ is the dual exponent defined by $1/p + 1/p' = 1$. Elements of the continuous Banach dual space, $(W^{k,p}(M; \mathbb{R}^n))^*$, may be characterized via [2, Section 3.10] as distributions in the Schwartz space, $\mathcal{D}'(M; \mathbb{R}^n)$ [2, Section 1.57].

Spaces of Hölder continuous maps, $C^{k,\lambda}(M; N)$ for $\lambda \in (0, 1)$ and integers $k \geq 0$, and norms,

$$\|f\|_{C^{k,\lambda}(M; \mathbb{R}^n)},$$

may be defined as in [2, Section 1.29].

We note that if (N, h) is real analytic, then the isometric embedding, $(N, h) \subset \mathbb{R}^n$, may also be chosen to be analytic by the analytic isometric embedding theorem due to Nash [32], with a simplified proof due to Greene and Jacobowitz [15]).

Definition 2.1 (Harmonic map). (See [17, Definition 1.4.9].) A map $f \in W^{1,2}(M; N)$ is called *weakly harmonic* if it is a critical point of the L^2 -energy functional (1.1), that is,

$$\mathcal{E}'(f)(u) = 0 \quad \forall u \in C^\infty(M; f^*TN),$$

and a map $f \in W^{2,p}(M; N)$, for $p \in [1, \infty]$, is called *harmonic* if

$$(2.3) \quad \Delta_g f - A_h(df, df) = 0 \quad \text{a.e. on } M.$$

A well-known result due to Hélein [17, Theorem 4.1.1] tells us that if M has dimension $d = 2$, then $f \in C^\infty(M; N)$; for $d \geq 3$, regularity results are far more limited — see, for example, [17, Theorem 4.3.1] due to Bethuel. From [12], we recall the following theorem.

Theorem 2.2 (Lojasiewicz–Simon gradient inequality for the energy function for maps between pairs of Riemannian manifolds). (See [12, Theorem 5].) *Let $d \geq 2$ and $k \geq 1$ be integers and $p \in (1, \infty)$ be such that $kp > d$. Let (M, g) and (N, h) be closed, smooth Riemannian manifolds, with M of dimension d . If (N, h) is real analytic (respectively, C^∞) and $f \in W^{k,p}(M; N)$, then the gradient map \mathcal{M} in (2.1) for the energy function, $\mathcal{E} : W^{k,p}(M; N) \rightarrow \mathbb{R}$, in (1.1),*

$$W^{k,p}(M; N) \ni f \mapsto \mathcal{M}(f) \in W^{k-2,p}(M; f^*TN) \subset W^{k-2,p}(M; \mathbb{R}^n),$$

is a real analytic (respectively, C^∞) map of Banach spaces. If (N, h) is real analytic and $f_\infty \in W^{k,p}(M; N)$ is a weakly harmonic map, then there are positive constants $Z \in (0, \infty)$, and $\sigma \in (0, 1]$, and $\theta \in [1/2, 1)$, depending on f_∞, g, h, k, p , with the following significance. If $f \in W^{k,p}(M; N)$ obeys

$$(2.4) \quad \|f - f_\infty\|_{W^{k,p}(M; \mathbb{R}^n)} < \sigma,$$

then the gradient \mathcal{M} in (2.1) of the harmonic map energy function \mathcal{E} in (1.1) obeys

$$(2.5) \quad \|\mathcal{M}(f)\|_{W^{k-2,p}(M; f^*TN)} \geq Z|\mathcal{E}(f) - \mathcal{E}(f_\infty)|^\theta.$$

Remark 2.3 (On the hypotheses of Theorem 2.2). When $k = d$ and $p = 1$, then $W^{d,1}(M; \mathbb{R}) \subset C(M; \mathbb{R})$ is a continuous embedding by [2, Theorem 4.12] and $W^{d,1}(M; \mathbb{R})$ is a Banach algebra by [2, Theorem 4.39]. In particular, $W^{d,1}(M; N)$ is a real analytic Banach manifold by [12, Proposition 3.2] and the harmonic map energy function, $\mathcal{E} : W^{d,1}(M; N) \rightarrow \mathbb{R}$, is real analytic by [12, Proposition 3.5]. However, the operator $\mathcal{M}'(f_\infty) : W^{d,1}(M; f_\infty^*TN) \rightarrow W^{d-2,1}(M; f_\infty^*TN)$ may not be Fredholm.

Theorem 2.2 extends a version of the Lojasiewicz–Simon gradient inequality that is stated by Simon in [39, Equation (4.27)] and can be derived from his more general [38, Theorem 3].

Theorem 2.4 (Lojasiewicz–Simon gradient inequality for the energy function for maps between pairs of Riemannian manifolds). (See [12, Corollary 6], [38, Theorem 3], [39, Equation (4.27)].) *Let $d \geq 2$ and $\lambda \in (0, 1)$ be constants, let (M, g) be a closed, smooth Riemannian manifold of dimension d , and (N, h) is a closed, real analytic Riemannian manifold. If $f_\infty \in C^{2,\lambda}(M; N)$ is a harmonic map, then there are positive constants $Z \in (0, \infty)$, and $\sigma \in (0, 1]$, and $\theta \in [1/2, 1)$, depending on f_∞, g, h, λ , with the following significance. If $f \in C^{2,\lambda}(M; N)$ obeys*

$$(2.6) \quad \|f - f_\infty\|_{C^{2,\lambda}(M; \mathbb{R}^n)} < \sigma,$$

then the gradient \mathcal{M} in (2.1) of the harmonic map energy function \mathcal{E} in (1.1) obeys

$$(2.7) \quad \|\mathcal{M}(f)\|_{L^2(M; f^*TN)} \geq Z|\mathcal{E}(f) - \mathcal{E}(f_\infty)|^\theta.$$

Remark 2.5 (Other versions of the Lojasiewicz–Simon gradient inequality for the harmonic map energy function). Topping [45, Lemma 1] proved a Lojasiewicz-type gradient inequality for maps, $f : S^2 \rightarrow S^2$, with small energy, with the latter criterion replacing the usual small $C^{2,\lambda}(M; \mathbb{R}^n)$ norm criterion of Simon for the difference between a map and a critical point. Topping’s result is generalized by Liu and Yang in [29, Lemma 3.3]. Kwon [26, Theorem 4.2] obtains a Lojasiewicz-type gradient inequality for maps, $f : S^2 \rightarrow N$, that are $W^{2,p}(S^2; \mathbb{R}^n)$ -close to a harmonic map, with $1 < p \leq 2$.

When $d = 2$ in the hypotheses of Theorem 2.2, the reader will note that the two cases that are most directly applicable to a proof of Theorem 1 are omitted, namely the cases $k = 2$ and $p = 1$ or $k = 1$ and $p = 2$, which are both critical since $kp = d$. We shall briefly comment on each of these two cases.

When $d = 2$, $k = 1$, and $p = 2$, it appears very difficult to verify the hypotheses of [12, Theorem 2]. The analytical difficulties are very much akin to those confronted by Hélein [17] in his celebrated proof of smoothness of weakly harmonic maps from Riemann surfaces. However, it is unclear that Hélein's methods could be used to extend Theorem 2.2 to the case $d = 2$, $k = 1$, and $p = 2$.

Similarly, when $d = 2$, $k = 2$, and $p = 1$, it is very difficult to verify the hypotheses of [12, Theorem 2]. One might speculate that a version of Theorem 2.2 could hold if the role of the pair of Sobolev spaces, $W^{2,1}(M; f_\infty^*TN)$ and $L^1(M; f_\infty^*TN)$, were replaced by suitably defined *local Hardy spaces*. We refer the reader to Semmes [37], Stein [41], and Taylor [44] for introductions to Hardy spaces of functions on Euclidean space and to Hélein [17] for their application to the problem of regularity for weakly harmonic maps from Riemann surfaces. Auscher, McIntosh, and Morris [3], Carbonaro, McIntosh, and Morris [7], and Taylor [43] provide definitions of local Hardy spaces on Riemannian manifolds. However, the analytical difficulties appear formidable in any such approach.

Fortunately, in our proof of Theorem 1, we can apply Theorem 2.2 with *non-critical* Sobolev exponents, namely $d = 2$, $k = 2$, and $p \in (1, \infty)$ by exploiting certain a priori estimates for harmonic maps similar to those used by Sacks and Uhlenbeck [34].

Theorem 2.2 is proved by the author and Maridakis in [12] as a consequence of a more general abstract Łojasiewicz–Simon gradient inequality for an analytic function on a Banach space, namely [12, Theorem 2], while Theorem 2.4 may be deduced as a consequence of [12, Theorem 2].

To state the abstract [12, Theorem 2], we let \mathcal{X} be a Banach space and let \mathcal{X}^* denote its continuous dual space. We call a bilinear form, $b : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$, *definite* if $b(x, x) \neq 0$ for all $x \in \mathcal{X} \setminus \{0\}$. We say that a continuous *embedding* of a Banach space into its continuous dual space, $j : \mathcal{X} \rightarrow \mathcal{X}^*$, is *definite* if the pullback of the canonical pairing, $\mathcal{X} \times \mathcal{X} \ni (x, y) \mapsto \langle x, j(y) \rangle_{\mathcal{X} \times \mathcal{X}^*} \rightarrow \mathbb{R}$, is a definite bilinear form. (This hypothesis on the continuous embedding, $\mathcal{X} \subset \mathcal{X}^*$, is easily achieved given a continuous embedding of \mathcal{X} into a Hilbert space \mathcal{H} but the increased generality is often convenient.)

Definition 2.6 (Gradient map). (See [4, Section 2.5], [19, Definition 2.1.1].) Let $\mathcal{U} \subset \mathcal{X}$ be an open subset of a Banach space, \mathcal{X} , and let $\tilde{\mathcal{X}}$ be a Banach space with continuous embedding, $\tilde{\mathcal{X}} \subseteq \mathcal{X}^*$. A continuous map, $\mathcal{M} : \mathcal{U} \rightarrow \tilde{\mathcal{X}}$, is called a *gradient map* if there exists a C^1 function, $\mathcal{E} : \mathcal{U} \rightarrow \mathbb{R}$, such that

$$(2.8) \quad \mathcal{E}'(x)v = \langle v, \mathcal{M}(x) \rangle_{\mathcal{X} \times \mathcal{X}^*} \quad \forall x \in \mathcal{U}, \quad v \in \mathcal{X},$$

where $\langle \cdot, \cdot \rangle_{\mathcal{X} \times \mathcal{X}^*}$ is the canonical bilinear form on $\mathcal{X} \times \mathcal{X}^*$. The real-valued function, \mathcal{E} , is called a *potential* for the gradient map, \mathcal{M} .

Theorem 2.7 (Łojasiewicz–Simon gradient inequality for analytic functions on Banach spaces). (See [12, Corollary 3].) Let \mathcal{X} and $\tilde{\mathcal{X}}$ be Banach spaces with continuous embeddings, $\mathcal{X} \subset \tilde{\mathcal{X}} \subset \mathcal{X}^*$, and such that the embedding, $\mathcal{X} \subset \mathcal{X}^*$, is definite. Let $\mathcal{U} \subset \mathcal{X}$ be an open subset, let $\mathcal{E} : \mathcal{U} \rightarrow \mathbb{R}$ be a C^2 function with real analytic gradient map, $\mathcal{M} : \mathcal{U} \rightarrow \tilde{\mathcal{X}}$, and let $x_\infty \in \mathcal{U}$ be a critical point of

\mathcal{E} , that is, $\mathcal{M}(x_\infty) = 0$. If $\mathcal{M}'(x_\infty) : \mathcal{X} \rightarrow \tilde{\mathcal{X}}$ is a Fredholm operator with index zero, then there are constants, $Z \in (0, \infty)$, and $\sigma \in (0, 1]$, and $\theta \in [1/2, 1)$, with the following significance. If $x \in \mathcal{U}$ obeys

$$(2.9) \quad \|x - x_\infty\|_{\mathcal{X}} < \sigma,$$

then

$$(2.10) \quad \|\mathcal{M}(x)\|_{\tilde{\mathcal{X}}} \geq Z|\mathcal{E}(x) - \mathcal{E}(x_\infty)|^\theta.$$

Theorem 2.2 follows from Theorem 2.7 by choosing

$$\mathcal{X} = W^{k,p}(M; f_\infty^*TN) \quad \text{and} \quad \tilde{\mathcal{X}} = W^{k-2,p}(M; f_\infty^*TN).$$

Theorem 2.4 follows from Theorem 2.2 when one can choose $k \geq 1$ and $p \in (1, \infty)$ with $kp > d$ so that there are continuous Sobolev embeddings, $C^{2,\lambda}(M; f_\infty^*TN) \subset W^{k,p}(M; f_\infty^*TN)$, and $L^2(M; f_\infty^*TN) \subset W^{k-2,p}(M; f_\infty^*TN)$. For example, if $d = 2$, then $k = p = 2$ will do. For $d \geq 2$, we may choose $k = 1$ and $d < p < \infty$ provided $L^2(M; \mathbb{R}) \subset W^{-1,p}(M; \mathbb{R})$ is a continuous embedding or, equivalently, $W^{1,p'}(M; \mathbb{R}) \subset L^2(M; \mathbb{R})$ is a continuous embedding, where $p' = p/(p - 1)$. According to [2, Theorem 4.12] when $1 \leq p' < d$, the latter embedding is continuous if $(p')^* = dp'/(d-p') = dp/(d(p-1)-p) \geq 2$. But we must choose $p > d$ when $k = 1$ in Theorem 2.2 and if $p = d$, then $dp/(d(p-1)-p) = d^2/(d(d-1)-d) = d/(d-2) \geq 2$ implies $d \geq 2d-4$ or $d \leq 4$. Hence, Theorem 2.2 implies Theorem 2.4 when $d = 2, 3$ (the case $d = 4$ is excluded since $p > d$ leads to $d < 4$ in the preceding inequalities). For arbitrary $d \geq 2$, another abstract Lojasiewicz–Simon gradient inequality [19, Theorem 2.4.2 (i)] due to Huang implies Theorem 2.4 with the choices

$$\mathcal{X} = C^{2,\lambda}(M; f_\infty^*TN), \quad \tilde{\mathcal{X}} = C^\lambda(M; f_\infty^*TN), \quad \mathcal{H} = L^2(M; f_\infty^*TN),$$

and $\mathcal{H}_{\mathcal{A}} = W^{2,2}(M; f_\infty^*TN)$ with $\mathcal{A} = \Delta_g + 1$ in [19, Hypotheses (H1)–(H3), pages 34–35]. We refer the reader to [13] for an exposition of Huang’s [19, Theorem 2.4.2 (i)].

Alternatively, our [12, Theorem 3] implies Theorem 2.4, as we show in the proof of [12, Corollary 6].

3. A PRIORI ESTIMATE FOR THE DIFFERENCE OF TWO HARMONIC MAPS

In this section, we give two proofs of Theorem 1, based on Theorems 2.2 and 2.4, respectively. We begin with the

Lemma 3.1 (A priori $W^{2,p}$ estimate for the difference of two harmonic maps). *Let (M, g) be a closed Riemann surface, let (N, h) be a closed, smooth Riemannian manifold, and let $p \in (1, 2]$ be a constant. Then there is a constant $C = C(g, h, p) \in [1, \infty)$ with the following significance. If $f, f_\infty \in C^\infty(M; N)$ are harmonic maps and $q = 2p/(2 - p) \in (2, \infty]$, then*

$$(3.1) \quad \|f - f_\infty\|_{W^{2,p}(M; \mathbb{R}^n)} \leq C (\|df\|_{L^q(M; \mathbb{R}^n)} + \|df_\infty\|_{L^q(M; \mathbb{R}^n)} + 1) \|f - f_\infty\|_{W^{1,2}(M; \mathbb{R}^n)}.$$

Proof. Because f and f_∞ are harmonic, equation (2.3) implies that

$$\begin{aligned} \Delta_g f - A_h(df, df) &= 0, \\ \Delta_g f_\infty - A_h(df_\infty, df_\infty) &= 0, \end{aligned}$$

and, therefore,

$$(3.2) \quad \Delta_g(f - f_\infty) - A_h(df, d(f - f_\infty)) - A_h(d(f - f_\infty), df_\infty) = 0.$$

Because $1/p = 1/2 + 1/q$ by hypothesis, the preceding equality yields the estimate,

$$\|\Delta_g(f - f_\infty)\|_{L^p(M; \mathbb{R}^n)} \leq C (\|df\|_{L^q(M; \mathbb{R}^n)} + \|df_\infty\|_{L^q(M; \mathbb{R}^n)}) \|d(f - f_\infty)\|_{L^2(M; \mathbb{R}^n)},$$

with $C = C(h) \in [1, \infty)$. The standard a priori $W^{2,p}$ estimate for an elliptic, linear, scalar, second-order partial differential operator over a bounded domain in Euclidean space [14, Theorem 9.13] yields the bound,

$$\|f - f_\infty\|_{W^{2,p}(M; \mathbb{R}^n)} \leq C (\|\Delta_g(f - f_\infty)\|_{L^p(M; \mathbb{R}^n)} + \|f - f_\infty\|_{L^p(M; \mathbb{R}^n)}),$$

for a constant $C = C(g, p) \in [1, \infty)$. Combining the preceding two inequalities gives

$$\begin{aligned} \|f - f_\infty\|_{W^{2,p}(M; \mathbb{R}^n)} &\leq C (\|df\|_{L^q(M; \mathbb{R}^n)} + \|df_\infty\|_{L^q(M; \mathbb{R}^n)}) \|d(f - f_\infty)\|_{L^2(M; \mathbb{R}^n)} \\ &\quad + C \|f - f_\infty\|_{L^p(M; \mathbb{R}^n)}, \end{aligned}$$

for $C = C(g, h, p) \in [1, \infty)$. Since $p \leq 2$, this yields the desired estimate. □

Lemma 3.2 (A priori $W^{1,q}$ estimate for a harmonic map). *Let (M, g) be a closed Riemann surface, let (N, h) be a closed, smooth Riemannian manifold, and let $q \in (2, \infty)$ be a constant. Then there is a constant $\epsilon = \epsilon(g, h, q) \in (0, 1]$ with the following significance. If $f, f_\infty \in C^\infty(M; N)$ are harmonic maps obeying (1.2), then*

$$(3.3) \quad \|f\|_{W^{1,q}(M; \mathbb{R}^n)} \leq 1 + 3\|f_\infty\|_{W^{1,q}(M; \mathbb{R}^n)}.$$

Proof. For $p \in (1, 2)$ defined by $p^* := 2p/(2-p) = q$, we observe that $W^{1,p}(M; \mathbb{R}) \subset L^q(M; \mathbb{R})$ is a continuous Sobolev embedding by [2, Theorem 4.12]. Hence, the estimate (3.1) yields

$$\begin{aligned} \|f - f_\infty\|_{W^{1,q}(M; \mathbb{R}^n)} &\leq C \|f - f_\infty\|_{W^{2,p}(M; \mathbb{R}^n)} \\ &\leq C (\|df\|_{L^q(M; \mathbb{R}^n)} + \|df_\infty\|_{L^q(M; \mathbb{R}^n)} + 1) \|f - f_\infty\|_{W^{1,2}(M; \mathbb{R}^n)}. \end{aligned}$$

Therefore,

$$\begin{aligned} \|f\|_{W^{1,q}(M; \mathbb{R}^n)} &\leq \|f - f_\infty\|_{W^{1,q}(M; \mathbb{R}^n)} + \|f_\infty\|_{W^{1,q}(M; \mathbb{R}^n)} \\ &\leq C \|df\|_{L^q(M; \mathbb{R}^n)} \|f - f_\infty\|_{W^{1,2}(M; \mathbb{R}^n)} \\ &\quad + C (\|df_\infty\|_{L^q(M; \mathbb{R}^n)} + 1) \|f - f_\infty\|_{W^{1,2}(M; \mathbb{R}^n)} + \|f_\infty\|_{W^{1,q}(M; \mathbb{R}^n)}. \end{aligned}$$

Choosing $\epsilon = \epsilon(g, h, q) \leq 1/(2C)$ in (1.2) and applying rearrangement in the preceding inequality yields

$$\|f\|_{W^{1,q}(M; \mathbb{R}^n)} \leq \|df_\infty\|_{L^q(M; \mathbb{R}^n)} + 1 + 2\|f_\infty\|_{W^{1,q}(M; \mathbb{R}^n)},$$

as desired. □

It remains to complete the

Proof of Theorem 1 using Theorem 2.2. Combining the inequalities (3.1) and (3.3) yields

$$\|f - f_\infty\|_{W^{2,p}(M; \mathbb{R}^n)} \leq C (1 + \|f_\infty\|_{W^{1,q}(M; \mathbb{R}^n)}) \|f - f_\infty\|_{W^{1,2}(M; \mathbb{R}^n)}.$$

We now fix $p \in (1, 2)$ and $q = 2p/(2 - p)$ (say with $p = 3/2$) and choose $\epsilon = \epsilon(f_\infty, g, h) \in (0, 1]$ in (1.2) small enough that

$$\epsilon C (1 + \|f_\infty\|_{W^{1,q}(M;\mathbb{R}^n)}) \leq \sigma,$$

where the constant $\sigma \in (0, 1]$ is as in Theorem 2.2. Consequently,

$$\|f - f_\infty\|_{W^{2,p}(M;\mathbb{R}^n)} < \sigma,$$

and the hypothesis (2.4) is satisfied. The Lojasiewicz–Simon gradient inequality (2.5) (with $d = k = 2$) in Theorem 2.2 therefore yields

$$\|\mathcal{M}(f)\|_{L^p(M;f^*TN)} \geq Z|\mathcal{E}(f) - \mathcal{E}(f_\infty)|^\theta.$$

But $\mathcal{M}(f) = 0$ since f is harmonic and thus $\mathcal{E}(f) = \mathcal{E}(f_\infty)$. □

It is possible to give an alternative proof of Theorem 1 using Theorem 2.4 with the aid of an elliptic bootstrapping argument to fulfill the stronger hypothesis (2.6). We first observe that Lemma 3.1 can be strengthened to give

Lemma 3.3 (A priori $W^{k,p}$ estimate for the difference of two harmonic maps). *Let (M, g) be a closed Riemann surface, let (N, h) be a closed, smooth Riemannian manifold, let $p \in (1, \infty)$ be a constant, let $k \geq 2$ be an integer, and let $f_\infty \in C^\infty(M; N)$ be a harmonic map. Then there are constants $\epsilon = \epsilon(f_\infty, g, h, k, p) \in (0, 1]$ and $C = C(f_\infty, g, h, k, p) \in [1, \infty)$ with the following significance. If $f \in C^\infty(M; N)$ is a harmonic map that obeys (1.2), then*

$$(3.4) \quad \|f - f_\infty\|_{W^{k,p}(M;\mathbb{R}^n)} \leq C\|f - f_\infty\|_{W^{1,2}(M;\mathbb{R}^n)}.$$

Proof. For $k = 2$ and $p \in (1, 2]$, the conclusion follows by combining (3.1) and (3.3). For $k \geq 3$ and $p \in (1, \infty)$, the conclusion follows by taking derivatives of (3.2) and applying a standard elliptic bootstrapping argument. □

We can now give the

Proof of Theorem 1 using Theorem 2.4. For $p \in (1, \infty)$ and $\lambda \in (0, 1)$ and large enough $k = k(g, p, \lambda) \geq 2$, there is a continuous Sobolev embedding, $W^{k,p}(M; \mathbb{R}^n) \subset C^{2,\lambda}(M; \mathbb{R}^n)$, and thus a constant $C = C(g, k, p, \lambda) \in [1, \infty)$ such that

$$\|f - f_\infty\|_{C^{2,\lambda}(M;\mathbb{R}^n)} \leq C\|f - f_\infty\|_{W^{k,p}(M;\mathbb{R}^n)}.$$

Combining the preceding inequality with (3.4) yields the bound

$$\|f - f_\infty\|_{C^{2,\lambda}(M;\mathbb{R}^n)} \leq C\|f - f_\infty\|_{W^{1,2}(M;\mathbb{R}^n)},$$

for a constant $C = C(f_\infty, g, h, k, p, \lambda) \in [1, \infty)$.

We now fix k, p, λ and choose $\epsilon = \epsilon(f_\infty, g, h) \in (0, 1]$ in (1.2) small enough that $C\epsilon \leq \sigma$, where the constant $\sigma \in (0, 1]$ is as in Theorem 2.4. Consequently,

$$\|f - f_\infty\|_{C^{2,\lambda}(M;\mathbb{R}^n)} < \sigma,$$

and the hypothesis (2.6) is satisfied. The Lojasiewicz–Simon gradient inequality (2.7) in Theorem 2.4 therefore yields

$$\|\mathcal{M}(f)\|_{L^2(M;f^*TN)} \geq Z|\mathcal{E}(f) - \mathcal{E}(f_\infty)|^\theta.$$

Again, $\mathcal{M}(f) = 0$ since f is harmonic and thus $\mathcal{E}(f) = \mathcal{E}(f_\infty)$. □

ACKNOWLEDGMENTS

The author is very grateful to the Institute for Advanced Study, Princeton, for their support during the preparation of this article. The author thanks Sagun Chanillo, Fernando Codá Marques, Joel Hass, Tobias Lamm, Paul Larain, Fang-Hua Lin, Thomas Parker, Tristan Rivière, Michael Taylor, Peter Topping, Karen Uhlenbeck, and especially Manouosos Maridakis for helpful questions and comments that influenced the development of this article.

REFERENCES

- [1] Toshiaki Adachi and Toshikazu Sunada, *Energy spectrum of certain harmonic mappings*, *Compositio Math.* **56** (1985), no. 2, 153–170. MR809864
- [2] Robert A. Adams and John J. F. Fournier, *Sobolev spaces*, 2nd ed., *Pure and Applied Mathematics* (Amsterdam), vol. 140, Elsevier/Academic Press, Amsterdam, 2003. MR2424078
- [3] Pascal Auscher, Alan McIntosh, and Andrew J. Morris, *Calderón reproducing formulas and applications to Hardy spaces*, *Rev. Mat. Iberoam.* **31** (2015), no. 3, 865–900. MR3420479
- [4] Melvin S. Berger, *Nonlinearity and functional analysis*, Academic Press [Harcourt Brace Jovanovich, Publishers], New York-London, 1977. Lectures on nonlinear problems in mathematical analysis; *Pure and Applied Mathematics*. MR0488101
- [5] Volker Branding, *Some aspects of Dirac-harmonic maps with curvature term*, *Differential Geom. Appl.* **40** (2015), 1–13. MR3333092
- [6] Volker Branding, *The heat flow for the full bosonic string*, *Ann. Global Anal. Geom.* **50** (2016), no. 4, 347–365. MR3573990
- [7] Andrea Carbonaro, Alan McIntosh, and Andrew J. Morris, *Local Hardy spaces of differential forms on Riemannian manifolds*, *J. Geom. Anal.* **23** (2013), no. 1, 106–169. MR3010275
- [8] Isaac Chavel, *Eigenvalues in Riemannian geometry*, *Pure and Applied Mathematics*, vol. 115, Academic Press, Inc., Orlando, FL, 1984. Including a chapter by Burton Randol; With an appendix by Jozef Dodziuk. MR768584
- [9] Qun Chen, Jürgen Jost, Jiayu Li, and Guofang Wang, *Dirac-harmonic maps*, *Math. Z.* **254** (2006), no. 2, 409–432. MR2262709
- [10] J. Eells and L. Lemaire, *A report on harmonic maps*, *Bull. London Math. Soc.* **10** (1978), no. 1, 1–68. MR495450
- [11] J. Eells and L. Lemaire, *Another report on harmonic maps*, *Bull. London Math. Soc.* **20** (1988), no. 5, 385–524. MR956352
- [12] P. M. N. Feehan and M. Maridakis, *Lojasiewicz–Simon gradient inequalities for analytic and Morse–Bott functionals on Banach spaces and applications to harmonic maps*, arXiv:1510.03817v6.
- [13] ———, *Lojasiewicz–Simon gradient inequalities for coupled Yang–Mills energy functionals*, *Memoirs of the American Mathematical Society*, American Mathematical Society, Providence, RI, in press, arXiv:1510.03815v4.
- [14] David Gilbarg and Neil S. Trudinger, *Elliptic partial differential equations of second order*, 2nd ed., *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*, vol. 224, Springer-Verlag, Berlin, 1983. MR737190
- [15] Robert E. Greene and Howard Jacobowitz, *Analytic isometric embeddings*, *Ann. of Math.* (2) **93** (1971), 189–204. MR0283728
- [16] Richard S. Hamilton, *Harmonic maps of manifolds with boundary*, *Lecture Notes in Mathematics*, Vol. 471, Springer-Verlag, Berlin-New York, 1975. MR0482822
- [17] Frédéric Hélein, *Harmonic maps, conservation laws and moving frames*, 2nd ed., *Cambridge Tracts in Mathematics*, vol. 150, Cambridge University Press, Cambridge, 2002. Translated from the 1996 French original; With a foreword by James Eells. MR1913803
- [18] Frédéric Hélein and John C. Wood, *Harmonic maps*, *Handbook of global analysis*, Elsevier Sci. B. V., Amsterdam, 2008, pp. 417–491, 1213. MR2389639
- [19] Sen-Zhong Huang, *Gradient inequalities*, *Mathematical Surveys and Monographs*, vol. 126, American Mathematical Society, Providence, RI, 2006. With applications to asymptotic behavior and stability of gradient-like systems. MR2226672

- [20] Jürgen Jost, *Harmonic mappings between Riemannian manifolds*, Proceedings of the Centre for Mathematical Analysis, Australian National University, vol. 4, Australian National University, Centre for Mathematical Analysis, Canberra, 1984. MR756629
- [21] Jürgen Jost, *Harmonic maps between surfaces*, Lecture Notes in Mathematics, vol. 1062, Springer-Verlag, Berlin, 1984. MR754769
- [22] Jürgen Jost, *Two-dimensional geometric variational problems*, Pure and Applied Mathematics (New York), John Wiley & Sons, Ltd., Chichester, 1991. A Wiley-Interscience Publication. MR1100926
- [23] Jürgen Jost, *Unstable solutions of two-dimensional geometric variational problems*, Differential geometry: partial differential equations on manifolds (Los Angeles, CA, 1990), Proc. Sympos. Pure Math., vol. 54, Amer. Math. Soc., Providence, RI, 1993, pp. 205–244. MR1216586
- [24] Jürgen Jost, *Riemannian geometry and geometric analysis*, 6th ed., Universitext, Springer, Heidelberg, 2011. MR2829653
- [25] J. Jost, L. Liu, and M. Zhu, *Geometric analysis of the action functional of the nonlinear supersymmetric sigma model*, (2015), Max Planck Institute for Mathematics in the Sciences preprint, www.mis.mpg.de/de/publications/preprints/2015/prepr2015-77.html.
- [26] Heaseung Kwon, *Asymptotic convergence of harmonic map heat flow*, ProQuest LLC, Ann Arbor, MI, 2002. Thesis (Ph.D.)—Stanford University. MR2703296
- [27] Yuxiang Li and Youde Wang, *A counterexample to the energy identity for sequences of α -harmonic maps*, Pacific J. Math. **274** (2015), no. 1, 107–123. MR3320872
- [28] Fanghua Lin, *Mapping problems, fundamental groups and defect measures*, Acta Math. Sin. (Engl. Ser.) **15** (1999), no. 1, 25–52. MR1701132
- [29] Qingyue Liu and Yunyan Yang, *Rigidity of the harmonic map heat flow from the sphere to compact Kähler manifolds*, Ark. Mat. **48** (2010), no. 1, 121–130. MR2594589
- [30] Roger Moser, *Partial regularity for harmonic maps and related problems*, World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2005. MR2155901
- [31] John Nash, *The imbedding problem for Riemannian manifolds*, Ann. of Math. (2) **63** (1956), 20–63. MR0075639
- [32] J. Nash, *Analyticity of the solutions of implicit function problems with analytic data*, Ann. of Math. (2) **84** (1966), 345–355. MR0205266
- [33] Thomas H. Parker, *Bubble tree convergence for harmonic maps*, J. Differential Geom. **44** (1996), no. 3, 595–633. MR1431008
- [34] J. Sacks and K. Uhlenbeck, *The existence of minimal immersions of 2-spheres*, Ann. of Math. (2) **113** (1981), no. 1, 1–24. MR604040
- [35] J. Sacks and K. Uhlenbeck, *Minimal immersions of closed Riemann surfaces*, Trans. Amer. Math. Soc. **271** (1982), no. 2, 639–652. MR654854
- [36] Richard M. Schoen, *Analytic aspects of the harmonic map problem*, Seminar on nonlinear partial differential equations (Berkeley, Calif., 1983), Math. Sci. Res. Inst. Publ., vol. 2, Springer, New York, 1984, pp. 321–358. MR765241
- [37] Stephen Semmes, *A primer on Hardy spaces, and some remarks on a theorem of Evans and Müller*, Comm. Partial Differential Equations **19** (1994), no. 1-2, 277–319. MR1257006
- [38] Leon Simon, *Asymptotics for a class of nonlinear evolution equations, with applications to geometric problems*, Ann. of Math. (2) **118** (1983), no. 3, 525–571. MR727703
- [39] Leon Simon, *Isolated singularities of extrema of geometric variational problems*, Harmonic mappings and minimal immersions (Montecatini, 1984), Lecture Notes in Math., vol. 1161, Springer, Berlin, 1985, pp. 206–277. MR821971
- [40] Leon Simon, *Theorems on regularity and singularity of energy minimizing maps*, Lectures in Mathematics ETH Zürich, Birkhäuser Verlag, Basel, 1996. Based on lecture notes by Norbert Hungerbühler. MR1399562
- [41] Elias M. Stein, *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*, Princeton Mathematical Series, vol. 43, Princeton University Press, Princeton, NJ, 1993. With the assistance of Timothy S. Murphy; Monographs in Harmonic Analysis, III. MR1232192
- [42] Michael Struwe, *Variational methods*, 4th ed., Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], vol. 34, Springer-Verlag, Berlin, 2008. Applications to nonlinear partial differential equations and Hamiltonian systems. MR2431434

- [43] Michael Taylor, *Hardy spaces and BMO on manifolds with bounded geometry*, J. Geom. Anal. **19** (2009), no. 1, 137–190. MR2465300
- [44] Michael E. Taylor, *Tools for PDE*, Mathematical Surveys and Monographs, vol. 81, American Mathematical Society, Providence, RI, 2000. Pseudodifferential operators, paradifferential operators, and layer potentials. MR1766415
- [45] Peter Miles Topping, *Rigidity in the harmonic map heat flow*, J. Differential Geom. **45** (1997), no. 3, 593–610. MR1472890
- [46] Karen Uhlenbeck, *Harmonic maps into Lie groups: classical solutions of the chiral model*, J. Differential Geom. **30** (1989), no. 1, 1–50. MR1001271
- [47] Hajime Urakawa, *Calculus of variations and harmonic maps*, Translations of Mathematical Monographs, vol. 132, American Mathematical Society, Providence, RI, 1993. Translated from the 1990 Japanese original by the author. MR1252178
- [48] Giorgio Valli, *On the energy spectrum of harmonic 2-spheres in unitary groups*, Topology **27** (1988), no. 2, 129–136. MR948176
- [49] Yuanlong Xin, *Geometry of harmonic maps*, Progress in Nonlinear Differential Equations and their Applications, vol. 23, Birkhäuser Boston, Inc., Boston, MA, 1996. MR1391729

DEPARTMENT OF MATHEMATICS, RUTGERS, THE STATE UNIVERSITY OF NEW JERSEY, 110 FREL-
INGHUYSEN ROAD, PISCATAWAY, NEW JERSEY 08854-8019

Email address: `feehan@math.rutgers.edu`