

THE EXPONENTIAL INTEGRABILITY OF QUASI-ADDITIVE FUNCTIONALS OF GAUSSIAN VECTORS

UDC 519.21

V. V. BULDYGIN

ABSTRACT. We study the exponential integrability of quasi-additive functionals of Gaussian random vectors.

1. DEFINITIONS AND AUXILIARY RESULTS

Let (V, \mathcal{F}) be a *measurable vector space*. This means that V is a real vector space where the addition of its elements and the multiplication of its elements by real numbers agree with the σ -field \mathcal{F} of subsets of the space V . Let (Ω, F, \mathbf{P}) be a probability space.

A mapping X from Ω to V measurable with respect to the σ -fields F and \mathcal{F} is called a *V -valued random vector*. The independence of a family of V -valued random vectors is defined in a standard way (see [1]). A V -valued random vector Y is called a *copy* of a V -valued random vector X if X and Y have the same distribution.

Definition 1. A V -valued random vector X is called *Gaussian* if, for each pair (X_1, X_2) of two independent copies of the random vector X , the pair (Z_1, Z_2) , where

$$Z_1 = 2^{-1/2}(X_1 + X_2), \quad Z_2 = 2^{-1/2}(X_1 - X_2),$$

is also a pair of independent copies of X .

If $V = \mathbf{R}$ is the set of real numbers and $\mathcal{F} = B(\mathbf{R})$ is the σ -field of Borel sets, then the zero mean Gaussian random variables (and only they) satisfy Definition 1 (see, for example, [1]). This implies that for Banach spaces as well as for some general topological vector spaces V Definition 1 is equivalent to the definition of a zero mean Gaussian random vector expressed in terms of the characteristic functional (see, for example, [2]).

We consider the problem on the integrability of quasi-additive functionals of V -valued Gaussian random vectors.

Definition 2. Let (V, \mathcal{F}) be a measurable vector space. A real-valued functional $g(\cdot) = (g(x), x \in V)$ is called a *measurable quasi-additive functional* (in other words, a *measurable C -quasi-additive functional*) if

- 1) $g(x) \geq 0, x \in V$;
- 2) there exists $C \geq 1$ such that $g(x + y) \leq C(g(x) + g(y))$ for $x, y \in V$;
- 3) the mapping $x \mapsto g(x)$ is measurable with respect to the σ -fields \mathcal{F} and $B(\mathbf{R})$.

If the following additional condition holds:

- 4) $g(x) = g(-x), x \in V$,

then the functional $g(\cdot)$ is called a *C -measurable quasi-additive symmetric functional*.

2000 *Mathematics Subject Classification.* Primary 60B11, 60G15.

Remark 1. The constant C involved in condition 2) is not unique. If condition 2) holds for some constant C , then it also holds for all $C' > C$. The minimum constant C_{\min} for which condition 2) holds can be defined as the infimum of the set of numbers C satisfying that condition. This minimum constant C_{\min} can be viewed as a kind of characteristic of the functional g . However the exact value of C_{\min} does not play any role in the further discussion.

Remark 2. One can consider C -quasi-additive functionals for $0 < C < 1$, too. However

$$0 \leq \sup_{x \in V} g(x) \leq \frac{Cg(0)}{1-C} < \infty$$

for $C \in (0, 1)$, since $g(x) = g(x + 0) \leq C(g(x) + g(0))$, that is, the functional $g(\cdot)$ is uniformly bounded on V . This is a trivial case, thus we consider the case of $C \geq 1$ in condition 2).

A measurable quasi-additive functional $g(\cdot)$ is called *measurable semiadditive* if $C = 1$. Measurable seminorms give a well-known example of semiadditive symmetric functionals. Measurable seminorms satisfy conditions 1)–4) and an extra (homogeneity) condition: $g(\lambda x) = |\lambda|g(x)$ for all $x \in V$ and $\lambda \in \mathbf{R}$. Measurable quasi-norms are a more general example of semiadditive functionals for which $g(\lambda x) \leq g(x)$ for all $x \in V$ and $\lambda \in [-1, 1]$.

Simple examples of *measurable* C -quasi-additive functionals for $C > 1$ are given by convex measurable functionals $(g(x), x \in V)$ such that $g(x) > 0$, $x \neq 0$, and

$$K = \sup_{x \neq 0} (g(2x)/g(x)) \in (2, \infty).$$

In this case, $C = K/2$. In particular, $g(\cdot) = q^r(\cdot)$ is a C -quasi-additive functional if $q(\cdot)$ is a measurable quasi-norm and $r > 1$. In this case, $C = 2^{r-1} > 1$.

It is proved in [3, 4, 5] that for various infinite-dimensional vector spaces V the following holds: *if $g(\cdot)$ is a measurable seminorm and X is a Gaussian random vector, then there exists $a_0 > 0$ such that $\mathbf{E} \exp\{ag^2(X)\} < \infty$ for all $a \in [0, a_0)$.* The Fernique [4] proof of this result is based on Definition 1. A generalization of Fernique's method allows one to prove an analogous result for semiadditive functionals (see [6]).

The aim of this paper is to prove that there are numbers b such that

$$\mathbf{E} \exp\{ag^b(X)\} < \infty$$

for every measurable C -quasi-additive functional g of a Gaussian random vector X and to show how b depends on the constant C .

2. MAIN RESULTS

The first result concerns a general C -quasi-additive functional.

Theorem 1. *Let (V, \mathcal{F}) be a measurable vector space, X be a V -valued Gaussian random vector, $C \geq 1$, and let $g(\cdot)$ be a measurable C -quasi-additive functional on V . Then for any $\varepsilon > 0$ there is a number $a_0 > 0$ (depending, generally speaking, on X , g , and ε) such that*

$$(1) \quad \mathbf{E} \exp\{ag^b(X)\} < \infty$$

for all $a \in [0, a_0)$, where

$$b = b(\varepsilon, C) = \frac{2}{1 + (1 + \varepsilon) \log_2 C}.$$

Note that $b = 2$ if $C = 1$. Therefore Theorem 1 generalizes the result on the exponential integrability of measurable semiadditive functionals.

Inequality (1) holds for all $b \in [0, b(C))$ where

$$b(C) = \frac{2}{1 + \log_2 C}$$

and $C > 1$. Note however that inequality (1) may hold with $b = b(C)$ for some measurable C -quasi-additive functionals.

Example 1. Let

$$g(x) = p^r(x), \quad x \in V,$$

where $p(\cdot)$ is a measurable semiadditive functional and $r > 1$. Then $g(\cdot)$ is a measurable 2^{r-1} -quasi-additive functional, that is, $C = 2^{r-1}$. Due to the theorem on the exponential integrability of semiadditive functionals, inequality (1) holds for $b = 2/r$. It remains to note that

$$b = \frac{2}{r} = \frac{2}{1 + \log_2 2^{r-1}} = \frac{2}{1 + \log_2 C}.$$

Example 1 shows that not only the constant b in inequality (1) can attain the value $b(C)$ but also that b does not exceed $b(C)$ in the class of C -quasi-additive functionals. A natural question arises whether inequality (1) may hold for $b = b(C)$ in the class of quasi-additive functionals. The answer is positive under certain additional restrictions on the functional $g(\cdot)$.

Theorem 2. *Let (V, \mathcal{F}) be a measurable vector space, X be a V -valued Gaussian random vector, $C \in [1, 2)$, and let $g(\cdot)$ be a measurable convex C -quasi-additive functional on V . Then there is a number $a_0 > 0$ (depending, generally speaking, on X and g) such that*

$$\mathbb{E} \exp \{ a g^b(X) \} < \infty$$

for all $a \in [0, a_0)$, where

$$b = \frac{2}{1 + \log_2 C}.$$

An answer to the question on whether Theorem 2 holds for $C \geq 2$ requires an additional investigation.

3. PROOFS OF THE THEOREMS

We need several auxiliary results in order to prove Theorem 1.

Let X_1, \dots, X_n be independent copies of a V -valued Gaussian random vector X , where $n = 2^l$ and $l \geq 1$. We introduce random vectors S_1, \dots, S_n as follows:

$$(2) \quad (S_1, \dots, S_n)^\top = U_n (X_1, \dots, X_n)^\top,$$

where “ \top ” denotes the transposition,

$$U_n = n^{-1/2} H_n$$

are real $n \times n$ matrices, and $H_n = (h_{kj})_{k,j=1}^n$ are defined by

$$H_2 = \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}, \quad H_n = \begin{pmatrix} H_{\frac{n}{2}} & H_{\frac{n}{2}} \\ H_{\frac{n}{2}} & -H_{\frac{n}{2}} \end{pmatrix}, \quad n = 2^l, \quad l > 1.$$

Therefore

$$S_k = 2^{-l/2} \sum_{j=1}^{2^l} h_{kj} X_j$$

for all $k = 1, \dots, 2^l$, where $h_{kj} = \pm 1$.

Lemma 1 ([6]). *The random vectors S_1, \dots, S_n are independent copies of the Gaussian random vector X .*

Lemma 2. *Let $C \geq 1$ and let $(g(x), x \in V)$ be a measurable C -quasi-additive functional. Then*

$$(3) \quad g(2^n x) \leq (2C)^n g(x),$$

$$(4) \quad g(x_1 + \dots + x_n) \leq Cg(x_1) + \dots + C^{n-2}g(x_{n-2}) + C^{n-1}g(x_{n-1}) + C^{n-1}g(x_n),$$

$$(5) \quad g(x_1 + \dots + x_n) \geq \frac{g(x_1)}{C} - C^{n-2} \sum_{k=2}^n g(-x_k)$$

for all positive integers n and all $x, x_1, \dots, x_n \in V$.

Proof of Lemma 2. Inequalities (3) and (4) can be proved by induction. Inequality (5) follows from (4). \square

Lemma 3. *Let X be a V -valued Gaussian random vector, $C \geq 1$, and let $(g(x), x \in V)$ be a measurable C -quasi-additive functional on V . Then*

$$(6) \quad (\mathbb{P}\{g(X) \leq s\})^{4^m-1} \mathbb{P}\{g(X) > t\} \leq \left(\mathbb{P}\left\{g(X) > \frac{t - (4^m - 1)C^{4^m-1}s}{C(2C)^m}\right\} \right)^{4^m}$$

for all positive integers m and all $t > 0$ and $s > 0$.

Proof of Lemma 3. Let X_1, \dots, X_{4^m} be independent copies of X . In view of Lemma 1, the random vectors

$$S_k = 2^{-m} \sum_{j=1}^{4^m} h_{kj} X_j, \quad k = 1, \dots, 4^m,$$

are independent copies of X .

Since $g(\cdot)$ is a C -quasi-additive symmetric functional, Lemma 2 implies

$$(2C)^m g(S_k) \geq g(2^m S_k) = g\left(\sum_{j=1}^{4^m} h_{kj} X_j\right) \\ \geq \frac{1}{C} g(h_{k1} X_1) - C^{4^m-2} \sum_{j=2}^{4^m} g(-h_{kj} X_j) = \frac{1}{C} g(X_1) - C^{4^m-2} \sum_{j=2}^{4^m} g(X_j)$$

for all $k = 1, \dots, 4^m$. This yields

$$(\mathbb{P}\{g(X) \leq s\})^{4^m-1} \mathbb{P}\{g(X) > t\} = \mathbb{P}\{g(X_1) > t, g(X_2) \leq s, \dots, g(X_{4^m}) \leq s\} \\ \leq \mathbb{P}\left\{\frac{1}{C} g(X_1) - C^{4^m-2} \sum_{j=2}^{4^m} g(X_j) > \frac{t - (4^m - 1)C^{4^m-1}s}{C}\right\} \\ \leq \mathbb{P}\left\{\bigcap_{k=1}^{4^m} \left\{g(S_k) > \frac{t - (4^m - 1)C^{4^m-1}s}{C(2C)^m}\right\}\right\} \\ = \left(\mathbb{P}\left\{g(X) > \frac{t - (4^m - 1)C^{4^m-1}s}{C(2C)^m}\right\}\right)^{4^m}$$

for all $t > 0$ and $s > 0$. \square

Proof of Theorem 1. First we consider the case of a C -quasi-additive symmetric functional $g(\cdot)$. Consider a positive number $s = t_0$ such that

$$c = \mathbb{P}\{g(X) \leq t_0\} > 1/2.$$

For a given positive integer m let the sequence of positive numbers $(t_n, n \geq 1)$ be defined as follows:

$$t_n = ((\gamma + 1)\beta^n - \gamma)t_0, \quad n \geq 1,$$

where

$$\gamma = \frac{\alpha}{\beta - 1}, \quad \alpha = (4^m - 1)C^{4^m - 1}, \quad \beta = C(2C)^m \geq 2.$$

The sequence $(t_n, n \geq 1)$ is such that

$$\frac{t_n - \alpha t_0}{\beta} = t_{n-1}, \quad n \geq 1,$$

and $t_n \rightarrow \infty, n \rightarrow \infty$.

Let $z_n, n \geq 0$, be real numbers such that

$$z_0 = \frac{1 - c}{c}, \quad cz_n = P\{g(X) > t_n\}, \quad n \geq 1.$$

Note that $z_0 < 1$. It follows from (6) that $z_n \leq (z_{n-1})^{4^m}, n \geq 1$. Thus

$$P\{g(X) > t_n\} \leq c(z_0)^{4^{mn}}, \quad n \geq 1,$$

whence

$$\begin{aligned} E \exp\{ag^b(X)\} &\leq c \left[\exp\{at_0^b\} + \sum_{n=1}^{\infty} z_0^{4^{m(n-1)}} \exp\{a(t_0(1 + \gamma))^b \beta^{bn}\} \right] \\ &= c \left[\exp\{at_0^b\} + \sum_{n=1}^{\infty} z_0^{4^{m(n-1)}} \exp\{a(t_0(1 + \gamma))^b 2^{nb \log_2 \beta}\} \right] \\ &= c \left[\exp\{at_0^b\} + \sum_{n=1}^{\infty} z_0^{4^{m(n-1)}} \exp\{a(t_0(1 + \gamma))^b 4^{mn}\} \right] \\ &= c \left[\exp\{at_0^b\} + \sum_{n=1}^{\infty} \exp\{4^{mn}[a(t_0(1 + \gamma))^b - (|\ln z_0|/4^m)]\} \right] \end{aligned}$$

for

$$b = b\left(\frac{1}{m}, C\right) = \frac{2}{1 + (1 + \frac{1}{m}) \log_2 C}.$$

This implies that the series converges and

$$E \exp\{ag^b(X)\} < \infty$$

for $a < |\ln z_0|/[4^m a(t_0(1 + \gamma))^b]$. This completes the proof of the theorem for C -quasi-additive symmetric functionals $g(\cdot)$, since m is arbitrary.

Now let $g(\cdot)$ be a measurable C -quasi-additive functional on V . Then

$$\tilde{g}(x) = \max\{g(x), g(-x)\}, \quad x \in V,$$

is a measurable C -quasi-additive symmetric functional on V . Thus there is a number $a_0 > 0$ such that

$$E \exp\{ag^b(X)\} \leq E \exp\{a\tilde{g}^b(x)\} < \infty$$

for all $a \in [0, a_0]$. □

Remark 3. The proof of Theorem 1 is simpler for $C = 1$. In this case, Theorem 1 follows from Lemma 3 for $m = 1$ (see [6]).

The following result is needed for the proof of Theorem 2.

Lemma 4. *Let X be a V -valued Gaussian random vector, $C \in [1, 2)$, and let $g(\cdot)$ be a measurable convex C -quasi-additive functional on V . If*

$$(7) \quad \mathbf{E} \exp\{a_1 g(X)\} < \infty$$

for some $a_1 > 0$, then there is a $a > 0$ such that

$$\mathbf{E} \exp\{ag^b(X)\} < \infty,$$

where

$$b = \frac{2}{1 + \log_2 C}.$$

Proof of Lemma 4. It follows from Lemma 1 that

$$\mathbf{P}\{g(X) \geq t2^s\} = \mathbf{P}\left\{g\left(\frac{X_1 + \cdots + X_{4^n}}{2^n}\right) \geq t2^s\right\}$$

for all positive integers n and all $s, t > 0$ where X_1, \dots, X_{4^n} are independent copies of the random vector X .

Since $g(\cdot)$ is a convex C -quasi-additive functional,

$$\begin{aligned} \mathbf{P}\left\{g\left(\frac{X_1 + \cdots + X_{4^n}}{2^n}\right) \geq t2^s\right\} &= \mathbf{P}\left\{g\left(\frac{2^n(X_1 + \cdots + X_{4^n})}{4^n}\right) \geq t2^s\right\} \\ &\leq \mathbf{P}\{g(2^n X_1) + \cdots + g(2^n X_{4^n}) \geq t2^{s+2n}\} \\ &\leq \mathbf{P}\left\{g(X_1) + \cdots + g(X_{4^n}) \geq \frac{t2^{s+n}}{C^n}\right\}. \end{aligned}$$

Condition (7) and the Markov inequality imply that

$$\begin{aligned} \mathbf{P}\{g(X) \geq t2^s\} &\leq \mathbf{P}\left\{g(X_1) + \cdots + g(X_{4^n}) \geq \frac{t2^{s+n}}{C^n}\right\} \\ &\leq \exp\left\{-a_1 t \frac{2^{s+n}}{C^n}\right\} \mathbf{E} \exp\{a_1(g(X_1) + \cdots + g(X_{4^n}))\} \\ &= \exp\left\{-a_1 t \frac{2^{s+n}}{C^n}\right\} (\mathbf{E} \exp\{a_1 g(X)\})^{4^n} = \exp\left\{-a_1 t \frac{2^{s+n}}{C^n} + a_2 4^n\right\} \end{aligned}$$

for all positive integers n and all $s, t > 0$ where $a_2 = \ln \mathbf{E} \exp\{a_1 g(X)\} \in (0, \infty)$. In the case of $s = n(1 + \log_2 C) = 2n/b$ one has

$$\mathbf{P}\{g(X) \geq t4^{n/b}\} \leq \exp\{-(a_1 t - a_2)4^n\}$$

for all positive integers n and all $t > 0$. This implies that

$$\mathbf{E} \exp\{ag^b(X)\} \leq \exp\{4at^b\} + \sum_{n=1}^{\infty} \exp\{-(a_1 t - 4at^b - a_2)4^n\}.$$

The series converges and

$$\mathbf{E} \exp\{ag^b(X)\} < \infty$$

if $t > a_2/a_1$ and $0 \leq a < (a_1 t - a_2)/4t^b$. \square

Proof of Theorem 2. Theorem 2 for $C = 1$ follows from Theorem 1 immediately. Condition (7) holds by Theorem 1 if $C \in (1, 2)$, in which case one needs to apply Lemma 4. \square

BIBLIOGRAPHY

1. V. P. Skitovich, *Linear forms of independent variables and the normal law of distribution*, *Izv. Akad. Nauk SSSR, Ser. Mat.* **18** (1954), 952. (Russian) MR **16**:52a
2. N. N. Vakhania, V. I. Tarieladze, and S. A. Chobanyan, *Probability Distributions on Banach Spaces*, “Nauka”, Moscow, 1985; English transl., Kluwer, Dordrecht, 1987. MR **86j**:60014; MR **97k**:60007
3. A. V. Skorokhod, *A remark on Gaussian measures in a Banach space*, *Teor. Veroyatnost. i Primenen.* **15** (1970), no. 3, 519–520; English transl., *Theory Probab. Appl.* **15** (1971), 508–509. MR **43**:3417
4. X. Fernique, *Intégrabilité des vecteurs gaussiens*, *C. R. Acad. Sci. Paris Sér. I. Math.* **270** (1970), no. 7, 1698–1699. MR **42**:1170
5. H. J. Landau and L. A. Shepp, *On the supremum of a Gaussian process*, *Sankhya* **32** (1970), no. 4, 369–378. MR **44**:3381
6. V. V. Buldygin and D. M. Severnyuk, *On the exponential integrability of semiadditive functionals of Gaussian vectors* *Teor. Imovirnost. Matem. Statist.* **60** (1999), 11–16; English transl., *Theory Probab. Math. Statist.* **60** (2000), 13–18.

DEPARTMENT OF ANALYSIS AND PROBABILITY THEORY, NATIONAL TECHNICAL UNIVERSITY OF UKRAINE (KPI), PROSPEKT PEREMOGY 37, KYIV-56 02056, UKRAINE
E-mail address: valbuld@comsys.ntu-kpi.kiev.ua

Received 13/DEC/2002

Translated by OLEG KLESOV