

**EXPLICIT EXTRAPOLATION FORMULAS
FOR CORRELATION MODELS OF HOMOGENEOUS ISOTROPIC
RANDOM FIELDS**

UDC 519.21

N. V. SEMENOVSKA AND M. I. YADRENKO

ABSTRACT. For some correlation models of homogeneous isotropic random fields, we obtain explicit formulas for linear extrapolation of a random field to the center of a sphere from observations on the sphere.

1. HOMOGENEOUS ISOTROPIC RANDOM FIELDS ON \mathbf{R}^n

A random function $\xi(x)$ ($x \in \mathbf{R}^n$) is called a homogeneous isotropic random field if $\mathbf{E} \xi(x) = \text{const}$ (in what follows we assume that $\mathbf{E} \xi(x) = 0$), $\mathbf{E} \xi^2(x) < \infty$, and $\mathbf{E} \xi(x)\xi(y) = \varphi(r)$ depends only on the distance $r = |x - y|$ between points x and y .

The correlation function $\varphi(r)$ of a homogeneous isotropic random field admits the following representation (see, for example, [1, 2]):

$$(1) \quad \varphi(r) = 2^{(n-2)/2} \Gamma\left(\frac{n}{2}\right) \int_0^\infty \frac{J_{(n-2)/2}(\lambda r)}{(\lambda r)^{(n-2)/2}} dF(\lambda)$$

where F is a nondecreasing bounded function on $[0, +\infty]$ and J is the Bessel function of a real argument. Relation (1) is called the *spectral representation* of the correlation function of the homogeneous isotropic random field, while the function $F(\lambda)$ is called the *spectral function* of the field. If $F(\lambda)$ is absolutely continuous, that is, there exists a function $f(\lambda)$ such that

$$F(\lambda) = \int_0^\lambda f(u) du,$$

then $f(\lambda)$ is called the *spectral density* of the homogeneous and isotropic random field. Relation (1) in the case of $n = 1$ is reduced to the well-known expansion

$$\varphi(r) = \int_0^\infty \cos(\lambda r) dF(\lambda)$$

of the correlational function of a real stationary process.

The spectral density $f(\lambda)$ can be expressed in terms of the correlational function $\varphi(r)$, namely

$$(2) \quad f(\lambda) = \int_0^\infty \sqrt{\lambda r} J_{(n-2)/2}(\lambda r) \frac{(\lambda r)^{(n-1)/2}}{2^{(n-2)/2} \Gamma(n/2)} \varphi(r) dr.$$

2000 *Mathematics Subject Classification.* Primary 60J60, 60J12, 60J25.

Key words and phrases. Random fields, Whittle–Mattern correlation model, linear extrapolation.

The spectral function $F(\lambda)$ can also be expressed in terms of $\varphi(r)$ as follows:

$$(3) \quad F(\lambda) = \frac{1}{2^{(n-2)/2}\Gamma(n/2)} \int_0^\infty \sqrt{\lambda r} J_{(n-2)/2}(\lambda r) (\lambda r)^{n/2} \frac{\varphi(r)}{r} dr$$

(see [1]). Note that relation (1) implies that $\varphi(0) = F(+\infty)$.

Let Φ_n be the class of correlation functions of mean square continuous, homogeneous isotropic random fields defined on the Euclidean space \mathbf{R}^n . The classes Φ_n are non-increasing in n ,

$$(4) \quad \Phi_1 \supset \Phi_2 \supset \Phi_3 \supset \dots \supset \Phi_n \supset \dots$$

The class

$$\Phi_\infty = \bigcap_{n \geq 1} \Phi_n$$

consists of functions $\varphi(r)$ of the following form:

$$(5) \quad \varphi(r) = \int_0^\infty e^{-\lambda^2 r^2} dF(\lambda)$$

(see [1]).

Assume that a homogeneous isotropic random field $\xi(x)$ is observed on a sphere and the problem is to find the linear extrapolation formula for $\xi(0)$ that is the best estimator for $\xi(0)$ in the sense of the minimum mean square error.

The following result is obtained in [1].

Theorem 1. *The extrapolation to the center O of the sphere*

$$S_r = \{x: |x| = r\}$$

of a mean square continuous and homogeneous random field $\xi(x)$ defined on the n -dimensional Euclidean space and observed on the sphere S_r is given by

$$(6) \quad \eta(0) = \frac{r^{(n-2)/2}}{2\pi^{n/2}} \cdot \frac{\int_0^\infty \lambda^{(2-n)/2} J_{(n-2)/2}(\lambda r) dF(\lambda)}{\int_0^\infty \lambda^{2-n} J_{(n-2)/2}^2(\lambda r) dF(\lambda)} \int_{S_n} \xi(x) dm_n$$

where $m_n(\cdot)$ is Lebesgue measure on the unit sphere in \mathbf{R}^n . The mean square error of extrapolation

$$\sigma^2(0, r) = E[\xi(0)]^2 - E[\eta(0)]^2$$

is given by

$$(7) \quad \sigma^2(0, r) = F(+\infty) - \frac{[\int_0^\infty \lambda^{(2-n)/2} J_{(n-2)/2}(\lambda r) dF(\lambda)]^2}{\int_0^\infty \lambda^{2-n} J_{(n-2)/2}^2(\lambda r) dF(\lambda)}.$$

Using (1) we rewrite $\eta(0)$ and $\sigma^2(0, r)$ as follows:

$$(8) \quad \eta(0) = \frac{1}{2^{n-1}\pi^{n/2}\Gamma(n/2)} \cdot \frac{r^{n-2}B(r)}{\int_0^\infty \lambda^{2-n} J_{(n-2)/2}^2(\lambda r) dF(\lambda)} \int_{S_n} \xi(x) dm_n,$$

$$(9) \quad \sigma^2(0, r) = F(+\infty) - \frac{1}{2^{n-2}\Gamma^2(n/2)} \cdot \frac{r^{n-2}B^2(r)}{\int_0^\infty \lambda^{2-n} J_{(n-2)/2}^2(\lambda r) dF(\lambda)}.$$

Note that one must calculate the integral

$$(10) \quad \int_0^\infty \lambda^{2-n} J_{(n-2)/2}^2(\lambda r) dF(\lambda)$$

to evaluate $\eta(0)$ and $\sigma^2(0, r)$ according to (8) and (9), respectively, in the case of a known correlation function.

TABLE 1. Buell’s list of correlation functions.

Index	Model $\varphi(r)$	Membership in Φ_n
1	e^{-r^2}	belongs to Φ_∞
2	$(1 + r^2)^{-\nu}$	belongs to Φ_n if $\nu \geq 0$
3	$(1 + r + \kappa r^2)e^{-r}$	belongs to Φ_n if $-\frac{1}{n} \leq \kappa \leq \frac{1}{3}$
3a	$(1 + r)e^{-r}$	belongs to Φ_∞
3b	$(1 + r + r^2/3)e^{-r}$	belongs to Φ_∞
3c	$(1 + r - r^2/2)e^{-r}$	belongs to Φ_2
4	$(ae^{-br} - be^{-ar})/(a - b)$	belongs to Φ_n if $b \neq a = 0$ or $a \neq b = 0$ or $a > b > 0$ or $b > a > 0$
5	$\sin r/r$	belongs to Φ_3
6	$\frac{2^{1/3}}{\Gamma(2/3)}r^{2/3}K_{2/3}(r)$	belongs to Φ_∞

2. ON MODELS OF CORRELATION FUNCTIONS OF HOMOGENEOUS AND ISOTROPIC RANDOM FIELDS

Isotropic random fields play an important role in modeling various spatial phenomena in natural science and engineering. Isotropic correlation models are used in various branches of science, for example, in geodesy, geology, geomorphology, hydrology, meteorology, agronomy, and geophysics. Buell’s list of “empirical correlation functions” ([3, p. 53]) became a standard source of correlation models in meteorology. Consider the Whittle–Mattern class of correlation functions

$$(11) \quad \varphi_\nu(r) = \frac{2^{1-\nu}}{\Gamma(\nu)}(r)^\nu K_\nu(r)$$

where K is the modified Bessel function, $\nu > 0$. If

$$\nu = k + \frac{1}{2}$$

and k is a positive integer, then (11) can be written as the product of the exponent e^{-r} and a polynomial of order k with respect to r (see [4, 8.468]). For example, the correlation functions

$$\varphi_{1/2}(r) = e^{-r}$$

and

$$\varphi_{3/2}(r) = (1 + r)e^{-r}$$

can be obtained in this way for $\nu = 1/2$ and $\nu = 3/2$, respectively.

In this paper we consider the Whittle–Mattern and Buell correlation models and, for all of them, find spectral densities and extrapolation formulas for the center of a sphere from observations of the field on the sphere.

Note that the Whittle–Mattern class contains the correlation functions of homogeneous isotropic random fields that are solutions of the following stochastic equation:

$$(12) \quad (\Delta - c^2)^m \xi(x) = w'(x)$$

where Δ is the Laplace operator on \mathbf{R}^n and $w'(x)$ is the white noise on \mathbf{R}^n ([1, 5]).

Buell [3] proposed to apply the correlation models shown in Table 1 in meteorology.

Now we turn to the Whittle–Mattern family of correlation functions (11). If $\nu = 2/3$, $3/2$, or $5/2$, then formula (11) coincides with the Buell model of indices 6, 3a, or 3b,

respectively (see [4, 8.468]). Note that the Buell model of index 3 can be written as a linear combination of the Whittle–Mattern models for $\nu = 3/2$ and $5/2$, namely

$$(13) \quad (1 + r + \kappa r^2) e^{-r} = (1 - 3\kappa)\varphi_{3/2}(r) + 3\kappa\varphi_{5/2}(r).$$

3. EXPLICIT EXTRAPOLATION FORMULAS FOR SOME CORRELATION MODELS

First we prove the following auxiliary result.

Lemma 1. *If $n > \nu > 0$, then*

$$(14) \quad \int_0^\infty J_{(n-2)/2}^2(\lambda r) \frac{\lambda}{(\lambda^2 + a^2)^{n/2+\nu}} d\lambda \\ = \frac{r^{n-2} B(1/2, (n-1)/2 + \nu)}{2\pi} \\ \times \left(\frac{B((n-1)/2, \nu)}{a^{2\nu} \Gamma(n-1)} \cdot {}_1F_2((n-1)/2; 1-\nu, n-1; a^2 r^2) \right. \\ \left. + \frac{r^{2\nu} \Gamma(-\nu)}{\Gamma(n+\nu-1)} \cdot {}_1F_2((n-1)/2 + \nu; n+\nu-1, \nu+1; a^2 r^2) \right),$$

where ${}_pF_q$ is the generalized hypergeometric series

$${}_pF_q(a_1, \dots, a_p; b_1, \dots, b_q; z) = \sum_{k=0}^{\infty} \frac{(a_1)_k \cdots (a_p)_k}{(b_1)_k \cdots (b_q)_k} \cdot \frac{z^k}{k!}.$$

Proof. We use the following integral representation for the product of Bessel functions:

$$J_\mu(z) J_\nu(z) = \frac{2}{\pi} \int_0^{\pi/2} J_{\mu+\nu}(2z \cos \theta) \cos(\mu - \nu)\theta d\theta,$$

and a representation of an integral of a Bessel function in terms of a sum of hypergeometric functions

$$\int_0^\infty J_{(n-2)/2}^2(\lambda r) \frac{\lambda d\lambda}{(\lambda^2 + a^2)^{n/2+\nu}} = \frac{2}{\pi} \int_0^{\pi/2} \int_0^\infty J_{n-2}(2\lambda r \cos \theta) \frac{\lambda d\lambda d\theta}{(\lambda^2 + a^2)^{n/2+\nu}} \\ = \frac{2}{\pi} \int_0^{\pi/2} \left(\frac{r^{n-2} B(n/2, \nu)}{2a^{2\nu} \Gamma(n-1)} \cos^{n-2} \theta \cdot {}_1F_2(n/2; 1-\nu, n-1; a^2 r^2 \cos^2 \theta) \right. \\ \left. + \frac{r^{n-2+2\nu} \Gamma(-\nu)}{2\Gamma(n+\nu-1)} \cos^{n-2+2\nu} \theta \right. \\ \left. \times {}_1F_2(n/2 + \nu; n+\nu-1, \nu+1; a^2 r^2 \cos^2 \theta) \right) d\theta$$

(see [6]). We consider separately the integrals

$$\int_0^{\pi/2} \cos^{n-2} \theta \cdot {}_1F_2(n/2; 1-\nu, n-1; a^2 r^2 \cos^2 \theta) d\theta$$

and

$$\int_0^{\pi/2} \cos^{n-2+2\nu} \theta \cdot {}_1F_2(n/2 + \nu; n+\nu-1, \nu+1; a^2 r^2 \cos^2 \theta) d\theta.$$

Using the definition of the generalized hypergeometric series ${}_pF_q$ we expand the integrand into the series, integrate the series term by term (see [4]), and then evaluate its sum:

$$\begin{aligned} & \int_0^{\pi/2} \cos^{n-2} \theta \cdot {}_1F_2 \left(n/2; 1 - \nu, n - 1; a^2 r^2 \cos^2 \theta \right) d\theta \\ &= \sum_{k=0}^{\infty} \frac{(n/2)_k}{(1 - \nu)_k (n - 1)_k} \frac{(a^2 r^2)^k}{k!} \int_0^{\pi/2} \cos^{n-2+2k} \theta d\theta \\ &= \sum_{k=0}^{\infty} \frac{(n/2)_k}{(1 - \nu)_k (n - 1)_k} \frac{(a^2 r^2)^k}{k!} \cdot \frac{1}{2} B(1/2, (n - 1)/2 + k) \\ &= \frac{1}{2} B(1/2, (n - 1)/2) \cdot {}_1F_2 \left((n - 1)/2; 1 - \nu, n - 1; a^2 r^2 \right), \\ & \int_0^{\pi/2} \cos^{n-2+2\nu} \theta \cdot {}_1F_2 \left(n/2 + \nu; n + \nu - 1, \nu + 1; a^2 r^2 \cos^2 \theta \right) d\theta \\ &= \sum_{k=0}^{\infty} \frac{(n/2 + \nu)_k}{(n + \nu - 1)_k (\nu + 1)_k} \frac{(a^2 r^2)^k}{k!} \int_0^{\pi/2} \cos^{n-2+2\nu+2k} \theta d\theta \\ &= \sum_{k=0}^{\infty} \frac{(n/2 + \nu)_k}{(n + \nu - 1)_k (\nu + 1)_k} \frac{(a^2 r^2)^k}{k!} \cdot \frac{1}{2} B(1/2, (n - 1)/2 + \nu + k) \\ &= \frac{1}{2} B(1/2, (n - 1)/2 + \nu) \cdot {}_1F_2 \left((n - 1)/2 + \nu; n + \nu - 1, \nu + 1; a^2 r^2 \right). \end{aligned}$$

Now relation (14) follows easily. □

Model 1. Consider a homogeneous isotropic random field whose correlation function is given by (11). The spectral density for this model is of the form

$$(15) \quad f_\nu(\lambda) = \frac{2\lambda^{n-1}}{B(\nu, n/2)(\lambda^2 + 1)^{n/2+\nu}}, \quad n > \nu.$$

Using (14) we obtain for $a = 1$ that

$$\begin{aligned} & \int_0^\infty \lambda^{2-n} J_{(n-2)/2}^2(\lambda r) dF(\lambda) \\ &= \frac{B(1/2, (n - 1)/2 + \nu) r^{n-2}}{\pi B(\nu, n/2)} \\ (16) \quad & \times \left(\frac{B((n - 1)/2, \nu)}{\Gamma(n - 1)} \cdot {}_1F_2 \left((n - 1)/2; 1 - \nu, n - 1; r^2 \right) \right. \\ & \left. + \frac{r^{2\nu} \Gamma(-\nu)}{\Gamma(n + \nu - 1)} \cdot {}_1F_2 \left((n - 1)/2 + \nu; n + \nu - 1, \nu + 1; r^2 \right) \right). \end{aligned}$$

Let

$$\begin{aligned} G_1(r) &= \frac{B((n - 1)/2, \nu)}{\Gamma(n - 1)} \cdot {}_1F_2 \left((n - 1)/2; 1 - \nu, n - 1; r^2 \right), \\ G_2(r) &= \frac{\Gamma(-\nu)}{\Gamma(n + \nu - 1)} \cdot {}_1F_2 \left((n - 1)/2 + \nu; n + \nu - 1, \nu + 1; r^2 \right). \end{aligned}$$

According to (8) the extrapolation formula for the center of a sphere is of the following form:

$$(17) \quad \eta(0) = \frac{2^{2-n-\nu} \pi^{(2-n)/2}}{\Gamma(n/2 + \nu) B(1/2, (n - 1)/2 + \nu)} \cdot \frac{K_\nu(r)}{r^{-\nu} G_1(r) + r^\nu G_2(r)} \int_{S_n} \xi(x) dm_n.$$

As was mentioned before, $F(+\infty) = 1$. According to (9), the mean square error of the extrapolation is

$$(18) \quad \sigma^2(0, r) = 1 - \frac{\pi 2^{4-n-2\nu} \Gamma(n/2 + \nu)}{\Gamma(n/2) \Gamma(\nu) B(1/2, (n-1)/2 + \nu)} \cdot \frac{K_\nu^2(r)}{r^{-2\nu} G_1(r) + G_2(r)}.$$

Model 2. Let $\varphi(r) = \exp\{-r^2\}$. Using relation (2) and formula (10) of Section 8.6 in [7] we get the spectral density

$$f(\lambda) = \frac{\lambda^{n-1} \exp\{-\lambda^2/4\}}{2^{n-1} \Gamma(n/2)}.$$

Taking into account relation (39) of Section 4.14 in [7], we get

$$\int_0^\infty \lambda^{2-n} J_{(n-2)/2}^2(\lambda r) f(\lambda) d\lambda = \frac{e^{-2r^2} I_{(n-2)/2}(2r^2)}{2^{n-2} \Gamma(n/2)}.$$

After simple calculations we obtain from (8) the following extrapolation formula for the center of the sphere:

$$\eta(0) = \frac{1}{2\pi^{n/2}} \cdot \frac{r^{n-2} e^{r^2}}{I_{(n-2)/2}(2r^2)} \cdot \int_{S_n} \xi(x) dm_n.$$

In view of (9) the mean square error of the extrapolation is given by

$$\sigma^2(0, r) = 1 - \frac{1}{\Gamma(n/2)} \cdot \frac{r^{n-2}}{I_{(n-2)/2}(2r^2)}.$$

Model 3. Let

$$\varphi(r) = (1 + r^2)^{-\nu}, \quad \nu > 0.$$

We apply (2) to find the spectral density. Using relation (20) of Section 8.5 in [8] we get

$$f(\lambda) = \frac{\lambda^{(n-2)/2+\nu} K_{n/2-\nu}(\lambda)}{2^{(n-2)/2+\nu-1} \Gamma(n/2) \Gamma(\nu)}.$$

According to equality (25) of Section 10.3 in [8],

$$\int_0^\infty \lambda^{2-n} J_{(n-2)/2}^2(\lambda r) f(\lambda) d\lambda = \frac{2^{2-n} r^{n-2}}{\Gamma^2(n/2)} F\left(\frac{n-1}{2}, \nu; n-1; -4r^2\right)$$

where F is the Gauss hypergeometric series

$$F(a, b; c; z) = {}_2F_1(a, b; c; z).$$

After a simple calculation we obtain from (8) the following extrapolation formula for the center of the sphere

$$\eta(0) = \frac{\Gamma(n/2)}{2\pi^{n/2}} \cdot \frac{(1 + r^2)^{-\nu}}{F((n-1)/2, \nu; n-1; -4r^2)} \int_{S_n} \xi(x) dm_n.$$

In view of (9) the mean square error of the extrapolation is given by

$$\sigma^2(0, r) = 1 - \frac{(1 + r^2)^{-2\nu}}{F((n-1)/2, \nu; n-1; -4r^2)}.$$

Model 4. Let $\varphi(r) = (1 + r)e^{-r}$. As mentioned before, this correlation model corresponds to the Whittle–Mattern model $\varphi_{3/2}(r)$. Thus we use (15) and obtain

$$f_{3/2}(\lambda) = \frac{2\lambda^{n-1}}{B(3/2, n/2) (\lambda^2 + 1)^{(n+3)/2}}, \quad n > 3/2.$$

It follows from (16) that

$$\int_0^\infty \lambda^{2-n} J_{(n-2)/2}^2(\lambda r) dF(\lambda) = \frac{nr^{n-2}}{\pi} (G_1(r) + r^3 G_2(r))$$

where

$$G_1(r) = \frac{B((n-1)/2, 3/2)}{\Gamma(n-1)} \cdot {}_1F_2\left(\frac{n-1}{2}; -\frac{1}{2}, n-1; r^2\right),$$

$$G_2(r) = \frac{4\sqrt{\pi}}{3\Gamma(n+1/2)} \cdot {}_1F_2\left(\frac{n}{2} + 1; n+1/2, \frac{5}{2}; r^2\right).$$

Since $K_{3/2} = (\pi/(2r^3))^{1/2}(1+r)e^{-r}$, we obtain from (17) the linear extrapolation formula for the center of the sphere:

$$\eta(0) = \frac{\pi^{(2-n)/2}}{2^n \Gamma(n/2 + 1)} \frac{(1+r)e^{-r}}{(G_1(r) + r^3 G_2(r))} \int_{S_n} \xi(x) dm_n.$$

The mean square error of the extrapolation is equal to

$$\sigma^2(0, r) = 1 - \frac{\pi 2^{1-n}}{\Gamma(n/2)\Gamma(n/2 + 1)} \cdot \frac{(1+r)^2 e^{-2r}}{(G_1(r) + r^3 G_2(r))}$$

(see (18)).

Model 5. Let $\varphi(r) = (1+r+r^2/3)e^{-r}$. This correlation model corresponds to the Whittle–Mattern model $\varphi_{5/2}(r)$. It follows from (15) that

$$f_{5/2}(\lambda) = \frac{2\lambda^{n-1}}{B(5/2, n/2)(\lambda^2 + 1)^{(n+5)/2}}, \quad n > 5/2.$$

Relation (16) implies that

$$\int_0^\infty \lambda^{2-n} J_{(n-2)/2}^2(\lambda r) dF(\lambda) = \frac{n(n+2)r^{n-2}}{3\pi} (g_1(r) - r^5 g_2(r))$$

where

$$g_1(r) = \frac{B((n-1)/2, 5/2)}{\Gamma(n-1)} \cdot {}_1F_2\left(\frac{(n-1)}{2}; -\frac{3}{2}, n-1; r^2\right),$$

$$g_2(r) = \frac{8\sqrt{\pi}}{15\Gamma(n+3/2)} \cdot {}_1F_2\left(\frac{n}{2} + 2; n + \frac{3}{2}, \frac{7}{2}; r^2\right).$$

According to (17) the linear extrapolation formula for the center of the sphere is given by

$$\eta(0) = \frac{3\pi^{(2-n)/2}}{2^{n+1}\Gamma(n/2 + 2)} \frac{(1+r+r^2/3)e^{-r}}{(g_1(r) - r^5 g_2(r))} \int_{S_n} \xi(x) dm_n,$$

since

$$K_{5/2} = (\pi/(8r^5))^{1/2} \left(1+r+\frac{r^2}{3}\right) e^{-r}.$$

The mean square error of the extrapolation is equal to

$$\sigma^2(0, r) = 1 - \frac{3\pi 2^{-n}}{\Gamma(n/2)\Gamma(n/2 + 2)} \cdot \frac{(1+r+r^2/3)^2 e^{-2r}}{(g_1(r) - r^5 g_2(r))}$$

(see (18)).

Model 6. Consider the following correlation model:

$$\varphi(r) = (1 + r + \kappa r^2) e^{-r}.$$

As mentioned before, this model is a linear combination of the models $\varphi_{3/2}(r)$ and $\varphi_{5/2}(r)$, namely

$$(1 + r + \kappa r^2) e^{-r} = (1 - 3\kappa)\varphi_{3/2}(r) + 3\kappa\varphi_{5/2}(r).$$

Thus the spectral density $f(\lambda)$ of this correlation model is the linear combination of $\varphi_{3/2}(r)$ and $\varphi_{5/2}(r)$ with the same coefficients. Thus

$$f(\lambda) = \frac{2(n+1)\lambda^{n-1}}{B(1/2, n/2)(1+\lambda^2)^{(n+5)/2}} ((1-3\kappa)\lambda^2 + n\kappa + 1)$$

by the results obtained for the two preceding models. The same arguments yield

$$(19) \quad \int_0^\infty \lambda^{2-n} J_{(n-2)/2}^2(\lambda r) dF(\lambda) = \frac{n}{\pi} r^{n-2} ((1-3\kappa)(G_1(r) + r^3 G_2(r)) + \kappa(n+2)(g_1(r) - r^5 g_2(r)))$$

where

$$\begin{aligned} G_1(r) &= \frac{B((n-1)/2, 3/2)}{\Gamma(n-1)} \cdot {}_1F_2\left(\frac{(n-1)}{2}; -\frac{1}{2}, n-1; r^2\right), \\ G_2(r) &= \frac{4\sqrt{\pi}}{3\Gamma(n+1/2)} \cdot {}_1F_2\left(\frac{n}{2} + 1; n + \frac{1}{2}, \frac{5}{2}; r^2\right); \\ g_1(r) &= \frac{B((n-1)/2, 5/2)}{\Gamma(n-1)} \cdot {}_1F_2\left(\frac{(n-1)}{2}; -\frac{3}{2}, n-1; r^2\right), \\ g_2(r) &= \frac{8\sqrt{\pi}}{15\Gamma(n+3/2)} \cdot {}_1F_2\left(n/2 + 2; n + \frac{3}{2}, \frac{7}{2}; r^2\right). \end{aligned}$$

Now we obtain from (8) the extrapolation formula for the center of the sphere:

$$(20) \quad \eta(0) = \frac{2^{-n} \pi^{(2-n)/2}}{\Gamma(n/2 + 1)} \times \frac{(1 + r + \kappa r^2) e^{-r}}{((1-3\kappa)(G_1(r) + r^3 G_2(r)) + \kappa(n+2)(g_1(r) - r^5 g_2(r)))} \int_{S_n} \xi(x) dm_n.$$

The mean square error of the extrapolation is given by

$$(21) \quad \sigma^2(0, r) = 1 - \frac{\pi 2^{1-n}}{\Gamma(n/2)\Gamma(n/2 + 1)} \times \frac{(1 + r + \kappa r^2)^2 e^{-2r}}{((1-3\kappa)(G_1(r) + r^3 G_2(r)) + \kappa(n+2)(g_1(r) - r^5 g_2(r)))}$$

(see (9)).

Model 7. Consider the following correlation model:

$$\varphi(r) = (1 + r - r^2/2) e^{-r}, \quad n = 2.$$

This is the preceding model with $\kappa = -1/2$, and we may use the preceding results:

$$f(\lambda) = \frac{15}{2} \lambda^3 (1 + \lambda^2)^{-7/2}.$$

Similarly we obtain from (19) that

$$\int_0^\infty \lambda^{2-n} J_{(n-2)/2}^2(\lambda r) dF(\lambda) = \frac{2}{\pi} \left(\frac{5}{2} (G_1(r) + r^3 G_2(r)) + 2 (r^5 g_2(r) - g_1(r)) \right)$$

where

$$\begin{aligned} G_1(r) &= \frac{\pi}{2} \cdot {}_1F_2\left(\frac{1}{2}; -\frac{1}{2}, 1; r^2\right), \\ G_2(r) &= \frac{2^5}{5 \cdot 3^2} \cdot {}_1F_2\left(2; \frac{5}{2}, \frac{5}{2}; r^2\right); \\ g_1(r) &= \frac{3\pi}{4} \cdot {}_1F_2\left(\frac{1}{2}; -\frac{3}{2}, 1; r^2\right), \\ g_2(r) &= \frac{2^6}{5^2 \cdot 3^2} \cdot {}_1F_2\left(3; \frac{7}{2}, \frac{7}{2}; r^2\right). \end{aligned}$$

The extrapolation formula for the center of the sphere is found from (20):

$$\eta(0) = \frac{(1 + r - r^2/2)e^{-r}}{(5(G_1(r) + r^3G_2(r)) + 4(r^5g_2(r) - g_1(r)))} \int_{S_n} \xi(x) dm_n.$$

The mean square error of the extrapolation is given by (21) as follows:

$$\sigma^2(0, r) = 1 - \frac{\pi}{2} \cdot \frac{(1 + r - r^2/2)^2 e^{-2r}}{(5/2(G_1(r) + r^3G_2(r)) + 2(r^5g_2(r) - g_1(r)))}.$$

Model 8. Consider the following correlation model:

$$\varphi(r) = (ae^{-br} - be^{-ar}) / (a - b).$$

Let $\varphi_a(r) = e^{-ar}$ and $\varphi_b(r) = e^{-br}$, and let the respective spectral densities be $f_a(\lambda)$ and $f_b(\lambda)$. The correlation function $\varphi(r)$ is a linear combination of the functions $\varphi_a(r)$ and $\varphi_b(r)$. Thus the spectral density $f(\lambda)$ corresponding to the correlation function $\varphi(r)$ is a linear combination of the functions $f_a(\lambda)$ and $f_b(\lambda)$. Using (2) and relation (4) of Section 8.6 in [8] we obtain

$$f_a(\lambda) = \frac{2a\Gamma((n + 1)/2)}{\sqrt{\pi}\Gamma(n/2)} \cdot \frac{\lambda^{n-1}}{(\lambda^2 + a^2)^{(n+1)/2}}.$$

We use equality (14) to evaluate the integral

$$\int_0^\infty \lambda^{2-n} J_{(n-2)/2}^2(\lambda r) f_a(\lambda) d\lambda.$$

We have

$$\begin{aligned} &\int_0^\infty \lambda^{2-n} J_{(n-2)/2}^2(\lambda r) \cdot \frac{a\Gamma((n + 1)/2)}{\sqrt{\pi}\Gamma(n/2)} \cdot \frac{\lambda^{n-1} d\lambda}{(a^2 + \lambda^2)^{(n+1)/2}} \\ &= \frac{ar^{n-2}}{\pi} \left(\frac{B((n - 1)/2, 1/2)}{a\Gamma(n - 1)} {}_1F_2\left(\frac{n - 1}{2}; \frac{1}{2}, n - 1; a^2r^2\right) \right. \\ &\quad \left. - \frac{2\sqrt{\pi}}{\Gamma(n - 1/2)} r {}_1F_2\left(\frac{n}{2}; \frac{3}{2}, n - \frac{1}{2}; a^2r^2\right) \right). \end{aligned}$$

Let

$$\begin{aligned} L(a, r) &= \left(\frac{B((n - 1)/2, 1/2)}{a\Gamma(n - 1)} {}_1F_2\left(\frac{n - 1}{2}; \frac{1}{2}, n - 1; a^2r^2\right) \right. \\ &\quad \left. - \frac{2\sqrt{\pi}}{\Gamma(n - 1/2)} r {}_1F_2\left(\frac{n}{2}; \frac{3}{2}, n - \frac{1}{2}; a^2r^2\right) \right). \end{aligned}$$

Then

$$\int_0^\infty \lambda^{2-n} J_{(n-2)/2}^2(\lambda r) f_a(\lambda) d\lambda = \frac{ar^{n-2}}{\pi} L(a, r).$$

Similarly

$$f_b(\lambda) = \frac{2b\Gamma((n+1)/2)}{\sqrt{\pi}\Gamma(n/2)} \cdot \frac{\lambda^{n-1}}{(\lambda^2 + b^2)^{(n+1)/2}}$$

and

$$\int_0^\infty \lambda^{2-n} J_{(n-2)/2}^2(\lambda r) f_b(\lambda) d\lambda = \frac{br^{n-2}}{\pi} L(b, r)$$

where

$$L(b, r) = \left(\frac{B((n-1)/2, 1/2)}{b\Gamma(n-1)} {}_1F_2 \left(\frac{n-1}{2}; \frac{1}{2}, n-1; b^2 r^2 \right) - \frac{2\sqrt{\pi}}{\Gamma(n-1/2)} r {}_1F_2 \left(\frac{n}{2}; \frac{3}{2}, n - \frac{1}{2}; b^2 r^2 \right) \right).$$

Thus the spectral density of the correlation model under consideration is of the following form:

$$f(\lambda) = \frac{2ab}{a-b} \frac{\Gamma((n+1)/2)}{\sqrt{\pi}\Gamma(n/2)} \left(\frac{\lambda^{n-1}}{(\lambda^2 + b^2)^{(n+1)/2}} - \frac{\lambda^{n-1}}{(\lambda^2 + a^2)^{(n+1)/2}} \right).$$

Moreover

$$\int_0^\infty \lambda^{2-n} J_{(n-2)/2}^2(\lambda r) f(\lambda) d\lambda = \frac{abr^{n-2}(L(b, r) - L(a, r))}{\pi(a-b)}.$$

The linear extrapolation formula (8) takes the form

$$\eta(0) = \frac{2^{1-n}\pi^{(2-n)/2}}{\Gamma(n/2)} \cdot \frac{ae^{-br} - be^{-ar}}{abr^{n-2}(L(b, r) - L(a, r))} \int_{S_n} \xi(x) dm_n.$$

The mean square error of extrapolation is given by (9) as follows:

$$\sigma^2(0, r) = 1 - \frac{\pi 2^{2-n}}{(a-b)\Gamma^2(n/2)} \cdot \frac{(ae^{-br} - be^{-ar})^2}{ab(L(b, r) - L(a, r))}.$$

Model 9. Let $\varphi(r) = \sin(r)/r$, $n = 3$. This is the so-called Markov type field. The problem of extrapolation for the Markov type field in \mathbf{R}^n (and, in particular, in \mathbf{R}^3) is considered in [1].

Model 10. Let $\varphi(r) = 2^{1/3}r^{2/3}K_{2/3}(r)/\Gamma(2/3)$. This correlation model corresponds to the Whittle–Matern model $\varphi_{2/3}(r)$. Thus (15) implies that

$$f_{2/3}(\lambda) = \frac{2\lambda^{n-1}}{B(2/3, n/2)(\lambda^2 + 1)^{n/2+2/3}}, \quad n > 2/3.$$

Let

$$G_1(r) = \frac{B((n-1)/2, 2/3)}{\Gamma(n-1)} \cdot {}_1F_2((n-1)/2; 1/3, n-1; r^2),$$

$$G_2(r) = \frac{\Gamma(-2/3)}{\Gamma(n-1/3)} \cdot {}_1F_2(n/2 + 1/6; n-1/3, 5/3; r^2).$$

The extrapolation formula is given by

$$\eta(0) = \frac{2^{1/3-n}\pi^{(2-n)/2}}{\Gamma(n/2 + 2/3)B(1/2, n/2 + 1/6)} \cdot \frac{K_{2/3}(r)}{r^{-2/3}G_1(r) + r^{2/3}G_2(r)} \int_{S_n} \xi(x) dm_n.$$

The mean square error of the extrapolation is given by (18) as follows:

$$\sigma^2(0, r) = 1 - \frac{\pi 2^{8/3-n}\Gamma(n/2 + 2/3)}{\Gamma(n/2)\Gamma(2/3)B(1/2, n/2 + 1/6)} \cdot \frac{K_{2/3}^2(r)}{r^{-4/3}G_1(r) + G_2(r)}.$$

BIBLIOGRAPHY

1. M. I. Yadrenko, *Spectral Theory of Random Fields*, “Vyshcha Shkola”, Kiev, 1980; English transl., Optimization Software, Inc., New York, 1983. MR0590889 (82e:60001)
2. A. M. Yaglom, *Some classes of random fields in n -dimensional space related to stationary random processes*, *Teor. Veroyatnost. i Primenen.* **2** (1957), no. 3, 292–338; English transl. in *Theory Probab. Appl.* **2** (1957), no. 3, 273–320. MR0094844 (20:1353)
3. C. E. Buell, *Correlation functions for wind and geopotential on isobaric surfaces*, *J. Appl. Meteor.* **11** (1972), 51–59.
4. I. S. Gradshteyn and I. M. Ryzhik, *Tables of Integrals, Series, and Products*, Gos. Izdat. Fiz. Mat. Literat., Moscow, 1963; English transl., Sixth edition, Academic Press, San Diego, CA, 2000. MR1773820 (2001c:00002)
5. M. I. Yadrenko, *Some statistical problems for the Whittle random field*, *Teor. Imovir. Mat. Stat.* **63** (2000), 163–167; English transl. in *Theory Probab. Math. Statist.* **63** (2001), 179–183. MR1870787 (2002m:60097)
6. G. N. Watson, *A Treatise on the Theory of Bessel Functions*, 2nd edition, Cambridge University Press, Cambridge, 1995. MR1349110 (96i:33010)
7. A. Erdélyi, W. Magnus, F. Oberhettinger, and F. G. Tricomi, *Higher Transcendental Functions*, Based in part on notes left by Harry Bateman, vol. 1, McGraw-Hill Book Company, Inc., New York–Toronto–London, 1953. MR0058756 (15:419i)
8. ———, *Higher Transcendental Functions*, Based in part on notes left by Harry Bateman, vol. 2, McGraw-Hill Book Company, Inc., New York–Toronto–London, 1953. MR0058756 (15:419i)
9. M. I. Yadrenko, *Isotropic random fields of Markov type in Euclidean and Hilbert spaces*, *Proc. All-Union Conf. Theory Prob. and Math. Statist.* (Erevan, 1958), Izdat. Akad. Nauk Armyan. SSR, Erevan, 1960, pp. 263–279. (Russian) MR0214123 (35:4974)

DEPARTMENT OF PROBABILITY THEORY AND MATHEMATICAL STATISTICS, FACULTY OF MECHANICS AND MATHEMATICS, NATIONAL TARAS SHEVCHENKO UNIVERSITY, ACADEMICIAN GLUSHKOV AVENUE 6, KYIV 03127, UKRAINE

E-mail address: `semenovsky@hotmail.ru`

DEPARTMENT OF PROBABILITY THEORY AND MATHEMATICAL STATISTICS, FACULTY OF MECHANICS AND MATHEMATICS, NATIONAL TARAS SHEVCHENKO UNIVERSITY, ACADEMICIAN GLUSHKOV AVENUE 6, KYIV 03127, UKRAINE

E-mail address: `yimi@mechmat.univ.kiev.ua`

Received 14/FEB/2003

Translated by V. SEMENOV