

EXISTENCE PROOF FOR A FIELD OF EXTREMALS TANGENT
TO A GIVEN CURVE*

BY

OSKAR BOLZA

In a recent paper,† Professor BLISS has given sufficient conditions for a minimum of the integral

$$(1) \quad J = \int_{t_0}^{t_1} F(x, y, x', y') dt$$

with respect to one-sided variations. His proof is based upon the construction of a *field of extremals tangent to a given curve*. He establishes the existence of such a field first for the special case where all curves considered are representable in the form $y = f(x)$, and then reduces the general case of parameter representation to the former by a point-transformation of the plane.

The object of the following note is to give a *direct proof for the existence of these fields* which play an important part also in other investigations of the calculus of variations.‡

§ 1. *The set of extremals tangent to a given curve.*

The terminology and assumptions concerning the function F being the same as in § 24 of my *Lectures on the Calculus of Variations*, we consider a curve of class C''

$$\tilde{C}: \quad x = \tilde{x}(a), \quad y = \tilde{y}(a), \quad A_1 \leq a \leq A_2,$$

without multiple points, which lies in the interior of the region of the x, y -plane in which the function F is supposed to be of class C''' for every $(x', y') \neq (0, 0)$, and satisfies the inequality

* Presented to the Society March 30, 1907. Received for publication February 27, 1907.

† Transactions of the American Mathematical Society, vol. 5 (1904), p. 477.

‡ Compare LINDBERG, *Mathematische Annalen*, vol. 59 (1904), p. 321.

$$(2) \quad F_1[\tilde{x}(a), \tilde{y}(a), \tilde{x}'(a), \tilde{y}'(a)] > 0 \text{ in } (A_1 A_2),$$

where $\tilde{x}' = d\tilde{x}/da$, $\tilde{y}' = d\tilde{y}/da$.

For simplicity, we suppose that the parameter a is the arc of the curve $\tilde{\mathcal{C}}$ measured from some fixed initial point.

Under these conditions it follows from the general existence theorems* for differential equations applied to the differential equation of the extremals† for the integral (1), that through every point $P(a)$ of the curve $\tilde{\mathcal{C}}$ one and but one extremal \mathcal{E}_a can be drawn which is tangent to $\tilde{\mathcal{C}}$ at P in such a manner that the positive tangents of the two curves coincide. For the parameter t on the extremal \mathcal{E}_a we may choose the arc of the extremal measured from the point P so that for every value of a the point P corresponds on \mathcal{E}_a to the value $t = 0$.

If we vary a , we thus obtain a set of extremals

$$(3) \quad x = \phi(t, a), \quad y = \psi(t, a),$$

for which the functions ϕ , ψ have the following properties:

1) The functions

$$\phi, \phi_t, \phi_a; \psi, \psi_t, \psi_a$$

are as functions of t and a of class C' in the domain

$$(4) \quad 0 \leq t \leq l, \quad A_1 \leq a \leq A_2,$$

where l is a sufficiently small positive quantity independent of a .‡

2) The functions ϕ , ψ satisfy the following initial conditions:

$$(5) \quad \begin{aligned} \phi(0, a) &= \tilde{x}(a), & \psi(0, a) &= \tilde{y}(a), \\ \phi_t(0, a) &= \tilde{x}'(a), & \psi_t(0, a) &= \tilde{y}'(a). \end{aligned}$$

From (5) we obtain by differentiation

$$(6) \quad \begin{aligned} \phi_a(0, a) &= \tilde{x}'(a), & \psi_a(0, a) &= \tilde{y}'(a), \\ \phi_{aa}(0, a) &= \tilde{x}''(a), & \psi_{aa}(0, a) &= \tilde{y}''(a). \end{aligned}$$

From these equations we derive for the Jacobian

$$\Delta(t, a) = \frac{\partial(\phi, \psi)}{\partial(t, a)}$$

* Compare BLISS, *Annals of Mathematics*, ser. 2, vol. 6 (1905), pp. 49-67.

† Compare KNESEER, *Lehrbuch der Variationsrechnung*, §§ 27, 29 and BOLZA, *Lectures on the Calculus of Variations*, § 25, b).

‡ Compare the corollary given by BLISS in the article on differential equations just referred to, p. 53, at the end of section 1.

the result :

$$(7) \quad \Delta(0, a) = 0, \quad \Delta_t(0, a) = \frac{1}{r} - \frac{1}{\bar{r}},$$

if we denote by $1/r$ and $1/\bar{r}$ the curvature at the point P of the curve \mathfrak{C}_a and of the curve $\tilde{\mathfrak{C}}$ respectively.

We make the *further assumption* that

$$\frac{1}{r} - \frac{1}{\bar{r}} \neq 0$$

along $\tilde{\mathfrak{C}}$, and in order to fix the ideas we suppose * that

$$(8) \quad \frac{1}{r} - \frac{1}{\bar{r}} > 0.$$

From this additional assumption it follows that two positive quantities $l_0 \leq l$ and m can be determined so that

$$(9) \quad \Delta(t, a) \geq tm$$

in the domain

$$(10) \quad 0 \leq t \leq l_0, \quad A_1 \leq a \leq A_2.$$

For if we define the function $\chi(t, a)$ for the domain (4) by the equations

$$\chi(t, a) = \begin{cases} \frac{\Delta(t, a)}{t}, & \text{when } t \neq 0, \\ \Delta_t(0, a), & \text{when } t = 0, \end{cases}$$

it is easily seen that $\chi(t, a)$ is continuous in the domain (4), and since moreover $\chi(0, a) > 0$ in (A_1, A_2) , it follows that a positive quantity $l_0 \leq l$ can be assigned such that $\chi(t, a) > 0$ in the domain (10). If we denote by m the minimum of $\chi(t, a)$ in the domain (10), we obtain (9).

§ 2. Proof that the set of extremals (3) furnishes a field.

We now choose two quantities a_1, a_2 so that

$$A_1 < a_1 < a_2 < A_2$$


and propose to prove that under the assumptions enumerated in § 1, a positive quantity $k \leq l_0$ can be assigned such that the equations (3) define a one-to-one correspondence between the rectangle

* In order that the curve $\tilde{\mathfrak{C}}$ may furnish a minimum for the integral (1) with respect to one-sided variations on the left of $\tilde{\mathfrak{C}}$, it is necessary that $1/r - 1/\bar{r} \geq 0$; compare BOLZA, *Lectures*, p. 194.

$$\mathbf{A}_\kappa: \quad 0 \leqq t \leqq \kappa, \quad a_1 \leqq a \leqq a_2$$

in the t, a -plane and its image \mathbf{S}_κ in the x, y -plane.

We suppose that it were not so; that is we suppose that, however small κ may be taken, there always exists in \mathbf{A}_κ at least one pair of distinct points whose images in the x, y -plane coincide. Reasoning then exactly as in the proof for the exist-



ence of a field which I have given in § 34 of my *Lectures* for the case where $\Delta(t, a) \neq 0$, we reach the result that under this hypothesis there would exist a point $\Pi (t = 0, a = \alpha)$ in the rectangle \mathbf{A}_κ such that every vicinity of Π contains at least one pair of distinct points of \mathbf{A}_κ whose images in the x, y -plane coincide.

We are going to prove that this leads to a contradiction with the inequality (9).

For this purpose we notice that our assumptions concerning the curve $\tilde{\mathcal{C}}$ imply* that $\tilde{x}'(\alpha), \tilde{y}'(\alpha)$ are not both zero; let $\tilde{x}'(\alpha) \neq 0$, or as we may write on account of (6),

$$\phi_a(0, \alpha) \neq 0.$$

We may then apply DINI's theorem on implicit functions to the function $\phi(t, a)$ and the point $t = 0, a = \alpha$. From this theorem it follows † that below any

* Compare the definition of "curve of class C'' " on p. 116 of my *Lectures*.

† Choose $d > 0$ so that $\phi(t, a)$ is of class C' and $\phi_a(t, a) \neq 0$ for $|t| \leqq d, |a - \alpha| \leqq d$. Let A be the maximum of $|\phi_t(t, a)|$, B the minimum of $|\phi_a(t, a)|$ in this domain. Choose $0 < d_0 < d$ and

$$\sigma < d_1 - d_0, \quad \rho < d_0, \quad \frac{\sigma}{2}, \frac{B\sigma}{2A}.$$

Compare PEANO-GENOCHI, *Differentialrechnung und Grundzüge der Integralrechnung*, pp. 138-141.

preassigned positive quantity δ two positive quantities ρ and σ can be determined having the following properties: If $P'(t', a')$ and $P''(t'', a'')$ be any two distinct points of the vicinity (ρ) of the point $\Pi(0, \alpha)$ for which

$$(11) \quad \phi(t', a') = \phi(t'', a''),$$

then in the first place $t'' \neq t'$ (say $t' < t''$) and in the second place the two points P', P'' can be joined by a curve representable in the form

$$a = a(t), \quad t' \leq t \leq t'',$$

such that

$$(12) \quad \phi[t, a(t)] = \phi(t', a') \text{ for } t' \leq t \leq t''.$$

The function $a(t)$ is of C' , and satisfies the inequality

$$|a(t) - a'| < \sigma \text{ for } t' \leq t \leq t''$$

and the initial conditions

$$(13) \quad a(t') = a', \quad a(t'') = a''.$$

Differentiating (12) we obtain

$$(14) \quad \phi_t[t, a(t)] + \phi_a[t, a(t)] a'(t) = 0.$$

On the other hand, it follows from the characteristic property of the point Π that there exists at least one pair of distinct points P', P'' in the domain

$$0 \leq t < \rho, \quad |a - \alpha| < \rho$$

for which not only (11) holds but at the same time

$$(15) \quad \psi(t', a') = \psi(t'', a'').$$

For such a pair of points the function $\psi[t, a(t)]$ is of class C' in (tt') and takes, according to (13) and (15), the same value for $t = t'$ and $t = t''$. Hence its derivative must vanish at least for one value $t = \tau$ between t' and t'' :

$$\psi_t(\tau, a(\tau)) + \psi_a[\tau, a(\tau)] a'(\tau) = 0.$$

Combining this equation with the equation derived from (14) by putting $t = \tau$, we obtain the result:

$$\Delta[\tau, a(\tau)] = 0.$$

But if we take ρ and σ sufficiently small, the point $t = \tau$, $a = a(\tau)$ lies in the domain (10); moreover, τ is positive since $0 \leq t' < \tau < t''$. Hence we have indeed reached a contradiction with the inequality (9), and therefore the statement enunciated at the beginning of this section is proved.

The image \mathcal{Q} of the boundary of the rectangle A_x is a continuous closed curve without multiple points. According to a theorem due to SCHÖNFLIES,* the point-set S_x is therefore identical with the interior of \mathcal{Q} together with the curve \mathcal{Q} itself.

THE UNIVERSITY OF CHICAGO,
February 27, 1907.

*Göttinger Nachrichten, 1899, p. 282; compare also OSGOOD, *ibid.*, 1900, p. 94; and BERNSTEIN, *ibid.*, 1900, p. 98.