

ON THE ZEROS OF THE FUNCTION $\beta(z)$ ASSOCIATED WITH THE GAMMA FUNCTION*

BY
T. H. GRONWALL

1. Nielsen † has raised the question whether the function $\beta(z)$ defined by

$$\beta(z) = \frac{1}{2} \left[\psi \left(\frac{1+z}{2} \right) - \psi \left(\frac{z}{2} \right) \right], \quad \left(\psi(z) = \frac{\Gamma'(z)}{\Gamma(z)} \right),$$

has any zeros, and has shown that there are no real zeros, and that the complex zeros, if any, must have their real part less than $-\frac{1}{2}$.

This question will be answered completely in the following, by showing that

The zeros of $\beta(z)$ are all complex, and their real part less than $-\frac{1}{2}$. For $n=1, 2, 3, \dots$, each of the infinite strips

$$-2n - \frac{1}{2} < \text{real part of } z < -2n + \frac{3}{2}$$

contains exactly two zeros, and for n sufficiently large, their asymptotic expression is

$$-2n + \frac{1}{2} + \frac{2 \log(8n+2)\pi}{(4n+1)\pi^2} \pm i \frac{\log(8n+2)\pi}{\pi} + \frac{\xi_n \pm i\eta_n}{2\pi n},$$

$$\xi_n^2 + \eta_n^2 < 1.$$

2. From the definition of $\beta(z)$ it follows at once that

$$(1) \quad \beta(z) = \sum_{n=0}^{\infty} \frac{(-1)^n}{z+n} = -\log 2 + \frac{1}{z} + \sum_{n=1}^{\infty} (-1)^n \left(\frac{1}{z+n} - \frac{1}{n} \right),$$

both series converging uniformly (and the second also absolutely) in any finite region in the z -plane to which the poles $z=0, -1, -2, \dots$ are exterior; and from (1) (or directly from the definition) we obtain the relation

$$(2) \quad \beta(z) + \beta(1-z) = \frac{\pi}{\sin \pi z}.$$

* Presented to the Society, September 5, 1916; received by the editors in April, 1925.

† N. Nielsen, *Handbuch der Theorie der Gammafunktion*, Leipzig, Teubner, 1906. See p. 101.

Let us begin by proving Nielsen's results. For z real and positive, the terms in (1) have alternating signs and decrease in absolute value, so that $\beta(z) > 0$. For $z = -m - \zeta$, where m is a positive integer or zero, and $0 < \zeta < 1$, we may write (1) in the form

$$(-1)^{m+1}\beta(-m-\zeta) = - \left[\frac{1}{1+\zeta} - \frac{1}{2+\zeta} + \dots + \frac{(-1)^m}{m+\zeta} \right] + \frac{1}{\zeta} + \frac{1}{1-\zeta} - \left[\frac{1}{2-\zeta} - \frac{1}{3-\zeta} + \dots \right];$$

the terms in brackets having alternating signs and decreasing in absolute value, we have

$$(-1)^{m+1}\beta(-m-\zeta) > \frac{1}{\zeta} + \frac{1}{1-\zeta} - \frac{1}{1+\zeta} - \frac{1}{2-\zeta} = \frac{1}{\zeta(1+\zeta)} + \frac{1}{(1-\zeta)(2-\zeta)} > 0.$$

Hence there are no real zeros. On the other hand, let $z = x + yi$, $y \neq 0$, be a complex zero; taking the imaginary part of (1) and dividing by y , we find

$$\frac{1}{x^2+y^2} - \frac{1}{(x+1)^2+y^2} + \left[\frac{1}{(x+2)^2+y^2} - \frac{1}{(x+3)^2+y^2} + \dots \right] = 0,$$

and assuming $x > -2$, the terms in the bracket have alternating signs and decrease in absolute value, whence

$$\frac{1}{x^2+y^2} - \frac{1}{(x+1)^2+y^2} < 0 \text{ or } x < -\frac{1}{2}.$$

3. It is convenient to determine first the zeros of $\beta(1-z)$ and then replace z by $1-z$. From the preceding results it follows that the zeros of $\beta(1-z)$ will be complex and have their real part greater than $\frac{3}{2}$. To obtain an asymptotic expression for $\beta(1-z)$, we observe that, for $\Re z > 0$ ($\Re z = x =$ real part of z), the expression (1) may be replaced by

$$\beta(z) = \int_0^\infty \frac{e^{-zu}}{1+e^{-u}} du,$$

whence, integrating twice by parts,

$$(3) \quad \beta(z) = \frac{1}{2z} + \frac{1}{4z^2} - \frac{1}{z^2} \int_0^\infty e^{-zu} \frac{e^{-u} - e^{-2u}}{(1+e^{-u})^3} du.$$

For $\Re z \geq 1$, we have

$$\left| \int_0^\infty e^{-su} \frac{e^{-u} - e^{-2u}}{(1+e^{-u})^3} du \right| < \int_0^\infty e^{-u} (e^{-u} - e^{-2u}) du = \frac{1}{6},$$

and (2) and (3) now give the asymptotic expression

$$(4) \quad \beta(1-z) = \frac{\pi}{\sin \pi z} - \frac{1}{2z} - \frac{h}{z^2}, \quad |h| < \frac{5}{12} \text{ for } \Re z \geq 1.$$

4. Let us consider first the zeros of the expression

$$(5) \quad \frac{\pi}{\sin \pi z} - \frac{1}{2z},$$

which constitutes the principal part of $\beta(1-z)$. Since $\sin \pi(x+yi) = \sin \pi x \operatorname{ch} \pi y + i \cos \pi x \operatorname{sh} \pi y$, a zero $z = x+yi$ of (5) implies the two equations

$$(6) \quad \sin \pi x \operatorname{ch} \pi y = 2\pi x,$$

$$(7) \quad \cos \pi x \operatorname{sh} \pi y = 2\pi y,$$

where, however, the solution $x=y=0$ must be discarded, since it corresponds to the pole $z=0$ of (5). It is evident that when $x+yi$ is a zero of (5), the same is true of $x-yi$, $-x+yi$ and $-x-yi$; we may therefore restrict the discussion of (6) and (7) to the case $x \geq 0$, $y \geq 0$. First assume $x=0$; since $y \geq 0$ in this case, and $(\operatorname{sh} \pi y)/\pi y$ increases steadily from 1 to ∞ as y increases from 0 to ∞ , (7) gives $y=y_0$, where y_0 is the unique positive root of

$$(8) \quad \frac{\operatorname{sh} \pi y_0}{\pi y_0} = 2.$$

Now assume $x > 0$; making $y=0$ in (6), we would obtain $(\sin \pi x)/\pi x = 2$, whereas the quotient to the left always lies between -1 and $+1$. We must consequently assume $x > 0$, $y > 0$, and (5) and (6) show that $\sin \pi x > 0$, $\cos \pi x > 0$, whence

$$(9) \quad x = 2n + \frac{1}{2} - \xi,$$

n a positive integer or zero, $0 < \xi < \frac{1}{2}$. Consider first the case $n=0$, whence $0 < x < \frac{1}{2}$; since $\pi x / \sin \pi x < \pi/2$ in this interval, (6) gives $\operatorname{ch} \pi y < \pi$, whence $\pi y < 1.812$, and we obtain from (7)

$$\cos \pi x > \frac{2 \times 1.812}{\operatorname{sh} 1.812} = 1.3336 > 1,$$

which is impossible.

Next, assume $n > 0$; the equations (6) and (7) become

$$(10) \quad \cos \pi \xi \operatorname{ch} \pi y = (4n+1)\pi - 2\pi \xi ,$$

$$(11) \quad \sin \pi \xi \operatorname{sh} \pi y = 2\pi y .$$

Since $\pi y / \operatorname{sh} \pi y$ is a decreasing function of y , equation (11) represents a curve in the rectangular coördinates ξ, y such that y decreases steadily from $+\infty$ to y_0 when ξ increases from 0 to $\frac{1}{2}$. On the curve (10), y is an extremum when

$$\cos^2 \pi \xi \operatorname{sh} \pi y \frac{dy}{d\xi} = [(4n+1)\pi - 2\pi \xi] \sin \pi \xi - 2 \cos \pi \xi = 0 ,$$

and writing this equation in the form $(4n+1)\pi - 2\pi \xi - 2 \cot \pi \xi = 0$, we see that it has a unique root ξ_0 in the interval $0 < \xi < \frac{1}{2}$, since the derivative of the left hand member is $2\pi \cot^2 \pi \xi > 0$. This root evidently corresponds to a minimum, since $y \rightarrow +\infty$ when $\xi \rightarrow \frac{1}{2}$ by (10), and we have

$$\sin \pi \xi_0 < \tan \pi \xi_0 = \frac{2}{(4n+1)\pi - 2\pi \xi_0} < \frac{1}{2n\pi} .$$

For $0 < \xi < \xi_0$, the value of y obtained from (10) decreases steadily from y_1 given by $\operatorname{ch} \pi y_1 = (4n+1)\pi$, to y_2 , given by $\cos \pi \xi_0 \operatorname{ch} \pi y_2 = (4n+1)\pi - 2\pi \xi_0$, while the value of y obtained from (11) decreases from $+\infty$ to y_3 , given by $\sin \pi \xi_0 \operatorname{sh} \pi y_3 = 2\pi y_3$. We shall now show that $y_3 > y_1$; it is evidently sufficient to prove that

$$\frac{\operatorname{sh} \pi y_3}{\pi y_3} > \frac{\operatorname{sh} \pi y_1}{\pi y_1} .$$

From $e^{\pi y_1} < e^{\pi y_1} + e^{-\pi y_1} = 2 \operatorname{ch} \pi y_1 = (8n+2)\pi$ it follows that $\pi y_1 < \log(8n+2)\pi$, and since $(\operatorname{sh} \pi y) / \pi y$ is a decreasing function,

$$\frac{\operatorname{sh} \pi y_1}{\pi y_1} < \frac{\operatorname{sh} [\log(8n+2)\pi]}{\log(8n+2)\pi} < \frac{\frac{1}{2} e^{\log(8n+2)\pi}}{\log(8n+2)\pi} = \frac{(4n+1)\pi}{\log(8n+2)\pi} .$$

On the other hand,

$$\frac{\operatorname{sh} \pi y_3}{\pi y_3} = \frac{2}{\sin \pi \xi_0} > 4n\pi ,$$

and since we have

$$4n\pi > \frac{(4n+1)\pi}{\log(8n+2)\pi}$$

for $n \geq 1$, the inequality in question is established, and it follows that the two curves (10) and (11) do not intersect in the interval $0 < \xi < \xi_0$. On the

contrary, in the interval $\xi_0 < \xi < \frac{1}{2}$ the y in (10) increases steadily from y_2 to $+\infty$, while the y in (11) decreases steadily from y_3 to y_0 , and since $y_3 > y_1 > y_2$, the two curves have a unique point of intersection in the interval considered.

Returning to (5), it is thus seen that the only zeros of this function are the following:

The two zeros $y_0i, -y_0i$;

In each of the strips $2n < \Re z < 2n + \frac{1}{2}$ ($n = 1, 2, 3, \dots$), two conjugate complex zeros;

In each of the strips $-2n - \frac{1}{2} < \Re z < -2n$ ($n = 1, 2, 3, \dots$), two conjugate complex zeros, equal to the preceding ones multiplied by -1 .

5. To find the distribution of the zeros of $\beta(1-z)$, we shall use the following theorem:

Let $f(z)$ and $g(z)$ be two functions, meromorphic inside a contour C , and holomorphic on C . When the inequality

$$|f(z) - g(z)| < |g(z)|$$

*is satisfied everywhere on C , then neither $f(z)$ nor $g(z)$ vanishes on C , and the difference between the number of zeros and the number of poles of $f(z)$ inside C equals the corresponding difference for $g(z)$.**

Let us apply this theorem to $f(z) = \beta(1-z)$ and $g(z) = (\pi/\sin \pi z) - (1/2z)$, C being the rectangle with vertices at $2n - \frac{1}{2} \pm bi$, $2n + \frac{3}{2} \pm bi$, where n is a positive integer, and b positive and very large. On the horizontal sides of the rectangle, we have $z = 2n - \frac{1}{2} + x \pm bi$, $0 \leq x \leq 2$, whence

$$\sin \pi z = -\cos \pi x \operatorname{ch} \pi b \pm i \sin \pi x \operatorname{sh} \pi b, \quad |\sin \pi z|^2 = \cos^2 \pi x + \operatorname{sh}^2 \pi b$$

and

$$\left| \frac{\pi}{\sin \pi z} \right| \leq \frac{\pi}{\operatorname{sh} \pi b} < 2\pi e^{-\pi b}.$$

* In the case when C is a circle and $f(z)$ and $g(z)$ have no poles in its interior, this theorem is due to E. Rouché, *Mémoire sur la série de Lagrange*, Journal de l'École Polytechnique, Cahier 39 (1862), pp. 193-224 (see Theorem III, p. 217). The theorem was rediscovered and generalized by A. Hurwitz, *Ueber die Nullstellen der Bessel'schen Function*, Mathematische Annalen, vol. 33 (1889), p. 246-266 (see p. 248). Incidentally, the proof is extremely simple: First, neither $f(z)$ nor $g(z)$ vanishes on C , since $|f(z) - g(z)| < |g(z)|$ yields the impossible inequalities $|f(z)| < 0$ for $g(z) = 0$ and $|g(z)| < |g(z)|$ for $f(z) = 0$. In the identity

$$\log f(z) = \log g(z) + \log [1 + (f(z) - g(z))/g(z)],$$

we perform the analytic continuation of both members along the contour C , described once in the positive sense. Then $\log f(z)$ will increase by $2\pi i$ times the difference between the number of zeros and the number of poles of $f(z)$ interior to C , the increase in $\log g(z)$ will be the corresponding expression for $g(z)$, and the third logarithm does not change, since, on account of the inequality $|f(z) - g(z)| < |g(z)|$, the point $\zeta = [f(z) - g(z)]/g(z)$ describes a closed path interior to the circle $|\zeta| < 1$ where $\log(1 + \zeta)$ is holomorphic.

Moreover, for $b > 2n + 2$,

$$\frac{1}{2|z|} > \frac{1}{4n-1+2x+2b} > \frac{1}{4b}, \quad \frac{1}{|z|^2} = \frac{1}{(2n-\frac{1}{2}+x)^2+b^2} < \frac{1}{b^2},$$

whence, for b sufficiently large and using (4),

$$\begin{aligned} \left| \frac{\pi}{\sin \pi z} - \frac{1}{2z} \right| &\geq \frac{1}{2|z|} - \left| \frac{\pi}{\sin \pi z} \right| > \frac{1}{4b} - 2\pi e^{-\pi b}, \\ \left| \beta(1-z) - \left(\frac{\pi}{\sin \pi z} - \frac{1}{2z} \right) \right| &< \frac{5}{12|z|^2} < \frac{5}{12b^2} < \frac{1}{4b} - 2\pi e^{-\pi b} \\ &< \left| \frac{\pi}{\sin \pi z} - \frac{1}{2z} \right|. \end{aligned}$$

On the vertical sides of the rectangle, we have $z = 2m - \frac{1}{2} + yi$, $m = n$ or $n + 1$ and $-b \leq y \leq b$, whence $\sin \pi z = -\operatorname{ch} \pi y$ and

$$\frac{\pi}{\sin \pi z} - \frac{1}{2z} = - \left(\frac{\pi}{\operatorname{ch} \pi y} + \frac{1}{4m-1+2yi} \right);$$

the real part of the second term to the right having the same sign as the first term, it follows that

$$\left| \frac{\pi}{\sin \pi z} - \frac{1}{2z} \right| > \left| \frac{1}{4m-1+2yi} \right| = \frac{1}{|2z|}.$$

Hence we see that

$$\left| \beta(1-z) - \left(\frac{\pi}{\sin \pi z} - \frac{1}{2z} \right) \right| < \frac{5}{12|z|^2} < \frac{1}{2|z|} < \left| \frac{\pi}{\sin \pi z} - \frac{1}{2z} \right|.$$

For b sufficiently large, the conditions of the theorem are thus fulfilled. According to paragraph 4, the function $(\pi/\sin \pi z) - (1/2z)$ has two zeros in our rectangle for b sufficiently large, viz., those with their real part between $2n$ and $2n + \frac{1}{2}$, and on the other hand, there are the two poles $2n$ and $2n + 1$. The function $\beta(1-z)$ has the same poles in the rectangle, and our theorem therefore shows that the rectangle contains exactly two zeros of $\beta(1-z)$. Letting b increase indefinitely, and replacing z by $1-z$, it is thus seen that for $n = 1, 2, 3, \dots$, each strip $-2n - \frac{1}{2} < \Re z < -2n + \frac{3}{2}$ contains two zeros of $\beta(z)$.

6. To obtain asymptotic expressions for the zeros, we begin with those of $(\pi/\sin \pi z) - (1/2z)$. As in paragraph 4, it is sufficient to consider the case $x > 0$, $y > 0$, and it follows from (10) that $\operatorname{ch} \pi y > 4n\pi$, so that y increases

indefinitely with n , and formula (11) shows that then $\lim \sin \pi \xi = 0$, whence $\lim \cos \pi \xi = 1$. By (10), we now have

$$\lim \frac{\operatorname{ch} \pi y}{(4n+1)\pi} = 1 = \lim \frac{e^{\pi y}}{2(4n+1)\pi},$$

$$\pi y = \log [(8n+2)\pi(1+\eta)], \quad \lim \eta = 0.$$

Substituting this value of πy in (11), it is seen that

$$\lim \frac{(4n+1)\pi}{2 \log (8n+2)\pi} \cdot \pi \xi = \lim \frac{\pi \xi}{\sin \pi \xi} \cdot \lim \frac{(4n+1)\pi}{2 \log (8n+2)\pi} \cdot \frac{2 \log [(8n+2)\pi(1+\eta)]}{\frac{1}{2} \left[(8n+2)\pi(1+\eta) - \frac{1}{(8n+2)\pi(1+\eta)} \right]}$$

$$= 1, \quad \pi \xi = \frac{2 \log (8n+2)\pi}{(4n+1)\pi} (1+\xi'), \quad \lim \xi' = 0.$$

Denoting, as usual, by $f(x) = O(g(x))$ the fact that two constants A and x_0 exist such that $|f(x)| < Ag(x)$ for $x > x_0$, we find by substituting the above values of πy and $\pi \xi$ in (10) and expanding $\cos \pi \xi$ in a power series, so that $\cos \pi \xi = 1 + O(\xi^2) = 1 + O((\log n)/n)^2$,

$$\left[1 + O\left(\left(\frac{\log n}{n}\right)^2\right) \right] \left[(4n+1)\pi(1+\eta) - \frac{1}{4(4n+1)\pi(1+\eta)} \right]$$

$$= (4n+1)\pi + O\left(\frac{\log n}{n}\right),$$

and it follows that

$$\eta = O\left(\left(\frac{\log n}{n}\right)^2\right).$$

Finally (11) gives, taking account of the order of magnitude of η ,

$$\left[\frac{2 \log (8n+2)\pi}{(4n+1)\pi} (1+\xi') + O\left(\left(\frac{\log n}{n}\right)^3\right) \right] \left\{ (4n+1)\pi \left[1 + O\left(\left(\frac{\log n}{n}\right)^2\right) \right] - \frac{1}{4(4n+1)\pi} \left[1 + O\left(\left(\frac{\log n}{n}\right)^2\right) \right] \right\}$$

$$= 2 \left[\log (8n+2)\pi + O\left(\left(\frac{\log n}{n}\right)^2\right) \right],$$

$$2 \log (8n+2)\pi \cdot \xi' = O\left(\frac{(\log n)^3}{n^2}\right),$$

whence

$$\xi' = O\left(\left(\frac{\log n}{n}\right)^2\right).$$

For the two zeros of (5) in the strip $2n < \Re z < 2n + \frac{1}{2}$, we consequently have the asymptotic expression

$$(12) \quad 2n + \frac{1}{2} - \frac{2 \log(8n+2)\pi}{(4n+1)\pi^2} \pm i \frac{\log(8n+2)\pi}{\pi} + O\left(\left(\frac{\log n}{n}\right)^2\right).$$

7. We shall now prove that for n sufficiently large, $\beta(1-z)$ has one zero in the neighborhood of each of the zeros (12) of its principal part (5). Consider the circle

$$(13) \quad z = 2n + \frac{1}{2} - \frac{2 \log(8n+2)\pi}{(4n+1)\pi^2} + i \frac{\log(8n+2)\pi}{\pi} + \frac{\xi + \eta i}{2n\pi},$$

$$\xi^2 + \eta^2 < 1;$$

by (12), this circle contains in its interior a single zero of (5) when n is sufficiently large. On the circumference $\xi^2 + \eta^2 = 1$, we have

$$\begin{aligned} \sin \pi z &= \frac{1}{2i} \left[\frac{i}{(8n+2)\pi} e^{-\frac{2i \log(8n+2)\pi}{(4n+1)\pi} + \frac{\xi i - \eta}{2n}} \right. \\ &\quad \left. + i(8n+2)\pi e^{\frac{2i \log(8n+2)\pi}{(4n+1)\pi} + \frac{\eta - \xi i}{2n}} \right] \\ &= (4n+1)\pi \left[1 + \frac{2i \log(8n+2)\pi}{(4n+1)\pi} + \frac{\eta - \xi i}{2n} + O\left(\left(\frac{\log n}{n}\right)^2\right) \right], \\ \frac{\pi}{\sin \pi z} &= \frac{1}{4n+1} \left[1 - \frac{2i \log(8n+2)\pi}{(4n+1)\pi} + \frac{\xi i - \eta}{2n} + O\left(\left(\frac{\log n}{n}\right)^2\right) \right], \\ \frac{1}{2z} &= \frac{1}{(4n+1) \left[1 + \frac{2i \log(8n+2)\pi}{(4n+1)\pi} + O\left(\frac{\log n}{n^2}\right) \right]} \\ &= \frac{1}{4n+1} \left[1 - \frac{2i \log(8n+2)\pi}{(4n+1)\pi} + O\left(\frac{\log n}{n^2}\right) \right], \end{aligned}$$

whence

$$\frac{\pi}{\sin \pi z} - \frac{1}{2z} = \frac{\xi i - \eta}{2n(4n+1)} + O\left(\frac{(\log n)^2}{n^3}\right) = \frac{\xi i - \eta}{8n^2} + O\left(\frac{(\log n)^2}{n^3}\right),$$

and since $\xi^2 + \eta^2 = 1$,

$$(14) \quad \left| \frac{\pi}{\sin \pi z} - \frac{1}{2z} \right| = \frac{1}{8n^2} + O\left(\frac{(\log n)^2}{n^3}\right).$$

On the other hand, we have on the same circumference, according to (4),

$$\left| \beta(1-z) - \left(\frac{\pi}{\sin \pi z} - \frac{1}{2z} \right) \right| < \frac{5}{12|z|^2} = \frac{5}{48n^2} + O\left(\frac{\log n}{n^3}\right),$$

and by comparison with (14)

$$\left| \beta(1-z) - \left(\frac{\pi}{\sin \pi z} - \frac{1}{2z} \right) \right| < \left| \frac{\pi}{\sin \pi z} - \frac{1}{2z} \right|$$

on the circumference $\xi^2 + \eta^2 = 1$ for n sufficiently large. Applying the theorem in paragraph 5, we see that the circle $\xi^2 + \eta^2 < 1$ contains a single zero of $\beta(1-z)$, and the same is evidently true of the circle obtained by changing the sign of i in (13). We have thus obtained asymptotic expressions for the two zeros of $\beta(1-z)$ contained in the strip $2n - \frac{1}{2} < \Re z < 2n + \frac{3}{2}$, and replacing z by $1-z$, we finally arrive at the result stated in paragraph 1.

It is clear that closer approximations to the zeros may be obtained by using, instead of (5), the general asymptotic expansion

$$\beta(z) = \frac{1}{2z} + \sum_{\nu=1}^n (-1)^{\nu-1} \frac{2^{2\nu}-1}{2\nu} B_{\nu} \cdot \frac{1}{z^{2\nu}} \\ + \frac{1}{z^{2n}} \int_0^{\infty} e^{-zu} \frac{d^n}{du^n} \left(\frac{1}{1+e^{-u}} \right) du,$$

valid for $\Re z > 0$, where B_{ν} are the Bernoulli numbers.

NEW YORK, N. Y.
