

SINGULAR CASE OF PAIRS OF BILINEAR, QUADRATIC, OR HERMITIAN FORMS*

BY

L. E. DICKSON

The main object of this paper is a treatment of the equivalence of pairs of bilinear forms in the singular case by a purely rational method. The problem was first discussed by Kronecker,† who employed the irrational canonical form due to Weierstrass for the auxiliary non-singular case, instead of the rational canonical form (13) employed here. It is then a simple matter to deduce in Parts II and III the criteria for the equivalence of pairs of symmetric or Hermitian bilinear forms, or quadratic or Hermitian forms, in the singular case.

I. PAIRS OF BILINEAR FORMS IN THE SINGULAR CASE

Let ϕ and ψ be bilinear forms in the same variables $x_1, \dots, x_r, y_1, \dots, y_s$ with coefficients in a field F , such that ϕ and ψ are not equivalent in F to forms both of which involve fewer than r variables x_i or fewer than s variables y_i . We shall treat the singular case in which either $r \neq s$, or else $r = s$ and the determinant of $f = u\phi + v\psi$ is zero identically in u and v . If $r \neq s$, we take $r > s$ without loss of generality, since we may interchange the letters x and y .

Write f_i, ϕ_i, ψ_i for the partial derivatives of f, ϕ, ψ with respect to x_i . The linear functions f_1, \dots, f_r of y_1, \dots, y_s are linearly dependent. Hence there exist homogeneous polynomials A_i in u and v of the same degree with coefficients in F , such that the A_i are not all zero identically and such that $\sum A_i f_i \equiv 0$ identically in u, v, y_1, \dots, y_s .

LEMMA 1. *The degree of A_1, \dots, A_r in u and v is not changed if we apply to f non-singular linear transformations on the x 's and the y 's separately whose coefficients are independent of u and v .*

*Presented to the Society, December 26, 1924; received by the editors in July, 1926.

† Sitzungsberichte der Akademie der Wissenschaften, Berlin, 1890, pp. 1225–37. His brief paper was elaborated by Muth, *Elementarteiler*, 1899, pp. 93–133. Some minor errata in Muth's work were reproduced in the account in the *Encyclopédie des Sciences Mathématiques*, tome I, vol. II, 1911, pp. 412–25.

Evidently each A_i is unchanged if we transform the y 's alone. Next, let f become g when the x 's are expressed linearly in terms of new variables X_j . Then

$$X_j = \sum_{i=1}^r a_{ji}x_i, \quad f_i = \sum_{j=1}^r \frac{\partial g}{\partial X_j} \frac{\partial X_j}{\partial x_i} = \sum_{j=1}^r a_{ji}g_j,$$

and $\sum B_j g_j \equiv 0$, where $B_j = \sum_i a_{ji}A_i$ is of the same degree as the A_i .

Of all relations $\sum A_i f_i \equiv 0$, select one

$$(1) \quad \sum_{i=1}^r A_i f_i \equiv \sum_{\alpha=0}^m (-1)^\alpha C_\alpha u^\alpha v^{m-\alpha} \equiv 0$$

in which the A_i are not all zero identically and are of minimal degree m in u and v , while

$$(2) \quad C_\alpha = \sum_{i=1}^r c_{\alpha i} f_i \quad (\alpha = 0, 1, \dots, m),$$

where the $c_{\alpha i}$ are numbers not all zero of the field F and are independent of u and v . By Lemma 1, m is an invariant of the pair ϕ and ψ under all non-singular linear transformations on the x 's and the y 's separately.

LEMMA 2. f_1, \dots, f_r are linearly independent in F .

For, if $\sum t_i f_i \equiv 0$, where the t_i are in F and are not all zero, the insertion of $f_i = u\phi_i + v\psi_i$ gives

$$\sum t_i \phi_i \equiv 0, \quad \sum t_i \psi_i \equiv 0.$$

Let $t_e \neq 0$ and apply the non-singular transformation

$$x_e = t_e X_e, \quad x_j = X_j + t_j X_e \quad (j = 1, \dots, r; j \neq e).$$

Then $\phi = \sum x_i \phi_i$ becomes

$$\sum_{i \neq e} X_i \phi_i + X_e \sum_{i=1}^r t_i \phi_i,$$

which lacks X_e since the final sum is zero. Likewise, ψ lacks X_e . But this contradicts our initial assumption.

COROLLARY. *The invariant m exceeds zero.*

For, if $m=0$, (1) becomes $C_0=0$ and (2) implies that every $c_{\alpha i}$ is zero, contrary to hypothesis.

LEMMA 3. C_0, \dots, C_m are linearly independent in F .

Suppose that a_0, \dots, a_m are numbers, not all zero, of F such that

$$(3) \quad \sum_{\alpha=0}^m a_\alpha C_\alpha \equiv 0.$$

Denote the first and second sums in (1) by Q and R . Write

$$(4) \quad P = \sum_{\alpha=0}^m (-1)^\alpha a_\alpha u^{m-\alpha} v^\alpha.$$

Each PA_i is a polynomial in u and v of degree $2m$; let k_i be the coefficient of $u^m v^m$. The terms $\sum k_i u^m v^m f_i$ of PQ are obtained from the terms $\sum a_\alpha C_\alpha u^m v^m$ of PR by replacing C_α by its value (2). Hence (3) implies $\sum k_i f_i \equiv 0$. By Lemma 2, each $k_i = 0$. Hence $PQ \equiv 0$ may be written in the form

$$(5) \quad v^{m+1} \sum f_i g_i + u^{m+1} \sum f_i h_i \equiv 0 \quad (i = 1, \dots, r),$$

in which the g_i and h_i are polynomials of degree $m-1$ in u and v . If they were all zero identically, every PA_i and hence every A_i would be zero identically. Replacing f_i by its value $u\phi_i + v\psi_i$, we see that the first sum in (5) is of degree $\leq m$ in u , while the second sum is evidently of degree $> m$ in u . Hence each sum is zero identically, and one of them gives a linear relation between f_1, \dots, f_r the degree in u and v of whose coefficients is $m-1$, while those coefficients (g_i or h_i) are not all zero identically. But this contradicts the definition of m as the minimal degree.

By Lemmas 2 and 3, the matrix of the coefficients in (2) has a non-vanishing determinant of order $m+1$. Hence we can annex further rows, also composed of numbers of F , and obtain a square matrix

$$(c_{\alpha\beta}) \quad (\alpha = 0, 1, \dots, r-1; \beta = 1, \dots, r)$$

whose determinant is not zero. Let the non-singular transformation

$$(6) \quad x_\beta = \sum_{\alpha=0}^{r-1} c_{\alpha\beta} X_\alpha \quad (\beta = 1, \dots, r),$$

whose coefficients are independent of u and v , replace f, ϕ, ψ by F, Φ, Ψ . The partial derivatives of the latter with respect to X_i are denoted by F_i, Φ_i, Ψ_i . Then

$$(7) \quad F_\alpha = \sum_{\beta=1}^r \frac{\partial f}{\partial x_\beta} \frac{\partial x_\beta}{\partial X_\alpha} = \sum_{\beta=1}^r c_{\alpha\beta} f_\beta = C_\alpha,$$

by (2). By (1),

$$\sum (-1)^\alpha F_\alpha u^\alpha v^{m-\alpha} \equiv 0.$$

Replacing F_α by $u\Phi_\alpha + v\Psi_\alpha$, and equating to zero the coefficients of v^{m+1} , uv^m , u^2v^{m-1} , \dots , u^mv , u^{m+1} , we get

$$(8) \quad \Psi_0 \equiv 0, \Psi_1 \equiv \Phi_0, \Psi_2 \equiv \Phi_1, \dots, \Psi_m \equiv \Phi_{m-1}, \Phi_m \equiv 0.$$

Hence

$$F_0 \equiv u\Psi_1, F_1 \equiv u\Psi_2 + v\Psi_1, \dots, F_{m-1} \equiv u\Psi_m + v\Psi_{m-1}, F_m \equiv v\Psi_m.$$

Multiply the first k equations by v^{k-1} , $-uv^{k-2}$, \dots , $(-u)^{k-2}v$, $(-u)^{k-1}$, respectively, and add; we get

$$(9) \quad -(-u)^k \Psi_k \equiv \sum_{i=0}^{k-1} F_i (-u)^i v^{k-i-1} \quad (k = 1, \dots, m).$$

If Ψ_1, \dots, Ψ_m satisfy a linear relation whose coefficients are in the field and are not all zero, multiply it by u^m and apply (9). We get a linear relation between F_0, \dots, F_{m-1} whose coefficients are polynomials, not all identically zero, of degree $m-1$ in u and v . Inserting the values (7) and applying (2), we get a similar linear relation between f_1, \dots, f_r , contrary to the definition of m . Hence Ψ_1, \dots, Ψ_m are linearly independent in the field. We may therefore choose linear functions Y_{m+1}, \dots, Y_s of y_1, \dots, y_s with coefficients in the field F such that the Ψ and Y are linearly independent in F . Then

$$(10) \quad \eta_\alpha = \Psi_\alpha \quad (\alpha = 1, \dots, m), \quad \eta_t = Y_t \quad (t = m+1, \dots, s)$$

define a non-singular linear transformation on y_1, \dots, y_s with coefficients in F . Transformation (6) replaces $u\phi + v\psi$ by

$$u\Phi + v\Psi = \sum_{i=0}^{r-1} u\Phi_i X_i + \sum_{i=0}^{r-1} v\Psi_i X_i.$$

For $i=0, \dots, m$, change i to $\alpha-1$ in the first sum, and i to α in the second. For $i > m$, change i to p . Then by (8) and (10),

$$u\Phi + v\Psi = \sum_{\alpha=1}^m (uX_{\alpha-1} + vX_\alpha)\eta_\alpha + \sum_{p=m+1}^{r-1} (u\Phi_p + v\Psi_p)X_p.$$

Here Φ_p and Ψ_p are linear in y_1, \dots, y_s and hence in η_1, \dots, η_s by (10):

$$\Phi_p = \sum_{\alpha=1}^s a_{p\alpha} \eta_\alpha, \quad \Psi_p = \sum_{\alpha=1}^s b_{p\alpha} \eta_\alpha \quad (p = m+1, \dots, r-1),$$

where the a 's and b 's are numbers in F . We have

$$\begin{aligned}
 (11) \quad u\Phi + v\Psi &= u\left(X_0 + \sum_p a_{p1}X_p\right)\eta_1 + u\sum_{\alpha=2}^m X_{\alpha-1}\eta_\alpha \\
 &+ u\sum_{\alpha=2}^m \sum_p b_{p\alpha-1}X_p\eta_\alpha + v\sum_{\alpha=1}^m \left(X_\alpha + \sum_p b_{p\alpha}X_p\right)\eta_\alpha + \rho + \sigma,
 \end{aligned}$$

where the summations with respect to p are from $m+1$ to $r-1$, and $\rho = u(l_2\eta_2 + \dots + l_m\eta_m)$, the l 's being linear in the X_p ($p > m$), while σ is a bilinear form in the X_p, η_q ($p > m, q > m$). In deriving (11), the terms involving η_α ($\alpha > m$) which arose from Ψ_p were carried to σ ; the terms involving η_α ($\alpha = 2, \dots, m$) which arose from Φ_p were carried to ρ , while those involving η_α ($\alpha > m$) were carried to σ . Finally we added the terms in the third sum in the second member and subtracted the same from ρ . To (11) apply the non-singular transformation

$$\begin{aligned}
 \xi_0 &= X_0 + \sum_p a_{p1}X_p, \quad \xi_\alpha = X_\alpha + \sum_p b_{p\alpha}X_p, \quad \xi_p = X_p \\
 &(\alpha = 1, \dots, m; \quad p = m+1, \dots, r-1).
 \end{aligned}$$

We get (12) in which $u\Phi + v\Psi$ is written for σ .

THEOREM 1. *If ϕ and ψ are bilinear forms in $x_1, \dots, x_r, y_1, \dots, y_s$ with coefficients in a field F , and $r \geq s$, and when $r = s$ the determinant $|u\phi + v\psi|$ is zero identically in u and v , there exist non-singular linear transformations on the x 's and on the y 's separately with coefficients in F which reduce $u\phi + v\psi$ to*

$$(12) \quad S = \sum_{\alpha=1}^m (u\xi_{\alpha-1} + v\xi_\alpha)\eta_\alpha + u\sum_{\alpha=2}^m l_\alpha\eta_\alpha + u\Phi + v\Psi,$$

where l_2, \dots, l_m are linear forms in $\xi_{m+1}, \dots, \xi_{r-1}$, and Φ and Ψ are bilinear forms in $\xi_{m+1}, \dots, \xi_{r-1}, \eta_{m+1}, \dots, \eta_s$, all with coefficients in F .

If $r < s$, $f = u\phi + v\psi$ can be reduced to a form derived from (12) by interchanging the letters ξ and η , since the form derived from f by interchanging the letters x and y can be reduced to (12).

Given a pair of bilinear forms λ and μ with coefficients in F such that $u\lambda + v\mu$ is of rank R , we can select independent linear combinations ϕ and ψ of λ and μ with coefficients in F such that the greatest common divisor of all R -rowed determinants of $u\phi + v\psi$ is not divisible by u , whence ψ is of rank R . In the corresponding form (12), let τ be the rank of $u\Phi + v\Psi$.

LEMMA 4. *The greatest common divisor of all τ -rowed determinants of $u\Phi + v\Psi$ is not divisible by u .*

In the matrix M of (12), the first m columns form the matrix M_1 of the coefficients of ξ_0, \dots, ξ_{r-1} in

$$\frac{\partial S}{\partial \eta_1} = u\xi_0 + v\xi_1, \quad \frac{\partial S}{\partial \eta_2} = u\xi_1 + v\xi_2 + u\eta_2, \quad \dots, \quad \frac{\partial S}{\partial \eta_m} = u\xi_{m-1} + v\xi_m + u\eta_m.$$

Since these are evidently linearly independent, M_1 is of rank m . The last $s-m$ columns of M form the matrix M_2 of the coefficients of ξ_0, \dots, ξ_{r-1} in the partial derivatives of $u\Phi + v\Psi$ with respect to $\eta_{m+1}, \dots, \eta_s$. Hence by the definition of τ , the rank of M_2 is τ . Thus not every $(m+\tau)$ -rowed determinant of M is zero. But every determinant with more than $m+\tau$ columns is zero, since it contains more than τ columns of M_2 . Hence the rank of M is $m+\tau=R$. Every non-vanishing $(m+\tau)$ -rowed determinant of M contains exactly τ columns of M_2 and is a linear function of τ -rowed determinants of M_2 and hence of $u\Phi + v\Psi$. If all τ -rowed determinants of the latter were divisible by u , the same would be true of all determinants with $m+\tau=R$ columns of M and hence of S , contrary to the remark before Lemma 4.

First, let $u\Phi + v\Psi$ fall under the non-singular case, so that it has the same number of variables in the two series and its determinant is not zero identically in u and v . By Lemma 4, this determinant is not divisible by u , whence $|\Psi| \neq 0$. Then there exist non-singular transformations on the variables of the two series separately, with coefficients in F , which reduce Φ and Ψ to the respective forms*

$$(13) \quad \rho = \sum_k \left(\sum_{i=1}^{a_k-1} x_{ki} y_{k\ i+1} + x_{ka_k} \sum_{i=1}^{a_k} b_{ki} y_{ki} \right), \quad \sigma = \sum_k \sum_{i=1}^{a_k} x_{ki} y_{ki}.$$

The transformations employed leave unaltered each variable occurring in the first sum in (12) and replace the second sum by $u \sum t_\alpha \eta_\alpha$, where t_α is a linear function of the x_{ki} . Hence if $r \geq s$, ϕ and ψ may be reduced rationally to

$$(14) \quad J = \sum_{\alpha=1}^m \xi_{\alpha-1} \eta_\alpha + \sum_{\alpha=2}^m \sum_k \sum_{i=1}^{a_k} d_{k\ i\ \alpha} x_{ki} \eta_\alpha + \rho, \quad K = \sum_{\alpha=1}^m \xi_\alpha \eta_\alpha + \sigma.$$

Second, let $u\Phi + v\Psi$ fall under the singular case. We apply to it our initial discussion of $u\phi + v\psi$ and conclude that it can be transformed rationally

*Dickson, *Modern Algebraic Theories*, Chicago, Sanborn, 1926, p. 116, Theorem IV. A direct verification follows from the fact that the invariant factors other than 1 of $\rho - \lambda\sigma$ are

$$\lambda^{a_k} - \sum_{i=1}^{a_k} b_{ki} \lambda^{i-1} \quad (k = 1, 2, \dots),$$

which may be identified with any assigned polynomials.

ally into a form $R_1 + uP_1 + f_1$ of type (12) or a form $S_1 + uQ_1 + f_1$ obtained from it by interchanging the letters ξ and η , where

$$f_1 = u\Phi_1 + v\Psi_1, \quad R_1 = \sum_{\alpha=1}^{m_1} (u\xi_{1\alpha-1} + v\xi_{1\alpha})\eta_{1\alpha}, \quad P_1 = \sum_{\alpha=2}^{m_1} P_{1\alpha}\eta_{1\alpha},$$

$$S_1 = \sum_{\alpha=1}^{n_1} (u\eta_{1\alpha-1} + v\eta_{1\alpha})\xi_{1\alpha}, \quad Q_1 = \sum_{\alpha=2}^{n_1} q_{1\alpha}\xi_{1\alpha}.$$

If f_1 falls under the non-singular case, Φ_1 and Ψ_1 can be transformed rationally into forms of type (13). If f_1 falls under the singular case, another repetition of the initial discussion shows that f_1 can be transformed rationally into $R_2 + uP_2 + f_2$ or $S_2 + uQ_2 + f_2$, where f_2, R_2, \dots, Q_2 are of the type f_1, R_1, \dots, Q_1 with the first subscripts 1 changed to 2.

This process must terminate, since the numbers of variables in Φ, Φ_1, \dots form a series of decreasing positive integers.

To prove by induction that $\sum l_\alpha \eta_\alpha$ (in (12)), P_1 or Q_1, P_2 or Q_2 , etc., can be removed by a linear transformation with coefficients in F which does not alter the sum of the remaining terms, we assume that all but $\sum l_\alpha \eta_\alpha$ have been so removed and shall prove that we can then remove it also.

We first show that R_1, S_1, R_2, \dots can all be written as terms of $u\rho + v\sigma$ in (13) by employing new values of k . To do this for R_1 , write

$$\xi_{1\alpha} = x_{k\alpha+1} \quad (\alpha = 0, 1, \dots, m_1), \quad \eta_{1\alpha} = y_{k\alpha+1} \quad (\alpha = 1, \dots, m_1),$$

and $m_1 = a_k - 1$. Then R_1 becomes

$$(15) \quad u \sum_{i=1}^{a_k-1} x_{ki} y_{k\ i+1} + v \sum_{i=1}^{a_k} x_{ki} y_{ki}$$

for $y_{k1} = 0$; the coefficients of u and v are the terms of (13) for the present value of k and with every $b_{ki} = 0$. In S_1 write

$$\xi_{1\alpha} = x_{k\ n_1+1-\alpha} \quad (\alpha = 1, \dots, n_1), \quad \eta_{1\alpha} = y_{k\ n_1+1-\alpha} \quad (\alpha = 0, 1, \dots, n_1),$$

and $n_1 = a_k - 1$. Then S_1 becomes (15) for $x_{ka_k} = 0$.

Whether $u\Phi + v\Psi$ falls under the singular or non-singular case, we may therefore reduce the initial pair ϕ, ψ with $r \geq s$ to a pair (14) such that, for certain values of k , either x_{ka_k} is suppressed or y_{k1} is suppressed and every b_{ki} replaced by zero in (13) and the second sum in (14).

It remains to prove that we may take every d to be zero in (14). We shall first prove that we may replace $d_{ki\alpha}$ by zero when $i < a_k$. We proceed by induction on α , keeping k fixed. For a given integer n such that $2 \leq n \leq m$,

assume that $d_{k\alpha} = 0$ for $i < a_k, \alpha = 2, \dots, n-1$. Apply the transformation*

$$\xi_n = X_n + \sum_{i=2}^{a_k} t_{ki} x_{ki}, \quad y_{ki} = Y_{ki} - t_{ki} \eta_n \quad (i = 2, \dots, a_k)$$

which leaves unaltered the variables not occurring in the left members. The increment to K is zero, while that to J is

$$\eta_{n+1} \sum_{i=2}^{a_k} t_{ki} x_{ki} - \sum_{i=1}^{a_k-1} t_{k \ i+1} x_{k \ i+1} \eta_n + c x_{ka_k} \eta_n,$$

where the first sum is to be omitted if $n = m$. Taking

$$t_{k \ i+1} = d_{kin} \quad (i = 1, \dots, a_k - 1),$$

we obtain forms of type (14) in which $d_{k\alpha} = 0$ for $i < a_k, \alpha = 2, \dots, n$. Hence the induction is complete and we now have

$$(16) \quad J = \sum_{\alpha=1}^m \xi_{\alpha-1} \eta_{\alpha} + \sum_{\alpha=2}^m \sum_k e_{k\alpha} x_{ka_k} \eta_{\alpha} + \rho, \quad K = \sum_{\alpha=1}^m \xi_{\alpha} \eta_{\alpha} + \sigma.$$

We shall prove that we may replace each $e_{k\alpha}$ by zero. No proof is needed when k is such that x_{ka_k} is to be suppressed. We proceed by induction on α , keeping k fixed. For a given positive integer n ($n < m$), assume that $e_{k\alpha} = 0$ for $\alpha = m, m-1, \dots, n+2$.

First, let $n < a_k$ and apply the transformation

$$(17) \quad \begin{aligned} \xi_{n-j} &= X_{n-j} + t x_{ka_{k-j}} & (j = 0, \dots, n), \\ y_{ka_{k-j}} &= Y_{ka_{k-j}} - t \eta_{n-j} & (j = 0, \dots, n-1), \end{aligned}$$

which leaves unaltered the remaining variables. The increment to K is zero, while that to J cancels to

$$(18) \quad t x_{ka_k} \eta_{n+1} - t x_{ka_k} \sum_{j=0}^{n-1} b_{ka_{k-j}} \eta_{n-j}.$$

Taking $t = -e_{k \ n+1}$, we obtain (16) with $e_{k\alpha} = 0$ ($\alpha = m, \dots, n+1$), but with a new term $c x_{ka_k} \eta_1$. The latter is removed by the transformation $\xi_0 = \Xi_0 - c x_{ka_k}$.

Second, let k be one of the values for which y_{k1} is not to be suppressed from (13) and let $n \geq a_k$. Apply transformation (17) with $j = 0, \dots, a_k - 1$ in both parts. Then K is unaltered, while the increment to J cancels to

*But let the summation extend only to $a_k - 1$ if k is such that x_{ka_k} is to be suppressed from (13) and (14). Then the text holds with $c = 0$.

(18) in which now j takes the values $0, \dots, a_k - 1$. We proceed as in the first case.

In either case we obtain forms of type (16) in which now $e_{k\alpha} = 0$ ($\alpha = m, m - 1, \dots, n + 1$). By induction on n , we obtain (16) in which every $e_{k\alpha} = 0$.

Third, let k be a value for which y_{k1} is to be suppressed from (16), while every $b_{ki} = 0$. Then $m < a_k$, whence $n < a_k$ and the second case is excluded. In the proof, we write

$$S = uJ + vK, \Sigma_\alpha = \frac{\partial S}{\partial \xi_\alpha} \quad (\alpha = 0, \dots, m), \quad S_{ki} = \frac{\partial S}{\partial x_{ki}} \quad (i = 1, \dots, a_k).$$

By (16) and (13),

$$(19) \quad \Sigma_0 = u\eta_1, \quad \Sigma_\alpha = u\eta_{\alpha+1} + v\eta_\alpha \quad (\alpha = 1, \dots, m - 1), \quad \Sigma_m = v\eta_m,$$

the solution of which is

$$(20) \quad u^{\alpha+1}\eta_{\alpha+1} = \sum_{j=0}^{\alpha} (-1)^j u^{\alpha-j} v^j \Sigma_{\alpha-j} \quad (\alpha = 0, \dots, m - 1).$$

Eliminating the resulting value of η_m from the last equation (19), we get

$$(21) \quad u^{m\Sigma_m} = \sum_{i=1}^m (-1)^{i-1} u^{m-i} v^i \Sigma_{m-i}.$$

Next,

$$(22) \quad S_{ki} = uy_{k\ i+1} + vy_{ki} \quad (i < a_k), \quad S_{ka_k} = u \sum_{\alpha=2}^m e_{k\alpha} \eta_\alpha + vy_{ka_k}.$$

The former with y_{k1} suppressed gives

$$(23) \quad u^{i+1}y_{k\ i+2} = \sum_{j=0}^i (-1)^j u^{i-j} v^j S_{k\ i+1-j} \quad (i = 0, \dots, a_k - 2).$$

By (20) and (23), $u^t \eta_t$ and $u^t y_{k\ t+1}$ are linear functions of the Σ 's and S 's whose coefficients are polynomials in u and v of degree $t - 1$. Multiply the last equation (22) by u^{m-1} . If $m \geq a_k$, we get

$$(24) \quad u^{m-1}S_{ka_k} = \sum_{\alpha=2}^m e_{k\alpha} u^{m-\alpha} u^\alpha \eta_\alpha + vu^{m-a_k} u^{a_k-1} y_{ka_k}.$$

By the remark just made, this yields a linear relation between the first derivatives of S whose coefficients are polynomials in u and v of degree $m - 1$. But the S on the left does not occur on the right. This contradicts the definition of m .

This completes the proof that we may take every d to be zero in (14). In stating the following theorem, we keep separate the forms R_i and S_i previously combined with $u\rho+v\sigma$. For each i only one of the pair R_i and S_i is a component of the canonical form. We shall change the notation of the variables occurring in a component S_j .

THEOREM 2. *Let ϕ and ψ be bilinear forms with coefficients in a field F such that ψ has* the same rank as $f=u\phi+v\psi$. In the singular case, f can be reduced by non-singular linear transformations with coefficients in F to $S=\sum R_i + \sum S_j + u\rho + v\sigma$, where ρ and σ are given by (13), and*

$$(25) \quad R_i = \sum_{\alpha=1}^{m_i} (u\xi_{i\alpha-1} + v\xi_{i\alpha})\eta_{i\alpha}, \quad S_j = \sum_{\alpha=1}^{n_j} (uY_{j\alpha-1} + vY_{j\alpha})X_{j\alpha}.$$

No variable appears in two of the forms $R_1, R_2, \dots, S_1, S_2, \dots, u\rho + v\sigma$, while all their variables are linearly independent in F .

We shall now prove that the invariant factors other than 1 of the matrix M of S coincide with those of $u\rho+v\sigma$. The determinant D of order d of the latter is not zero identically in u and v . Write $t=\sum m_i + \sum n_j$. Then the rank of M will be shown to be $d+t$. In M , the coefficient of a product $\xi\eta$ is placed at the intersection of the row $[\xi]$ corresponding to ξ and the column $[\eta]$ corresponding to η . Similarly for XY .

First, delete the rows $[\xi_{i,m_i}]$ corresponding to the ξ 's with maximal second subscripts in the R_i , and the columns $[Y_{j,n_j}]$ corresponding to Y 's with maximal second subscripts in the S_j . We obtain a square matrix which (apart from elements of D) has every diagonal element equal to u , and has zeros as elements to the right (left) of these u 's in the case of $R_i(S_j)$, whence its determinant is $u^t D$.

Second, delete the rows $[\xi_{i,0}]$ and the columns $[Y_{j,0}]$. We obtain a square matrix which (apart from elements of D) has every diagonal element equal to v , and has zeros as elements to the left (right) of these v 's in the case of $R_i(S_j)$, whence its determinant is $v^t D$.

Hence the greatest common divisor of all $(d+t)$ -rowed determinants of M is D . This yields†

LEMMA 5. *The invariant factors other than unity of S coincide with those of $u\rho+v\sigma$.*

Hence ρ and σ are uniquely determined by the invariant factors of $u\phi+v\psi$.

*We saw that this property may be secured by passing to suitably chosen linear combinations of ϕ and ψ with coefficients in F .

†See the writer's *Modern Algebraic Theories*, pp. 96, 97.

The variables ξ , X , x of the first series in S are linearly independent; likewise for the variables η , Y , y of the second series (Theorem 2). Write $W = u\rho + v\sigma$. No variable is common to two of the R_i , S_j , W . Hence a linear function $[S]$ of the first derivatives of $S = W + \sum R_i + \sum S_j$ with respect to the variables of either series is a sum of linear functions $[W]$, $[R_1]$, $[R_2]$, \dots , $[S_1]$, \dots of the first derivatives of W , R_1 , R_2 , \dots with respect to their own variables only. Also, $[S] = 0$ implies $[W] = 0$, $[R_1] = 0$, \dots . Since $|W|$ is not identically zero, the vanishing of $[W]$ implies that its coefficients are all zero.

If we suppress the first subscript i from m_i and the variables in R_i , and write Σ_α for $\partial R_i / \partial \xi_\alpha$, we have (19)-(21). Since $\Sigma_0, \dots, \Sigma_{m-1}$ are linearly independent, every linear relation between the Σ 's is obtained by multiplying (21) by a function of u and v .

The first derivatives of R_i with respect to the $\eta_{i\alpha}$ are evidently linearly independent. Since R_i becomes S_i by replacing ξ , η , m by Y , X , n , we have

LEMMA 6. *As a complete set of independent linear relations between the first derivatives of S with respect to the variables ξ , X , x of the first series we may take*

$$(26) \quad u^{m_i} \frac{\partial S}{\partial \xi_{i m_i}} = \sum_{j=1}^{m_i} (-1)^{i-1} u^{m_i-j} v^j \frac{\partial S}{\partial \xi_{i m_i-j}},$$

where i takes the same values as the subscript of R_i . Here m_i is the minimal degree in u and v of the coefficients in any linear relation between the first derivatives of S with respect to the variables ξ_i of R_i . Similarly, n_j is the minimal degree in u and v of the coefficients in any linear relation between the first derivatives of S with respect to the variables Y_j of S_j and these relations of minimal degrees form a complete set of independent linear relations between the first derivatives of S with respect to the variables η , Y , y of the second series.

These minimal degrees m_i and n_j uniquely determine the R_i and S_j . By Lemma 1, they have the same value for $u\phi + v\psi$ as for the canonical form S . They are called the *minimal numbers* of $u\phi + v\psi$ or of S .

By Lemmas 5 and 6, and the remarks just following them, we conclude that two pairs of bilinear forms ϕ , ψ and λ , μ in F , each satisfying the initial assumption and having the property about rank in Theorem 2, have the same canonical form and hence are equivalent in F if and only if they have the same minimal numbers and the same invariant factors.

Our next step is to prove this result when the two restrictions on the forms are removed.

Let f and g be any bilinear forms with coefficients in F in the singular case. After supplying zero coefficients, if necessary, we may assume that

there is the same number n of variables X_i as variables Y_i . If τ is the rank of $h=uf+vg$, there are exactly $n-\tau$ independent linear relations between the first derivatives of h with respect to X_1, \dots, X_n (or Y_1, \dots, Y_n). Let them be chosen so that their coefficients are of the least possible degrees $M_1, \dots, M_{n-\tau}$ (or $N_1, \dots, N_{n-\tau}$) in u and v . Let the M 's be arranged in order of increasing magnitude; likewise the N 's. The resulting numbers M_i and N_i are called the *minimal numbers* of the pair f, g .

For example, let $f=X_1Y_3+X_3Y_1, g=X_2Y_3+X_3Y_2$. Then h has the matrix

$$\begin{pmatrix} 0 & 0 & u \\ 0 & 0 & v \\ u & v & 0 \end{pmatrix}$$

of rank 2. Hence $vh_1-uh_2 \equiv 0$ is the single linear relation between the first partial derivatives h_1, h_2, h_3 , of h with respect to X_1, X_2, X_3 . The minimal numbers are $M_1=1, N_1=1$. Since the two-rowed determinants include $-u^2$ and $-v^2$, their greatest common divisor is 1. Hence the matrix has no invariant factor other than 1.

If the first ρ of the M 's and the first σ of the N 's are zero, while the others are positive, we can transform f, g into a pair ϕ, ψ involving only $r=n-\rho$ variables x_i and only $s=n-\sigma$ variables y_i (Lemma 2). Then ϕ, ψ satisfy our initial assumption that they cannot be transformed rationally into a pair of forms both of which involve fewer than r variables x_i or fewer than s variables y_i .

Since the M_i and N_i are invariant under every linear transformation on the X 's or on the Y 's (Lemma 1), the positive ones of M_i and N_i coincide with the above m_i and n_i of ϕ and ψ .

Introduce the linear combinations

$$(27) \quad \phi' = a\phi + b\psi, \quad \psi' = c\phi + d\psi \quad (ad - bc = 1).$$

Write $\chi = u\phi + v\psi, \chi' = u'\phi' + v'\psi'$. Then $\chi = \chi'$ if

$$(28) \quad u' = du - cv, \quad v' = -bu + av.$$

Before Lemma 4, we chose a, b, c, d so that the greatest common divisor of all R -rowed determinants of χ' is not divisible by u' , where R is the rank of χ and hence of ψ' .

Let λ, μ be another pair of bilinear forms in $x_1, \dots, x_r, y_1, \dots, y_s$ with coefficients in F , having the same minimal numbers and same invariant factors as the pair ϕ, ψ . Using the same a, b, c, d as in (27), write

$$\lambda' = a\lambda + b\mu, \quad \mu' = c\lambda + d\mu, \quad \nu = u\lambda + v\mu, \quad \nu' = u'\lambda' + v'\mu'.$$

Since χ' and ν' are transformed into χ and ν by (28), they have the same minimal numbers and same invariant factors, and in particular the same rank R and the same greatest common divisor of their R -rowed determinants. Hence also μ' is of rank R .

Theorem 2 and the remarks following Lemmas 5 and 6 show that χ' and ν' have the same canonical form and hence are equivalent in F . Thus the pairs ϕ, ψ and λ, μ are equivalent in F . This completes the proof of the fundamental

THEOREM 3. *In the singular case, two pairs of bilinear forms in the same $n+n$ variables with coefficients in any field F are equivalent in F if and only if they have the same invariant factors and the same minimal numbers.*

The canonical pair shows that these may be given any assigned values such that each invariant factor divides the next.

II. PAIRS OF SYMMETRIC OR HERMITIAN BILINEAR FORMS IN THE SINGULAR CASE

In order to treat simultaneously symmetric and Hermitian bilinear forms, we employ \bar{a} to denote a or the conjugate imaginary to a in the respective cases. Consider the form

$$(29) \quad f = \sum_{i,j=1}^n a_{ij}x_iy_j, \quad a_{ji} = \bar{a}_{ij},$$

whose matrix $A = (a_{ij})$ has the property $A' = \bar{A}$. Then

$$\frac{\partial f}{\partial x_i} = \sum_{j=1}^n a_{ij}y_j, \quad \frac{\partial f}{\partial y_i} = \sum_{j=1}^n a_{ji}x_j = \sum_{j=1}^n \bar{a}_{ij}x_j.$$

If

$$(30) \quad \sum_{i=1}^n c_i \frac{\partial f}{\partial x_i} \equiv 0,$$

where the c_i are independent of the x 's and y 's, then

$$(31) \quad \sum_{i=1}^n c_i a_{ij} = 0 \quad (j = 1, \dots, n),$$

whence $\sum \bar{c}_i \partial f / \partial y_i \equiv 0$. Take $f = u\phi + v\psi$. Hence each minimal number M_i of ϕ, ψ coincides with a minimal number N_i and vice versa.

Consider two pairs of forms ϕ, ψ and λ, μ whose n -rowed matrices A, B, C, D have the property $M' = \bar{M}$. Let $u\phi + v\psi$ and $u\lambda + v\mu$ have the same minimal numbers M_i (and hence the same N_i) and the same invariant factors. Then the pairs are equivalent by Theorem 3, so that there exist non-singular n -rowed matrices P and Q for which

$$C = PAQ, \quad D = PBQ.$$

Hence* there exists a non-singular n -rowed matrix E such that

$$C = E' A \tilde{E}, \quad D = E' B \tilde{E},$$

whence the linear transformation on the x 's with the matrix E and that on the y 's with the matrix \tilde{E} replace ϕ by λ and ψ by μ . Since this argument may be reversed, we have

THEOREM 4. *In the singular case, two pairs of symmetric (or Hermitian) bilinear forms are equivalent under a non-singular linear transformation on the x 's and the cogredient (or conjugate) transformation on the y 's if and only if they have the same invariant factors and the same minimal numbers M_i (which here coincide with the N_i).*

In seeking a canonical pair for such a pair ϕ, ψ , note that the number of the M_i which have the value zero is the number of the x 's (or y 's) which can be removed by linear transformation. Each M_i which exceeds zero will be denoted by m_i ; let their number be p . In the notations of Theorem 2, a canonical form is

$$C = \sum_{i=1}^p T_i + u\gamma + v\delta, \quad T_i = R_i + S_i,$$

with $n_i = m_i$ in S_i , while $u\gamma + v\delta$ is symmetric (or Hermitian), has a determinant not zero identically in u and v , and has the same invariant factors as $u\phi + v\psi$. Then the latter is equivalent to C in the sense of Theorem 4.

To emphasize the fact that each T_i is symmetric (or Hermitian), write x_α for ξ_α , $x_{m_i+\alpha}$ for X_α , y_α for Y_α , and $y_{m_i+\alpha}$ for η_α .

THEOREM 5. *In the singular case, any pair of symmetric (or Hermitian) bilinear forms can be reduced by a non-singular linear transformation on the x 's and the cogredient (or conjugate) transformation on the y 's to a canonical pair which are the coefficients of u and v in $\sum T_i + u\gamma + v\delta$, where*

$$T_i = \sum_{\alpha=1}^{m_i} [(ux_{i\alpha-1} + vx_{i\alpha})y_{i\ m_i+\alpha} + (uy_{i\ \alpha-1} + vy_{i\alpha})x_{i\ m_i+\alpha}]$$

is symmetric and Hermitian in its $2m_i+1$ pairs of variables, while γ, δ is a canonical pair† for the non-singular case.

**Modern Algebraic Theories*, pp. 122-3.

†For the quadratic case, we may employ Weierstrass's canonical pair, *Modern Algebraic Theories*, p. 131. For the Hermitian case, we may employ the corresponding pair, Logsdon, *American Journal of Mathematics*, vol. 44 (1922), p. 254. Or, without separating the two cases we may employ the writer's pair, *Modern Algebraic Theories*, pp. 126-8. In all of these cases, γ and δ have any prescribed invariant factors (or elementary divisors).

III. PAIRS OF QUADRATIC OR HERMITIAN FORMS IN THE SINGULAR CASE

Let $f = u\phi + v\psi$ be of type (29). To it corresponds

$$(32) \quad h = \sum_{i,j=1}^n a_{ij}x_i\tilde{x}_j, \quad a_{ji} = \tilde{a}_{ij},$$

which is a quadratic or Hermitian form according as \tilde{a} is identified with a or its conjugate imaginary, and likewise for \tilde{x} . In the respective cases, write $g=2$ or 1 . Then

$$\frac{\partial h}{\partial x_i} = g \sum_{j=1}^n a_{ij}\tilde{x}_j, \quad \sum_{i=1}^n c_i \frac{\partial h}{\partial x_i} = g \sum_{j=1}^n \left(\sum_{i=1}^n c_i a_{ij} \right) \tilde{x}_j = 0,$$

by (31). Hence any linear relation (30) between the first derivatives of f with respect to x_1, \dots, x_n , whose coefficients c_i are polynomials in u and v , implies a linear relation with the same coefficients c_i between the first derivatives of h , and conversely. Hence the minimal numbers $M_i = N_i$ of f coincide with those of h . By* Theorem 4, we have

THEOREM 6. *In the singular case, two pairs of quadratic or Hermitian forms are equivalent if and only if they have the same invariant factors and the same minimal numbers M_i .*

From Theorem 5 we obtain a canonical pair of Hermitian forms by identifying each y with the conjugate imaginary to x , and a canonical pair of quadratic forms by identifying each y with x . The latter pair is

$$(33) \quad \gamma + 2 \sum_{i=1}^p \sum_{\alpha=1}^{m_i} x_i x_{i-\alpha} x_{m_i+\alpha}, \quad \delta + 2 \sum_{i=1}^p \sum_{\alpha=1}^{m_i} x_{i\alpha} x_{i-m_i+\alpha},$$

where γ, δ is the canonical pair in the non-singular case (last foot-note in II). In particular, there exists a pair of quadratic or Hermitian forms with any prescribed invariant factors and any prescribed minimal numbers.

An example of (33) is given by $2x_0x_2, 2x_1x_2$. We have

$$v \frac{\partial f}{\partial x_0} - u \frac{\partial f}{\partial x_1} \equiv 0, \quad f = 2ux_0x_2 + 2vx_1x_2.$$

Here there is a single minimal number $M_1=1$ and no invariant factor.

*And the corollaries on pp. 66 and 68 of *Modern Algebraic Theories*.