

SOME ASYMPTOTIC FORMULAS IN THE THEORY OF NUMBERS⁽¹⁾

BY
ECKFORD COHEN

1. Introduction. It is the aim of this paper to generalize, unify, and refine a number of related but scattered results in asymptotic number theory. Special cases of the main theorems of the paper yield estimates due to Cesàro [2], Feller and Tornier [10], Kanold [12], Rényi [13], and the present author [4]. The method of proof is elementary and is essentially a variant of a method used in an earlier note [7].

Clearly, every positive integer n has a unique representation of the form

$$(1.1) \quad n = d^2e, \quad e \text{ square-free, } d > 0.$$

If in this relation we place $d = Q(n)$, then $Q^2(n)$ is the largest square divisor of n . Let α denote an arbitrary non-negative real number and let S be any nonvacuous subset of the set J of positive integers. In Theorem 2.1 we obtain approximations to the sum

$$(1.2) \quad B_\alpha(x, S) \equiv \sum_{n \leq x \cdot Q(n) \in S} \left(\frac{n}{Q^2(n)} \right)^\alpha, \quad x \geq 1,$$

with numerous special results deduced as corollaries.

We shall say that a divisor d of n is unitary, written $d \parallel n$ or $d * \delta = n$, if $d\delta = n$, $(d, \delta) = 1$. In §3 we consider the sum $B'_\alpha(x, S)$ arising from (1.2) if, in the summation, $Q^2(n)$ is required to be a unitary divisor of n . The main result is contained in Theorem 3.1.

Let the distinct prime factors of n be denoted p_1, \dots, p_r , and place

$$(1.3) \quad n = p_1^{e_1} \cdots p_r^{e_r} \quad (r = 0 \text{ if } n = 1).$$

The set of integers n for which $e_i \geq 2$ ($i = 1, \dots, r$), that is, the integers with no simple prime factors, will be denoted by L . Evidently, n is uniquely representable in the form

$$(1.4) \quad n = de, \quad d \in L, \quad e \text{ square-free, } (d, e) = 1.$$

The factor d defined by (1.4) is the largest divisor $Q^*(n)$ of n contained in L . If T is a nonvacuous subset of L , then one may define, analogous to (1.2),

Presented to the Society, March 29, 1961; received by the editors August 21, 1962.

(1) This research was supported in part by the National Science Foundation (G-19012).

$$(1.5) \quad B_\alpha^*(x, T) = \sum_{n \leq x; Q^*(n) \in T} \left(\frac{n}{Q^*(n)} \right)^\alpha.$$

An estimate for $B_\alpha^*(x, T)$ is proved in Theorem 4.1.

Known results which are deduced as special cases of the theorems discussed above are listed as follows: Corollaries 2.1.2., 2.3.2, 2.4.2, 2.7.2, 2.8.2, 3.2.4, 4.1.1, 4.2.1, 4.2.2. It will be observed that in the case $\alpha = 0$, the sums $B_\alpha(x, S)$, $B'_\alpha(x, S)$, and $B_\alpha^*(x, T)$ become enumerative functions of certain sequences of integers. This special case is given particular emphasis in each of the three problems considered.

We confine our description of specific results of the paper to a single type that has recently stimulated interest [1; 11; 13; 16]. Referring to (1.3), let us place $\omega(n) = r$ and $\Omega(n) = e_1 + \dots + e_r$. Further, place

$$(1.6) \quad \Delta(n) = \Omega(n) - \omega(n) = \Delta(Q^*(n)).$$

Rényi has proved [13,(3)] that if d_m^* is the asymptotic density of the sequence of integers n for which $\Delta(n) = m$, then d_m^* exists for each integer $m \geq 0$, and

$$(1.7) \quad \sum_{m=0}^\infty d_m^* z^m = \prod_p \left(1 - \frac{1}{p} \right) \left(1 + \frac{1}{p-z} \right), \quad |z| < 2,$$

where the product is extended over the primes p . A new proof of this result is given in §4.

We also prove two analogues of (1.7). In particular, let d_m denote the density of the set of integers n such that $\Delta(Q^2(n)) = m$; then, by §2, each d_m exists and

$$(1.8) \quad \sum_{m=0}^\infty d_m z^m = \prod_p \left(1 - \frac{1}{p^2} \right) \left(1 + \frac{z}{p^2 - z^2} \right), \quad |z| < 2.$$

Moreover, it is shown (§3) that if d'_m represents the density of the set of n for which $\Delta(Q^2(n)) = m$, where $Q^2(n) \parallel n$, then

$$(1.9) \quad \sum_{m=0}^\infty d'_m z^m = \prod_p \left(1 - \frac{1}{p} \right) \left(1 + \frac{1}{p} + \frac{z}{p^2 - z^2} \right), \quad |z| < 2.$$

REMARK 1.1. With $z = 1$ in (1.8) we note that

$$(1.10) \quad \sum_{m=0}^\infty d_m = 1;$$

by (1.7) the corresponding result for d_m^* is also valid (Rényi [13]).

2. **Problem I. Estimates for $B_\alpha(x, S)$.** Let $\Gamma_\alpha(n, S)$ be defined by

$$(2.1) \quad \Gamma_\alpha(n, S) = \begin{cases} \left(\frac{n}{Q^2(n)} \right)^\alpha & \text{if } Q(n) \in S, \\ 0 & \text{if } Q(n) \notin S. \end{cases}$$

In particular, $\Gamma_0(n, S)$ is the characteristic function of the set of integers such that $Q(n) \in S$.

With $\mu(n)$ denoting the Möbius function, we have the following relation which is basic for this section.

LEMMA 2.1.

$$(2.2) \quad \Gamma_\alpha(n, S) = \sum_{d^2 e = n; d \in S} e^\alpha \mu^2(e).$$

This formula is merely a restatement of the uniqueness of the representation (1.1). The following known estimate is also required.

LEMMA 2.2 [4, LEMMA 4.2, $r = 1$]. *If $s \geq 0$, then for $x \geq 1$,*

$$(2.3) \quad \mathfrak{U}_s(x) \equiv \sum_{n \leq x} n^s \mu^2(n) = \frac{6}{\pi^2} \left(\frac{x^{s+1}}{s+1} \right) + O(x^{s+1/2}).$$

Define $\zeta_s(s)$ to be the sum of the series,

$$(2.4) \quad \zeta_s(s) \equiv \sum_{n=1; n \in S}^{\infty} \frac{1}{n^s},$$

for those sequences S and real numbers s for which the series converges. Note that $\zeta_J(s)$ is the zeta-function $\zeta(s)$, provided $s > 1$.

We are now ready to prove the main theorem of this section.

THEOREM 2.1. *If $\alpha \geq 0$, then for $x \geq 2$,*

$$(2.5) \quad B_\alpha(x, S) = \frac{6}{\pi^2} \left(\frac{x^{\alpha+1}}{\alpha+1} \right) \zeta_S(2\alpha+2) + O((x^{\alpha+1/2} R_\alpha(x, S))),$$

uniformly in S , where

$$(2.6) \quad R_\alpha(x, S) = \sum_{n \leq \sqrt{x}; n \in S} n^{-2\alpha-1}$$

provided this sum is nonvacuous; otherwise $R_\alpha(x, S) = 1$.

REMARK 2.1. Evidently, $R_0(x, S) = O(1)$ for those S for which $\zeta_S(1)$ is finite. Moreover, for arbitrary S ,

$$(2.7) \quad R_\alpha(x, S) = \begin{cases} O(1) & \text{if } \alpha > 0, \\ O(\log x) & \text{if } \alpha = 0. \end{cases}$$

In particular, the estimate, $R_\alpha(x, S) = O(\log x)$, is always valid.

Proof. By (1.2), (2.1), (2.3), and Lemma 2.1,

$$(2.8) \quad B_\alpha(x, S) = \sum_{n \leq x} \Gamma_\alpha(n, S) = \sum_{d^2 e \leq x; d \in S} e^\alpha \mu^2(e) = \sum_{n \leq \sqrt{x}; n \in S} \mathfrak{U}_\alpha \left(\frac{x}{n^2} \right).$$

Then by Lemma 2.2

$$\begin{aligned}
 B_\alpha(x, S) &= \frac{6}{\pi^2} \left(\frac{x^{\alpha+1}}{\alpha+1} \right) \sum_{n \leq \sqrt{x}; n \in S} n^{-2\alpha-2} + O(x^{\alpha+1/2} R_\alpha(x, S)) \\
 &= \frac{6}{\pi^2} \left(\frac{x^{\alpha+1}}{\alpha+1} \right) \zeta_S(2\alpha+2) + O\left(x^{\alpha+1} \sum_{n > \sqrt{x}} n^{-2\alpha-2}\right) + O(x^{\alpha+1/2} R_\alpha(x, S)).
 \end{aligned}$$

But the summation in the middle term is $O(x^{-\alpha-1/2})$ and therefore (2.5) results.

We now specialize the theorem to some special sets of integers S . Let H denote the set of integral squares.

COROLLARY 2.1. *If $\alpha \geq 0$, then*

$$(2.9) \quad B_\alpha(x, H) = \frac{6}{\pi^2} \left(\frac{x^{\alpha+1}}{\alpha+1} \right) \zeta(4\alpha+4) + O(x^{\alpha+1/2}).$$

Place $B_0(x, S) = B(x, S)$ for all S . Since $\zeta(4) = \pi^4/90$, one obtains

COROLLARY 2.1.1 ($\alpha = 0$).

$$(2.10) \quad B(x, H) = \frac{\pi^2 x}{15} + O(\sqrt{x}).$$

COROLLARY 2.1.2 (CESÀRO [2, §17]). *The asymptotic density of the integers n for which $Q(n)$ is a square is $\pi^2/15$.*

Suppose that S_d is the set of all n divisible by the fixed positive integer d .

COROLLARY 2.2. *If $\alpha > 0$, then*

$$(2.11) \quad B_\alpha(x, S_d) = \frac{6}{\pi^2} \left(\frac{x^{\alpha+1}}{\alpha+1} \right) \frac{\zeta(2\alpha+2)}{d^{2\alpha+2}} + O(x^{\alpha+1/2}).$$

Noting that $S_1 = J$, we have

COROLLARY 2.2.1 ($d = 1$). *If $\alpha > 0$, then*

$$(2.12) \quad B_\alpha(x, J) = \frac{6}{\pi^2} \left(\frac{x^{\alpha+1}}{\alpha+1} \right) \zeta(2\alpha+2) + O(x^{\alpha+1/2}).$$

We say that an arithmetical function $f(n)$ has average order $E(x)$ if its summatory function $F(x) \sim xE(x)$ as $x \rightarrow \infty$. Note that $B_\alpha(x, J)$ is the summatory function of $\Gamma_\alpha(n, J)$.

COROLLARY 2.2.2. *If $\alpha > 0$, then the α th power of the divisor of n conjugate to the greatest square divisor of n has average order $(6x^\alpha/\pi^2)(\zeta(2\alpha+2)/(\alpha+1))$.*

COROLLARY 2.2.3 ($d = \alpha = 1$). *The conjugate divisor of the greatest square divisor of n has average order $\pi^2 x/30$.*

Next we consider the case in which S consists of a single positive integer r , $S \equiv r$.

COROLLARY 2.2. For all $\alpha \geq 0$,

$$(2.13) \quad B_\alpha(x, r) = \frac{6}{\pi^2} \left(\frac{x^{\alpha+1}}{\alpha+1} \right) r^{-2\alpha-2} + O\left(\frac{x^{\alpha+1/2}}{r^{2\alpha+1}} \right),$$

uniformly in $r \leq \sqrt{x}$.

COROLLARY 2.3.1 ($\alpha = 0$).

$$(2.14) \quad B(x, r) = \frac{6x}{\pi^2 r^2} + O(\sqrt{x}).$$

COROLLARY 2.3.2 (KANOLD [12, (13)]; ALSO CF. SCHERK [14, (3)]). The asymptotic density of the set of integers n with $Q(n) > r$ is $1 - (6/\pi^2) \sum_{a \leq r} a^{-2}$.

Let D denote the sequence of square-free integers. Since the generating function of D , $\sum_{n=1}^{\infty} \mu^2(n)/n^s = \zeta(s)/\zeta(2s)$, $s > 1$, we have

COROLLARY 2.4. For all $\alpha \geq 0$, $x \geq 2$,

$$(2.15) \quad B_\alpha(x, D) = \frac{6}{\pi^2} \left(\frac{x^{\alpha+1}}{\alpha+1} \right) \frac{\zeta(2\alpha+2)}{\zeta(4\alpha+4)} + \begin{cases} O(x^{\alpha+1/2}) & \text{if } \alpha > 0, \\ O(\sqrt{x} \log x) & \text{if } \alpha = 0. \end{cases}$$

COROLLARY 2.4.1 ($\alpha = 0$). If $x \geq 2$, then

$$(2.16) \quad B(x, D) = \frac{90x}{\pi^4} + O(\sqrt{x} \log x).$$

COROLLARY 2.4.2 (CESÀRO [2, §21]). The asymptotic density of the set of those n for which $Q(n)$ is square-free is $90/\pi^4$.

Let P denote the set of the primes. Since

$$(2.17) \quad \sum_{p \leq x} \frac{1}{p} = O(\log \log x), \quad x \geq 3,$$

we have

COROLLARY 2.5. If $x \geq 3$, then

$$(2.18) \quad B_\alpha(x, P) = \frac{6}{\pi^2} \left(\frac{x^{\alpha+1}}{\alpha+1} \right) \sum_p \frac{1}{p^{2\alpha+2}} + \begin{cases} O(x^{1/2+\alpha}) & \text{if } \alpha > 0, \\ O(\sqrt{x} \log \log x) & \text{if } \alpha = 0. \end{cases}$$

COROLLARY 2.5.1 ($\alpha = 0$). The asymptotic density of the integers for which $Q(n)$ is prime is equal to

$$\frac{6}{\pi^2} \sum_p \frac{1}{p^2}.$$

Let $\{a_1, \dots, a_s\}$ denote an ordered set of s fixed positive integers, $s \geq 0$, and let $N = N_s(a)$ denote the set of all n of the form $n = p_1^{a_1} \dots p_s^{a_s}$, where the p_i are

distinct primes. In case $s = 0$, it is understood that $N = (1)$. Then by (2.17) and the above theorem,

COROLLARY 2.6. If $x \geq 3$,

$$(2.19) \quad B_\alpha(x, N) = \frac{6}{\pi^2} \left(\frac{x^{\alpha+1}}{\alpha+1} \right) \zeta_N(2\alpha+2) + \begin{cases} O(x^{\alpha+1/2}) & \text{if } \alpha > 0, \\ O(\sqrt{x}(\log \log x)^t) & \text{if } \alpha = 0, \end{cases}$$

where $t = t(a)$ denotes the number of 1's in the set, a_1, \dots, a_s .

COROLLARY 2.6.1 ($\alpha=0$). The asymptotic density of the integers n for which $Q(n) \in N_s(a)$ is given by

$$(2.20) \quad \delta_s(a) = \frac{6}{\pi^2} \sum_{p_1, \dots, p_s} \frac{1}{(p_1^{a_1} \cdots p_s^{a_s})^2},$$

where the summation is over all s -tuples of distinct primes, p_1, \dots, p_s , for which the numbers, $p_1^{a_1} \cdots p_s^{a_s}$ are distinct; if $s = 0$, the sum is understood to have the value 1.

Note that Corollary 2.6.1 reduces to Corollary 2.5.1 in case $s = 1, a_1 = 1$. We now prove (1.8) as a consequence of (2.20).

COROLLARY 2.6.2. If d_m is defined as in the introduction, then for $|z| < 2$,

$$(2.21) \quad \sum_{m=0}^{\infty} d_m z^m = \frac{6}{\pi^2} \prod_p \left(1 + \frac{z}{p^2 - z^2} \right).$$

Proof. By definition of d_m and (2.20)

$$d_m = \frac{6}{\pi^2} \sum_{p_1, \dots, p_s; s \geq 0} \sum_{a_1, \dots, a_s} \frac{1}{(p_1^{a_1} \cdots p_s^{a_s})^2},$$

where for each set of s distinct primes, p_1, \dots, p_s , the numbers a_1, \dots, a_s range over all s -tuples of positive integers such that $(2a_1 - 1) + \dots + (2a_s - 1) = m$. Equivalently,

$$d_m = \frac{6}{\pi^2} \sum_{p_1, \dots, p_s; s \geq 0} \frac{1}{p_1 \cdots p_s} \sum_{b_1, \dots, b_s; b \text{ odd}} \frac{1}{p_1^{b_1} \cdots p_s^{b_s}},$$

where the numbers b_1, \dots, b_s , for each $s \geq 0$, range over all s -tuples of odd positive integers such that $b_1 + \dots + b_s = m$. It then follows that

$$\sum_{m=0}^{\infty} d_m z^m = \frac{6}{\pi^2} \prod_p \left(1 + \frac{1}{p} \left(\frac{z}{p} + \frac{z^3}{p^3} + \frac{z^5}{p^5} + \dots \right) \right),$$

from which (2.21) results.

The formula (1.8) follows from (2.21) by virtue of the relation $\zeta(2) = 6/\pi^2$ and the Euler product representation of $\zeta(s)$.

Let E denote the set of all n such that $\omega(n)$ is even, that is, those n with an even number of distinct prime factors.

COROLLARY 2.7. *If $x \geq 2$, then*

$$(2.22) \quad B_\alpha(x, E) = \left(\frac{3\zeta(2\alpha + 2)C_{\alpha+1}}{\pi^2(\alpha + 1)} \right) x^{\alpha+1} + \begin{cases} O(x^{\alpha+1/2}) & \text{if } \alpha > 0, \\ O(\sqrt{x \log x}) & \text{if } \alpha = 0, \end{cases}$$

where

$$(2.23) \quad C_t = 1 + \prod_p \left(1 - \frac{2}{p^{2t}} \right), \quad t > \frac{1}{2}.$$

Proof. It remains to evaluate $\zeta_E(s)$, $s > 1$. We have

$$\begin{aligned} \zeta_E(s) &= \sum_{n=1; \omega(n) \text{ even}}^{\infty} \frac{1}{n^s} = \frac{1}{2} \left\{ \sum_{n=1}^{\infty} \frac{1}{n^s} + \sum_{n=1}^{\infty} \frac{(-1)^{\omega(n)}}{n^s} \right\} \\ &= \frac{1}{2} \left\{ \zeta(s) + \prod_p \left(1 - \sum_{i=1}^{\infty} p^{-is} \right) \right\} = \frac{1}{2} \left\{ \zeta(s) + \prod_p \left(2 - \frac{1}{1 - p^{-s}} \right) \right\} \\ &= \frac{1}{2} \left\{ \zeta(s) + \prod_p \left(\frac{1}{1 - p^{-s}} \right) \cdot \prod_p (1 - 2p^{-s}) \right\} = \frac{\zeta(s)}{2} \left\{ 1 + \prod_p \left(1 - \frac{2}{p^s} \right) \right\}. \end{aligned}$$

This proves (2.22).

COROLLARY 2.7.1 ($\alpha = 0$). *If $x \geq 2$, then*

$$(2.24) \quad B(x, E) = \frac{Cx}{2} + O(\sqrt{x \log x}),$$

where $C = C_1$, as defined by (2.23).

COROLLARY 2.7.2 (FELLER AND TORNIER [10, §12]; ALSO CF. SCHOENBERG [15, §10]). *The asymptotic density of the sequence of n for which $\omega(Q(n))$ is even is*

$$(2.25) \quad \frac{1}{2} \left(1 + \prod_p \left(1 - \frac{2}{p^2} \right) \right).$$

Finally, we consider the case in which S is the set F of those n for which $\Omega(n)$ is even. One obtains in this case

COROLLARY 2.8. *If $x \geq 2$, then*

$$(2.26) \quad B_\alpha(x, F) = \left(\frac{3\zeta(2\alpha + 2)c_{\alpha+1}}{\pi^2(\alpha + 1)} \right) x^{\alpha+1} + \begin{cases} O(x^{\alpha+1/2}) & \text{if } \alpha > 0, \\ O(\sqrt{x \log x}) & \text{if } \alpha = 0, \end{cases}$$

where

$$(2.27) \quad c_t = 1 + \frac{\zeta(4t)}{\zeta^2(2t)}, \quad t \geq 1.$$

The proof is similar to that of Corollary 2.7.

COROLLARY 2.8.1 ($\alpha = 0$). *If $x \geq 2$, then*

$$(2.28) \quad B(x, F) = \frac{7x}{10} + O(\sqrt{x \log x}).$$

COROLLARY 2.8.2 (CESÀRO [2, §20]). *The asymptotic density of the set of n for which $\Omega(Q(n))$ is even is equal to $7/10$.*

3. Problem II. Estimates for $B'_\alpha(x, S)$. Corresponding to (2.1) we define

$$(3.1) \quad \Gamma'_\alpha(n, S) = \begin{cases} \left(\frac{n}{Q^2(n)}\right)^\alpha & \text{if } Q(n) \in S, Q(n) \parallel n, \\ 0 & \text{otherwise.} \end{cases}$$

In case $\alpha = 0$, $\Gamma'_\alpha(n, S)$ reduces to the characteristic function of the sequence of n for which $Q(n)$ is a unitary divisor of n and $Q(n) \in S$. From (1.1) one obtains

LEMMA 3.1.

$$(3.2) \quad \Gamma'_\alpha(n, S) = \sum_{d^2 \star e = n \cdot d \in S} e^\alpha \mu^2(e).$$

We shall also need the following generalization of Lemma 2.2.

LEMMA 3.2 [4, LEMMA 4.2]. *If $s \geq 0$ and r is a fixed positive integer, then*

$$(3.3) \quad \mathfrak{V}_{r,s}(x) \equiv \sum_{n \leq x; (n,r)=1} \mu^2(n) n^s = \frac{6r}{\pi^2 \psi(r)} \left(\frac{x^{s+1}}{s+1}\right) + O(\theta(r)x^{s+1/2}),$$

where $\theta(r)$ denotes the number of square-free divisors of r and $\psi(r)$ is Dedekind's ψ -function,

$$(3.4) \quad \psi(r) = r \prod_{p|r} \left(1 + \frac{1}{p}\right) = \sum_{d\delta=r} \mu^2(d)\delta.$$

REMARK 3.1. Note that $r/\psi(r) = O(1)$.

THEOREM 3.1. *If $\alpha \geq 0$, then for $x \geq 2$,*

$$(3.5) \quad B'_\alpha(x, S) = \frac{6}{\pi^2} \left(\frac{x^{\alpha+1}}{\alpha+1}\right) \zeta'_S(2\alpha+1) + O(x^{\alpha+1/2} R'_\alpha(x, S)),$$

where

$$(3.6) \quad \zeta'_S(s) = \sum_{n=1; n \in S}^{\infty} \frac{1}{n^s \psi(n)}, \quad s > 0,$$

and where

$$(3.7) \quad R'_\alpha(x, S) = \sum_{\sqrt{x} \cdot n \in S} \frac{\theta(n)}{n^{2\alpha+1}},$$

unless the sum is vacuous, in which case $R'_\alpha(x, S) = 1$. The O -constant is independent of the set S .

REMARK 3.2. Since $\sum_{n \leq x} \theta(n)/n = O(\log^2 x)$ (cf. [5, (3.4)]),

$$(3.8) \quad R'_\alpha(x, S) = \begin{cases} O(1) & \text{if } \alpha > 0, \\ O(\log^2 x) & \text{if } \alpha = 0. \end{cases}$$

In addition, if $\theta(n)$ is bounded on S , $R'_0(x, S) = O(\log x)$, while, if S is finite, $R'_0(x, S) = O(1)$.

Proof. By the definition of $B'_\alpha(x, S)$ and Lemma 2.1,

$$(3.9) \quad B'_\alpha(x, S) = \sum_{n \leq x} \Gamma'_\alpha(n, S) = \sum_{d^2 * e \leq x; d \in S} e^\alpha \mu^2(e) = \sum_{n \leq \sqrt{x}; n \in S} \mathbf{U}_{n, \alpha} \left(\frac{x}{n^2} \right),$$

so that by Lemma 3.2 and Remark 3.1,

$$\begin{aligned} B'_\alpha(x, S) &= \frac{6}{\pi^2} \left(\frac{x^{\alpha+1}}{\alpha+1} \right) \sum_{n \leq \sqrt{x}; n \in S} \frac{1}{n^{2\alpha+1} \psi(n)} + O(x^{\alpha+1/2} R'_\alpha(x, S)) \\ &= \frac{6}{\pi^2} \left(\frac{x^{\alpha+1}}{\alpha+1} \right) \zeta'_S(2\alpha+1) + O \left(x^{\alpha+1} \sum_{n > \sqrt{x}} \frac{1}{n^{2\alpha+2}} \right) + O(x^{\alpha+1/2} R'_\alpha(x, S)), \end{aligned}$$

from which the theorem results.

It is easily verified that for $s > 0, t > 0$,

$$(3.10) \quad \zeta'_J(s) \equiv \sum_{n=1}^{\infty} \frac{1}{n^s \psi(n)} = \zeta(s+1) \beta_{s+1}, \quad \beta_t \equiv \prod_p \left(1 - \frac{p^{-t}}{p+1} \right).$$

COROLLARY 3.1 ($S = J$). If $x \geq 2$, then

$$(3.11) \quad B'_\alpha(x, J) = \frac{6}{\pi^2} \left(\frac{x^{\alpha+1}}{\alpha+1} \right) \zeta(2\alpha+2) \beta_{2\alpha+2} + \begin{cases} O(x^{\alpha+1/2}) & \text{if } \alpha > 0, \\ O(\sqrt{x} \log^2 x) & \text{if } \alpha = 0. \end{cases}$$

Place $B'(x, S) = B'_0(x, S)$. Then

COROLLARY 3.1.1 ($\alpha = 0$).

$$(3.12) \quad B'(x, J) = \beta' x + O(\sqrt{x} \log^2 x) \quad (\beta' = \beta_2).$$

Letting J' denote the set of integers n whose greatest square divisor is unitary, one obtains

COROLLARY 3.1.2. The sequence J' has asymptotic density β_2 .

Next we consider the set S consisting of the single integer r .

COROLLARY 3.2. For all $\alpha \geq 0$,

$$(3.13) \quad B'_\alpha(x, r) = \frac{6}{\pi^2} \left(\frac{x^{\alpha+1}}{\alpha + 1} \right) \frac{r^{-2\alpha-1}}{\psi(r)} + O\left(\frac{x^{1/2+\alpha}\theta(r)}{r^{2\alpha+1}}\right),$$

uniformly in $r \leq \sqrt{x}$.

COROLLARY 3.2.1 ($\alpha=0$). The asymptotic density of the subset of J' consisting of those n for which $Q(n) = r$ is $6/r\psi(r)\pi^2$.

Let S_k denote the set of k th powers of the primes, $k \geq 1$, k fixed. Then

COROLLARY 3.2.2. For $x \geq 3$,

$$(3.14) \quad B'(x, S_k) = \frac{6x}{\pi^2} \left(\sum_p \frac{1}{p^{2k-1}(p+1)} \right) + \begin{cases} O(\sqrt{x}) & \text{if } k > 1, \\ O(\sqrt{x} \log \log x) & \text{if } k = 1. \end{cases}$$

Proof. This is a consequence of the theorem, in connection with (2.17).

COROLLARY 3.2.3. The asymptotic density of the set of integers all of whose prime divisors are simple, with the exception of a single divisor of multiplicity $2k$, is

$$\frac{6}{\pi^2} \sum_p \frac{1}{p^{2k-1}(p+1)}.$$

The case $k = 1$ yields

COROLLARY 3.2.4 (RÉNYI [13, (6)]; ALSO CF. [7, (1.1)]). The set of integers whose prime divisors are all simple, with the exception of a single double divisor, has asymptotic density,

$$\frac{6}{\pi^2} \sum_p \frac{1}{p(p+1)}.$$

Let $N_s(a)$ be defined as in §2.

COROLLARY 3.3. The asymptotic density of the integers n of J' for which $Q(n) \in N_s(a)$, is determined by

$$(3.15) \quad \delta'_s(a) = \frac{6}{\pi^2} \sum_{p_1, \dots, p_s} \frac{1}{p_1^{2a_1-1}(p_1+1) \cdots p_s^{2a_s-1}(p_s+1)},$$

where the summation satisfies the conditions described in Corollary 2.6.1.

Note that the case $s = 1$ of this result yields a second proof of Corollary 3.2.3.

Let now d'_m have the same significance as in the introduction. By Corollary 3.3, one finds that

COROLLARY 3.3.1. *If $|z| < 2$,*

$$(3.16) \quad \sum_{m=0}^{\infty} d'_m z^m = \frac{6}{\pi^2} \prod_p \left(1 + \frac{pz}{(p+1)(p^2-z^2)} \right).$$

The proof is analogous to that of Corollary 2.6.2 and is therefore omitted.

Let E denote the set of n with $\omega(n)$ even.

COROLLARY 3.4. *The asymptotic density of the set of n contained in J' , such that $\omega(Q(n))$ is even, is given by*

$$\frac{1}{2}(\beta' + \gamma'), \quad \beta' = \beta_2, \quad \gamma' = \prod_p \left(1 - \frac{2p+1}{p^2(p+1)} \right),$$

where β_i is defined by (3.10).

The proof is similar to that of Corollary 2.7.2.

As in §2, let F denote the sequence of n such that $\Omega(n)$ is even. Analogous to Corollary 3.4, one may prove

COROLLARY 3.5. *The asymptotic density of the sequence of n for which $n \in J'$ and $\Omega(Q(n))$ is even is $(5\beta' + 2\delta')/10$, where*

$$\beta' = \prod_p \left(1 - \frac{1}{p^2(p+1)} \right), \quad \delta' = \prod_p \left(1 + \frac{1}{p^2(p+1)} \right).$$

REMARK 3.3. Corollary 3.1.2 also results from the case $z = 1$ in (1.9), or equivalently (3.16), by virtue of (1.10) in connection with the following principle: Let G_1, G_2, \dots be mutually disjoint sets of integers with densities $D(G_i), i \geq 1$, such that $J = \bigcup_{i=1}^{\infty} G_i$ and $\sum_1^{\infty} D(G_i) = 1$; if $J^* \subseteq J$ and $\Gamma = \bigcup_{i \in J^*} G_i$, then $D(\Gamma)$ exists and

$$D(\Gamma) = \sum_{i \in J^*} D(G_i).$$

4. **Problem III. Estimates for $B_{\alpha}^*(x, T)$.** Let L have the meaning of §1 and suppose that T is a nonvacuous subset of L . We define

$$(4.1) \quad \Gamma_{\alpha}^*(n, T) = \begin{cases} \left(\frac{n}{Q^*(n)} \right)^{\alpha} & \text{if } Q^*(n) \in T, \\ 0 & \text{otherwise.} \end{cases}$$

Evidently, $\Gamma_{\alpha}^*(n, T)$ is the characteristic function of the sequence of n for which $Q^*(n) \in T$. The uniqueness of the factorization (1.4) leads to

LEMMA 4.1.

$$(4.2) \quad \Gamma_{\alpha}^*(n, T) = \sum_{d \bullet e = n; d \in T} e^{\alpha} \mu^2(e).$$

Let $L(x)$ denote the enumerative function of L , representing the number of $n \leq x$ contained in L . It is an elementary fact (cf. [8, §1]) that

$$(4.3) \quad R(x) = O(\sqrt{x}).$$

We now prove

THEOREM 4.1. *If $\alpha \geq 0$, then for $x \geq 2$,*

$$(4.4) \quad B_\alpha^*(x, T) = \frac{6}{\pi^2} \left(\frac{x^{\alpha+1}}{\alpha+1} \right) \zeta_T^*(\alpha) + O(x^{\alpha+1/2} R_\alpha^*(x, T)),$$

where $\zeta_T^*(\alpha) = \zeta_T'(\alpha)$, as defined by (3.6), and

$$(4.5) \quad R_\alpha^*(x, T) = \sum_{n \leq x; n \in T} \frac{\theta(n)}{n^{\alpha+1/2}},$$

if this sum is nonvacuous; otherwise, $R_\alpha^*(x, T) = 1$. The O -constant in (4.4) is not dependent upon the set T ; moreover, Remark 3.2 is valid here if S is replaced by T and $R'_\alpha(x, S)$ by $R_\alpha^*(x, T)$.

Proof. By Lemma 3.1

$$(4.6) \quad B_\alpha^*(x, T) = \sum_{n \leq x} \Gamma_\alpha^*(n, T) = \sum_{de \leq x; d \in T; (d, e) = 1} e^\alpha \mu^2(e) = \sum_{n \leq x; n \in T} \mathfrak{U}_{n, \alpha} \left(\frac{x}{n} \right),$$

where $\mathfrak{U}_{r, s}$ is defined as in Lemma 3.2. Application of (3.3) gives

$$(4.7) \quad \begin{aligned} B_\alpha^*(x, T) &= \frac{6}{\pi^2} \left(\frac{x^{\alpha+1}}{\alpha+1} \right) \sum_{n \leq \sqrt{x}; n \in T} \frac{1}{n^\alpha \psi(n)} + O(x^{\alpha+1/2} R_\alpha^*(x, T)) \\ &= \frac{6}{\pi^2} \left(\frac{x^{\alpha+1}}{\alpha+1} \right) \zeta_T^*(\alpha) + O \left(x^{\alpha+1} \sum_{n > x; n \in T} \frac{1}{n^{\alpha+1}} \right) + O(x^{\alpha+1/2} R_\alpha^*(x, T)). \end{aligned}$$

That $\zeta_T^*(\alpha)$ is finite for $\alpha \geq 0$ is evident from the identity [9]

$$(4.8) \quad \sum_{n \in L} \frac{1}{\psi(n)} = \frac{\pi^2}{6}.$$

By partial summation and (4.3),

$$\begin{aligned} \sum_{n > x; n \in T} \frac{1}{n^{\alpha+1}} &= o \left(\sum_{n > x; n \in L} \frac{1}{n^{\alpha+1}} \right) = o \left(\sum_{n > x} \frac{L(n)}{n^{\alpha+2}} \right) + o \left(\frac{L(x)}{x^{\alpha+1}} \right) \\ &= \sum_{n > x} \frac{1}{n^{\alpha+3/2}} + o \left(\frac{1}{x^{\alpha+1/2}} \right) = o \left(\frac{1}{x^{\alpha+1/2}} \right), \end{aligned}$$

and (4.4) results from (4.7).

The final statement of the theorem concerning $R_\alpha^*(x, T)$ is a consequence of the same type of argument carried out in detail in [5, §3] and will not be reproduced here.

Observe that $\Gamma_\alpha^*(n, L)$ is $\gamma_1^\alpha(n)$ where $\gamma_1(n)$ is the greatest unitary square-free divisor of n .

COROLLARY 4.1 (IN CASE $\alpha \geq 1$, CF. [4, COROLLARY 4.1.3]). *If $\alpha > 0$, then*

$$(4.9) \quad B_\alpha^*(x, L) \equiv \sum_{n \leq x} \gamma_1^\alpha(n) = \frac{6}{\pi^2} \left(\frac{x^{\alpha+1}}{\alpha+1} \right) \zeta(\alpha+1) k_\alpha + O(x^{\alpha+1/2}),$$

where k_α is defined by

$$(4.10) \quad k_\alpha = \prod_p \left(1 - \frac{p^{\alpha+1} + p^\alpha - 1}{(p+1)p^{2\alpha+1}} \right).$$

The case $\alpha = 1$ gives

COROLLARY 4.1.1 (COHEN [4, COROLLARY 4.1.4]).

$$(4.11) \quad \sum_{n \leq x} \gamma_1(n) = \frac{k_1 x^2}{2} + O(x^{3/2}).$$

Let $N = N_s(a)$ be defined as in §2 and place $B_0^*(x, T) = B^*(x, T)$.

COROLLARY 4.2. *Let a_1, \dots, a_s be an s -tuple of integers, each > 1 . Then for $x \geq 3$,*

$$(4.12) \quad B^*(x, N) = \left(\frac{6x}{\pi^2} \right) \zeta_N^*(1) + O(\sqrt{x} (\log \log x)^t),$$

where t is the number of 2's in the set a_1, \dots, a_s .

Proof. By (2.17) and the fact that $\theta(n)$ is bounded on $N_s(a)$.

COROLLARY 4.2.1 (RÉNYI [13, (17)]). *The asymptotic density of the integers n for which $Q^*(n) \in N_s(a)$, where a_1, \dots, a_s are all > 1 , is given by*

$$(4.13) \quad \frac{6}{\pi^2} \sum_{p_1, \dots, p_s} \frac{1}{p_1^{\alpha_1-1}(p_1+1) \cdots p_s^{\alpha_s-1}(p_s+1)},$$

where the summation satisfies the conditions of Corollary 2.6.1.

COROLLARY 4.2.2 (RÉNYI [13, (3')]).

$$(4.14) \quad \sum_{m=0}^{\infty} d_m^* z^m = \frac{6}{\pi^2} \prod_p \left(1 + \frac{z}{(p+1)(p-z)} \right).$$

The proof is similar to that of Corollary 2.6.2.

One may observe that $\omega(Q(n)) = \omega(Q^*(n))$; hence Theorem 3 yields no new result analogous to Corollary 2.7.2. The corresponding problem for the Ω -function, however, has the following solution,

COROLLARY 4.3. *The set of integers n for which $\Omega(Q^*(n))$ is even has asymptotic density,*

$$(4.15) \quad \frac{1}{2} + \frac{1}{5} \prod_p \left(1 + \frac{p^2 + p + 1}{p^3(p + 1)} \right).$$

BIBLIOGRAPHY

1. W. F. Bodmer, *The limiting frequencies of integers with a given partitional characteristic*, J. Roy. Statist. Soc. Ser. A **21** (1959), 134–143.
2. Ernesto Cesàro, *Le plus grand diviseur carré*, Ann. Mat. Pura Appl. **13** (1885), 251–268.
3. Eckford Cohen, *Arithmetical functions associated with arbitrary sets of integers*, Acta Arith. **5** (1959), 407–415.
4. ———, *On an elementary method in the asymptotic theory of numbers*, Duke Math. J. **28** (1961), 183–192.
5. ———, *Arithmetical notes. V. A divisibility property of the divisor function*, Amer. J. Math. **83** (1961), 693–697.
6. ———, *Arithmetical notes. VII. Some classes of even functions (mod r)*, Collect. Math. (to appear).
7. ———, *Arithmetical notes. VIII. An asymptotic formula of Rényi*, Proc. Amer. Math. Soc. **13** (1962), 536–539.
8. ———, *On the distribution of certain sequences of integers*, Amer. Math. Monthly **70** (1963), 516–521.
9. ———, *A property of Dedekind's ψ -function*, Proc. Amer. Math. Soc. **12** (1961), 996.
10. W. Feller and E. Tornier, *Mengentheoretische Untersuchungen von Eigenschaften der Zahlenreihe*, Math. Ann. **107** (1933), 188–232.
11. M. Kac, *A remark on the preceding paper by A. Rényi*, Acad. Serbe Sci. Publ. Inst. Math. **8** (1955), 163–165.
12. Hans-Joachim Kanold, *Über die Dichte von gewissen Zahlenmengen*, J. Reine Angew. Math. **193** (1954), 250–252.
13. Alfred Rényi, *On the density of certain sequences of integers*, Acad. Serbe Sci. Publ. Inst. Math. **8** (1955), 157–162.
14. Peter Scherk, *Bemerkungen zu einer Arbeit von Herrn Kanold*, J. Reine Angew. Math. **196** (1956), 133–136.
15. I. J. Schoenberg, *On asymptotic distributions of arithmetical functions*, Trans. Amer. Math. Soc. **39** (1936), 315–330.
16. ———, *On two theorems of P. Erdős and A. Rényi*, Illinois J. Math. **6** (1962), 53–57.

UNIVERSITY OF TENNESSEE,
KNOXVILLE, TENNESSEE