

THE TOMITA DECOMPOSITION OF RINGS OF OPERATORS

BY

JOSEPH L. TAYLOR⁽¹⁾

I. Introduction. It is known that if R is a symmetric ring of bounded operators on a separable Hilbert space H and ξ_0 is a vector in H which is cyclic with respect to R , then the positive functional $F(A) = (A\xi_0, \xi_0)$, for $A \in R$, may be written as a direct integral over a compact Hausdorff space M , i.e., $F(A) = \int_M f_m(A) d\mu(m)$ where μ is a positive regular Borel measure and the functionals f_m are indecomposable except, at most, for $m \in M_0 \subset M$ and $\mu(M_0) = 0$. This decomposition of F induces a representation of R as a direct integral of rings R_m of operators on a Hilbert space H_m and for almost all $m \pmod{\mu}$, R_m is an irreducible ring on H_m .

The problem of extending this type of decomposition to rings of operators on an arbitrary Hilbert space was attacked in 1954 by Tomita (cf. [6]) using extremely penetrating techniques. However, certain parts of Tomita's development of his decomposition theory require a special measure theoretic result which is not valid in general. Consequently, the question of whether or not this measure theoretic difficulty could be circumvented arose; i.e., did the Tomita decomposition hold for arbitrary rings and, if not, for what type of rings did it hold?

In this paper we shall first show (Theorem 2.3) that in case R is a weakly closed symmetric ring which contains its commutant, then the Tomita decomposition holds and, in fact, all of the rings R_m are irreducible; furthermore R is completely determined by the representation of H as a direct integral. In §3 of this paper we shall construct a symmetric ring R where the Tomita decomposition fails to hold (Theorem 3.3). In our example no one of the rings R_m is irreducible. We conclude the paper in §4 with some remarks indicating how recent results of Loomis [4] can be applied to obtain irreducible decomposition of rings of operators.

II. Notation and terminology. Throughout this paper we shall use the notation and terminology employed in Naimark's treatise, [5]. Also, we shall use the ring theory developed in [5] together with those parts of Tomita's formulation of the decomposition problem which are valid.

Let H denote an arbitrary Hilbert space, R_1 a symmetric Banach ring of operators on H , and E a maximal commutative subring contained in the commutant of R_1 . Let $R = R_1 \cup E$; hence, E is both the center and the commutant of R . We

Presented to the Society, August 30, 1963; received by the editors May 20, 1963.

⁽¹⁾ This research was supported in part by a National Science Foundation grant.

shall denote the maximal ideal space of E by M and the continuous functions on M by $C(M)$. N will denote the set of normalized indecomposable positive functionals on R and \bar{N} the closure of N in the weak- $*$ topology of R^* , where R^* is the dual of R . Finally, we shall assume that H contains a vector h_0 of norm one such that h_0 is cyclic with respect to R . We set $F(A) = (Ah_0, h_0)$, $A \in R$, so that $F \in R^*$. Theorem 2.1 and 2.2 below summarize certain parts of Tomita's theory that we shall need; for the details of these theorems, the reader is referred to [5].

THEOREM 2.1. *There exists a homeomorphism, denoted by $m \rightarrow f_m$, of M into \bar{N} and a positive, normalized regular Borel measure μ on M such that*

- (1) $F(A) = \int_M f_m(A) d\mu(m)$, for each $A \in R$,
- (2) $f_m(AB) = f_m(A)x_B(m)$, for each $A \in R$ and each $B \in E$, where $B \rightarrow x_B(m)$ is the natural isomorphism of E onto $C(M)$,
- (3) the mapping $m \rightarrow f_m$ and the measure μ are uniquely determined by (1) and (2), and
- (4) the carrier of μ is all of M .

For the verification of these statements see [5], in particular, II, p. 507; Theorem 1, p. 493, and III, p. 507.

Before stating Theorem 2.2, we shall use the above theorem to formulate the notion of a direct integral Hilbert space. This is done in the following manner. For each $m \in M$, the functional f_m determines a Hilbert space H_m , a ring of operators R_m on H_m , and a representation $A \rightarrow A_m$ of the ring R onto the ring R_m . There exists a vector $\xi_0(m) \in H_m$ of norm one such that $\xi_0(m)$ is cyclic with respect to R_m and $f_m(A) = (A_m \xi_0(m), \xi_0(m))$ for all $A \in R$. The set S of all vector valued functions of the form $\eta(m) = A_m \xi_0(m)$, $A \in R$, is a Euclidean space under the inner product

$$(\xi, \eta) = \int_M (\xi(m), \eta(m)) d\mu(m) = \int_M f_m(A^*C) d\mu(m),$$

where $A, C \in R$, $\xi(m) = C_m \xi_0(m)$, A^* is the adjoint of A . The completion of S , denoted by \hat{H} , is a Hilbert space and is called the direct integral of the spaces H_m with respect to μ . S is called the basis of the direct integral space \hat{H} .

THEOREM 2.2. *There exists a unitary operator U mapping H onto \hat{H} and an isometric mapping $A \rightarrow \hat{A}$ of R into $B(\hat{H})$ such that*

- (1) $A = U^{-1} \hat{A} U$, for $A \in R$,
- (2) $\hat{A} S \subset S$, for $A \in R$,
- (3) $(\hat{A} \xi)(m) = A_m \xi(m)$, for $A \in R$, $\xi \in S$, $m \in M$, and
- (4) if $B \in E$, then $(\hat{B} \xi)(m) = x_B(m) \xi(m)$ for $\xi \in S$ and $m \in M$; moreover, every continuous function on M defines an operator B in this fashion.

We refer the reader to pp. 512–515 in [5].

The image \hat{R} under the mapping $A \rightarrow \hat{A}$ is called a direct integral of the rings R_m . (3) shows that each operator \hat{A} may be considered as an operator valued function on M with values in the rings R_m . The essential problem in this type of decomposition is to determine if almost all of the rings R_m are irreducible; i.e., R_m is irreducible except for $m \in M_0$ where $\mu(M_0) = 0$. We shall now show that this is the case when R is weakly closed. We note that if R is weakly closed then R is the commutant of E .

We shall need two lemmas. In these lemmas we continue to use the notations and facts cited above.

LEMMA 2.1. *If y is an essentially bounded μ measurable function on M , then there exists $B \in E$ such that $y(m) = x_B(m)$ almost everywhere, where $B \rightarrow x_B(m)$ is in accordance with (2) of Theorem 2.1.*

Proof. Suppose $y \in L_\infty(\mu)$ and $0 \leq y(m) \leq 1$ for almost all $m \in M$. We define $G(A) = \int_M f_m(A)y(m)d\mu(m)$ for $A \in R$; hence, G is a positive functional on R and $G(A) \leq F(A) = (Ah_0, h_0)$, whenever A is a positive definite operator. Under these circumstances, there exists $B \in E$ such that $G(A) = F(AB)$ for $A \in R$ (cf. Theorem 1, p. 262 of [5]). Hence $\int_M f_m(A)y(m)d\mu(m) = \int_M f_m(A)x_B(m)d\mu(m)$ for $A \in R$.

In particular, the equality holds if we replace A by C , $C \in E$, and, using $f_m(C) = f_m(IC) = f_m(I)x_C(m) = x_C(m)$ ((2) of Theorem 2.1), we have that $\int_M k(m)y(m)d\mu(m) = \int_M k(m)x_B(m)d\mu(m)$ for every continuous function k on M . Hence, $x_B(m) = y(m)$ almost everywhere. The conclusion of Lemma 2.1 now follows immediately inasmuch as any function in $L_\infty(\mu)$ is a linear combination of functions of the above type.

LEMMA 2.2. *M is extremely disconnected, i.e., the closure of any open set is open.*

Proof. Suppose V is an open subset of M and χ_V the characteristic function of V on M . In accordance with Lemma 2.1, there exists a continuous function x_B , such that $0 \leq x_B \leq 1$, and $x_B = \chi_V$ almost everywhere. Hence, if $V_1 = \{m \in M; x_B(m) = 1\}$ and $V_2 = \{m \in M; x_B(m) > 0\}$, then V_1 is a closed set and V_2 is an open set. It follows that $\mu(V \cap (M - V_1)) = 0$ inasmuch as $x_B < \chi_V$ on $V \cap (M - V_1)$. However, the set $V \cap (M - V_1)$ is an open set and, in view of (4) of Theorem 2.1, we must have $V \cap (M - V_1) = \emptyset$ or, what is the same, $V \subset V_1$. Also, if \bar{V} denotes the closure of V , a similar argument shows that $V_2 \cap (M - \bar{V}) = \emptyset$, or $V_2 \subset \bar{V}$. Hence, $V \subset V_1 \subset V_2 \subset \bar{V}$ and, since V_1 is a closed set, $V_1 = \bar{V}$, so that $V_1 = V_2 = \bar{V}$. Since V_2 is an open set, \bar{V} is an open set.

THEOREM 2.3. *Suppose R is a symmetric ring of operator over a Hilbert space H such that*

- (1) *the center E of R is also the commutant of R ,*
- (2) *R is weakly closed, and*

(3) there exists a vector $h_0 \in H$ with norm one which is cyclic with respect to R and $F(A) = (Ah_0, h_0)$ for $A \in R$.

Then (a) the direct integral decomposition $F(A) = \int_M f_m(A) d\mu(m)$ has the property that each of the functionals f_m is indecomposable, and

(b) the direct integral decomposition of R given by Theorem 2.2 has the property that each of the rings R_m is irreducible over the Hilbert space H_m .

Proof. We will show that each of the rings R_m is irreducible and it will follow from this that each of the functionals f_m is indecomposable, inasmuch as the ring R_m is the image of R under the representation generated by f_m . In order to show that the ring R_m is irreducible, it is sufficient to prove that if $n \in M$, $\eta_n \in H_n$, $\gamma_n \in H_n$, $\gamma_n \neq 0$, and $\varepsilon > 0$, then there exists $A \in R$ such that $|A_n \gamma_n - \eta_n| < \varepsilon$. We shall prove this first for the case where η_n and γ_n are of the form $\eta_n = C_n \xi_0(n)$ and $\gamma_n = D_n \xi_0(n)$ for $C \in R$ and $D \in R$.

If $\gamma_n \neq 0$, then, since the function $|D_m \xi_0(m)|^2 = f(D_m^* D_m)$ is a continuous function on M , there exists an open set $V \subset M$ such that $n \in V$, $|D_m \xi_0(m)| \geq 2^{-1} |D_n \xi_0(n)| = \delta_1$, and $|C_m \xi_0(m)| \leq |C_n \xi_0(n)| + 1 = \delta_2$, for $m \in V$. In view of Lemma 2.2 we can select V so that it is compact and, hence, the characteristic function of V on M , say χ_V , is a continuous function and there exists a $P \in E$ such that $x_P = \chi_V$ everywhere. We note that $P^2 = P$.

We now define the subspace K of H to be the closed subspace generated by vectors of the form $h_1 = BPh_0$ for $B \in E$. For each such vector h_1 we have $|Dh_1|^2 = \int_M |D_m P_m B_m \xi_0(m)|^2 d\mu(m) = \int_M |x_P(m) x_B(m)|^2 |D_m \xi_0(m)|^2 d\mu(m) = \int_V |x_B(m)|^2 |D_m \xi_0(m)|^2 d\mu(m) \geq \delta_1^2 \int_V |x_B(m)|^2 d\mu(m) = \delta_1^2 \int_M |P_m B_m \xi_0(m)|^2 d\mu(m) = \delta_1^2 |h_1|^2$. Similarly, $|Ch_1| \leq \delta_2 |h_1|$.

Now each vector $h \in H$ can be uniquely expressed in the form $h = Dh_1 + h_2$ where $h_1 \in K$ and h_2 is in the orthogonal complement of the space DK . Thence, if we define a linear operator A on H by $Ah = Ch_1$, then

$$|Ah| = |Ch_1| = |Ch_1| (|Dh_1|)^{-1} |Dh_1| \leq \delta_2 \delta_1^{-1} |Dh_1 + h_2|,$$

so that A is a bounded operator and $|A| \leq \delta_2 \delta_1^{-1}$. Now, if $B \in E$, then $BAh = BCh_1 = CBh_1$, but $Bh_1 \in K$ and $Bh_2 \perp DK$, so that $CBh_1 = A(DBh_1 + Bh_2) = A(BDh_1 + Bh_2) = ABh$. Hence, A is in the commutant of E and, since R is weakly closed, the commutant of E is R ; i.e., $A \in R$. We also have

$$A_n \gamma_n = A_n D_n \xi_0(n) = (\hat{A} \hat{D} \xi_0)(n) = (\hat{C} \xi_0)(n) = C_n \xi_0(n) = \eta_n,$$

which establishes the assertion we made above for vectors of the form $\eta_n = C_n \xi_0(n)$ and $\gamma_n = D_n \xi_0(n)$. Finally, we also note that

$$|A| \leq \delta_2 \delta_1^{-1} \leq 2(|C_n \xi_0(n)| + 1) (|D_n \xi_0(n)|)^{-1}.$$

Suppose α_n and β_n are arbitrary vectors in H_n , $\alpha_n \neq 0$, and $\varepsilon > 0$. Since the vectors $C_n \xi_0(n)$, $C \in R$, are dense in H_n , there exist $C \in R$ and $D \in R$ such that

$|C_n \xi_0(n) - \beta_n| < 2^{-1} \varepsilon$, $|D_n \xi_0(n) - \alpha_n| < \varepsilon(4|\alpha_n|)^{-1}(|\beta_n| + 1)$, $|C_n \xi_0(n)| = |\beta_n|$, and $|D_n \xi_0(n)| = |\alpha_n|$. On the basis of the construction in the preceding paragraph there exists an operator $A \in R$ such that $A_n D_n \xi_0(n) = C_n \xi_0(n)$ and $|A| \leq 2(|C_n \xi_0(n)| + 1)(|D_n \xi_0(n)|^{-1}) = 2(|\beta_n| + 1)(|\alpha_n|^{-1})$. Hence,

$$\begin{aligned} |A_n \alpha_n - \beta_n| &\leq |A_n D_n \xi_0(n) - A_n \alpha_n| + |A_n D_n \xi_0(n) - C_n \xi_0(n)| + |C_n \xi_0(n) - \beta_n| \\ &= |A_n D_n \xi_0(n) - A_n \alpha_n| + |C_n \xi_0(n) - \beta_n| \\ &\leq |A| \cdot |D_n \xi_0(n) - \alpha_n| + |C_n \xi_0(n) - \beta_n| < \varepsilon. \end{aligned}$$

This completes the proof of Theorem 2.3.

III. An example. Suppose R_1 is a symmetric Banach ring of operators on the Hilbert space H , E a maximal commutative subring of the commutant of R_1 , $R_2 = R_1 \cup E$, and R the weak closure of R_2 (i.e., R is the commutant of E). Then, in accordance with Theorem 2.3, the Tomita decomposition of R yields a representation of R as a direct integral of irreducible rings over M ; the maximal ideal space of E ; moreover, R is completely determined by the representation of H as a direct integral over M , since R is the commutant of E and E is precisely (within unitary equivalence) the ring of operators (on this direct integral space) determined by the continuous functions on M . The ring R_2 also has a Tomita decomposition and it is related to the decomposition of the ring R in the following manner; in the representation of R as a direct integral of rings R_m , the image of R_2 is exactly the Tomita representation of R_2 as a direct integral of the rings $R_{2,m} \subset R_m$. However, the rings $R_{2,m}$ may no longer be irreducible; furthermore, R_2 is not characterized by the representation of H as a direct integral, and certainly R_1 is not characterized in this manner. We shall verify all of these statements by an example. Several lemmas and definitions will be needed in order to obtain our example.

LEMMA 3.1. *There exists a compact Hausdorff space M and a regular Borel measure μ on M such that*

- (1) *the carrier of μ is M ,*
- (2) *M has no isolated points, and*
- (3) *if y is an essentially bounded μ -measurable function on M , then there exists a continuous function x on M such that $x(m) = y(m)$ almost everywhere.*

Proof. Let ν denote the Lebesgue measure on $[0, 1]$. The ring $L_\infty(\nu)$ determines a commutative ring of operators, say Z , on $L_2(\nu)$ in the following manner; for $x \in L_\infty(\nu)$ and $y \in L_2(\nu)$, let $(B_x y)(t) = x(t)y(t)$ and $Z = \{B_x : x \in L_\infty(\nu)\}$. Z is isomorphic and isometric to $L_\infty(\nu)$ and we shall show that Z is its own commutant. To this end suppose $A \in B(L_2(\nu))$ and $AB_x = B_x A$ for all $x \in L_\infty(\nu)$, and let $x_A(t) = (AI)(t)$, where I is the identically one function in $L_2(\nu)$. Hence, if $y \in L_\infty(\nu) \subset L_2(\nu)$, then $(Ay)(t) = (AB_y I)(t) = (B_y AI)(t) = y(t)(AI)(t) = y(t)x_A(t)$.

Since $L_\infty(\nu)$ is dense in $L_2(\nu)$ it follows that A is the operator that is determined by pointwise multiplication with the function x_A , $x_A \in L_\infty(\nu)$, and Z is its own commutant.

It follows from Theorem 2.1 and Lemma 2.1 that if M is the maximal ideal space of Z and μ is the measure on M determined by the functional $F(B_x) = (B_x I, I) = \int_0^1 x(t) d\nu(t)$, then M and μ satisfy (1) and (3) of this lemma. That M has no isolated points follows from the fact that ν is a continuous measure; i.e., ν has no atomic parts.

This completes the proof of Lemma 3.1. Throughout the remainder of this paper, M and μ will remain fixed in accordance with Lemma 3.1.

DEFINITION 3.1. Let G denote the collection of all functions g from M to the two point group $I_2 = \{0, 1\}$. We define addition in G to be the pointwise addition and the topology of G to be the cross product topology of I_2^M , so that G is a compact topological Abelian group. Let ρ denote the normalized Haar measure on G and $\rho \times \mu$ the cross product measure induced by ρ and μ on the space $G \times M$. \mathfrak{H} denotes $L_2(\rho \times \mu)$, H denotes $L_2(\rho)$, and $B(\mathfrak{H})$ (resp. $B(H)$) denotes the space of bounded linear operators on \mathfrak{H} (resp. H).

DEFINITION 3.2. (a) If α is a continuous function on $G \times M$, then the operator $A_\alpha \in B(\mathfrak{H})$ is defined by $(A_\alpha \beta)(g, m) = \alpha(g, m) \beta(g, m)$ for $\beta \in \mathfrak{H}$. \mathfrak{C} is the collection of all such operators.

(b) If U is a compact-open set in M and $g' \in G$ such that $g'(m) = 0$ for $m \in U$, then the operator $T_{U, g'} \in B(\mathfrak{H})$ is defined by $(T_{U, g'} \beta)(g, m) = \beta(g + g', m)$ for $m \in U$ and $(T_{U, g'} \beta)(g, m) = 0$ for $m \notin U$. T is the collection of all such operators.

(c) $K \subset C(M) \subset \mathfrak{H}$ is defined to be the collection of all characteristic functions of the form $\chi_{V \times U}$, where U is a compact-open subset of M and V is a compact-open subset of G defined by $V = \{g \in G: g(m_i) = g_0(m_i)\}$ for $g_0 \in G$ and $\{m_i\}_{i=1}^n$ a finite subset of M .

(d) For $m \in M$, $V_m^0 = \{g \in G: g(m) = 0\}$, $V_m^1 = \{g \in G: g(m) = 1\}$, $H_m^0 = \{a \in H: a(g) = 0 \text{ for } g \in V_m^1\}$, and $H_m^1 = \{a \in H: a(g) = 0 \text{ for } g \in V_m^0\}$.

(e) R is the smallest norm-closed subring of $B(\mathfrak{H})$ containing $\mathfrak{C} \cup T$. We note since $\mathfrak{C}^* = \mathfrak{C}$ and $T^* = T$ that R is a symmetric ring.

LEMMA 3.2. *The linear subspace generated by K is dense in $C(G \times M)$ under the sup norm topology and is dense in the space \mathfrak{H} under the L_2 topology.*

Proof. It follows from Lemma 2.2 that M has topological basis of open-compact sets. Also, the sets $V = \{g \in G: g(m_i) = g_0(m_i), i = 1, \dots, n\}$ form a basis for G . Hence, the sets $V \times U$, U open-compact in M and V of the above form, constitute a compact-open neighborhood basis for $G \times M$. Thus the linear subspace generated by K is dense in $C(G \times M)$ and, consequently, in \mathfrak{H} .

LEMMA 3.3. *If $B \in B(\mathfrak{H})$ and B commutes with all the operators in \mathfrak{C} , then*

there exists $\beta \in L_\infty(\rho \times \mu)$ such that, for $\alpha \in \mathfrak{H}$, $(B\alpha)(g, m) = \beta(g, m)\alpha(g, m)$ almost everywhere (mod $\rho \times \mu$).

Proof. Let $\beta(g, m) = (BI)(g, m)$, where $I(g, m) \equiv 1$. Hence, if $\alpha \in C(G \times M)$, then $A_\alpha \in \mathfrak{C}$ and $(B\alpha)(g, m) = (BA_\alpha I)(g, m) = (A_\alpha BI)(g, m) = \alpha(g, m)\beta(g, m)$ almost everywhere. Since the extreme ends of the last chain of equalities holds for $\alpha \in C(G \times M)$, it must hold for $\alpha \in \mathfrak{H}$. Also, if $\alpha = \chi_{V \times U} \in K$, then

$$\int_{V \times U} |\beta(g, m)|^2 d\rho \times \mu = |B\alpha|^2 \leq |B|^2 |\alpha|^2 = |B|^2 (\rho \times \mu(V \times U))^2$$

so that $\beta \in L_\infty(\rho \times \mu)$ and, moreover, its L_∞ norm is $|B|$.

THEOREM 3.1. *If $B \in B(\mathfrak{H})$ and B commutes with all the operators in R , then there exists a continuous function x_B on M such that, for $\alpha \in \mathfrak{H}$, $(B\alpha)(g, m) = x_B(m)\alpha(g, m)$ almost everywhere (mod $\rho \times \mu$).*

Proof. In view of Lemma 3.3, there exists $\beta \in L_\infty(\rho \times \mu)$ such that $(B\alpha)(g, m) = \beta(g, m)\alpha(g, m)$. If $y(m) = \int_G \beta(g, m) d\rho(g)$, then $y \in L_\infty(\mu)$ and, by Lemma 3.1, there exists $x_B \in C(M)$ such that $x_B(m) = y(m)$ almost everywhere (mod μ). Let B' be the operator in $B(\mathfrak{H})$ such that, for $\alpha \in \mathfrak{H}$, $(B'\alpha)(g, m) = x_B(m)\alpha(g, m)$. We shall show that if $\alpha \in K$, then $(B\alpha, I) = (B'\alpha, I)$ and, since K generates \mathfrak{H} , it will follow that $(B\alpha, I) = (B'\alpha, I)$ for $\alpha \in \mathfrak{H}$. This, in turn, will imply that $(B'A_\gamma^* \alpha, I) = (BA_\gamma^* \alpha, I)$ for $A_\gamma \in \mathfrak{C}$, and, since B' and B commute with all operators in \mathfrak{C} , $(B'\alpha, \gamma) = (B\alpha, \gamma)$ for $\gamma \in C(G \times M)$; consequently, $(B'\alpha, \gamma) = (B\alpha, \gamma)$ for arbitrary $\alpha, \gamma \in \mathfrak{H}$ and, hence, $B = B'$.

In order to prove that $(B'\alpha, I) = (B\alpha, I)$ for $\alpha \in K$, let $\alpha = \chi_{V \times U}$ where U is compact-open in M and $V = \{g \in G : g(m_i) = g_0(m_i), i = 1, \dots, n\}$. Let (a_{ij}, \dots, a_{nj}) , $j = 1, \dots, 2^n$, be all possible n -tuples of zeros and ones and set $g_j(m_i) = a_{ij}$ and $g_j(m) = 0$ for $m \neq m_i, i = 1, \dots, n$. The collection of sets $\{V + g_j\}_{j=1}^{2^n}$ form a pairwise disjoint compact-open cover of G .

Now parts (1) and (2) of Lemma 3.1 imply that no point of M has positive μ measure; hence, for $\delta > 0$, there exists a compact open set $U' \subset U$ such that $\mu(U - U') < \delta$ and $m_i \notin U'$ for $i = 1, \dots, n$.

It follows now (cf. (b) of Definition 3.2) that the operator $T_{U', g_j} \in T$ for $j = 1, \dots, 2^n$; moreover $(BT_{U', g_j} \alpha, I) = (T_{U', g_j} B\alpha, I)$, i.e.,

$$\begin{aligned} & \int_M \int_G \beta(g, m) \chi_{V \times U'}(g + g_j, m) d\rho(g) d\mu(m) \\ &= \int_M \int_G \beta(g + g_j, m) \chi_{V \times U'}(g + g_j, m) d\rho(g) d\mu(m). \end{aligned}$$

Now using the invariance of the Haar measure and the facts $\rho(V) = 2^{-n}$ and $U - g = g$, we have:

$$\begin{aligned}
& \int_M \int_G \beta(g, m) \chi_{V \times U'}(g, m) d\rho(g) d\mu(m) \\
&= 2^{-n} \sum_{j=1}^{2^n} \int_M \int_G \beta(g + g_j, m) \chi_{V \times U'}(g + g_j, m) d\rho(g) d\mu(m) \\
&= 2^{-n} \sum_{j=1}^{2^n} \int_M \int_G \beta(g, m) \chi_{V \times U'}(g + g_j, m) d\rho(g) d\mu(m) \\
&= 2^{-n} \int_{U'} \sum_{j=1}^{2^n} \int_{V+g_j} \beta(g, m) d\rho(g) d\mu(m) = 2^{-1} \int_{U'} \int_G \beta(g, m) d\rho(g) d\mu(m) \\
&= 2^{-n} \int_{U'} x_B(m) d\mu(m) = \rho(V) \int_{U'} x_B(m) d\mu(m) \\
&= \int_M \int_G x_B(m) \chi_{V \times U'}(m, g) d\rho(g) d\mu(m).
\end{aligned}$$

Letting δ approach zero, this yields

$$\begin{aligned}
(B\alpha, I) &= \int_M \int_G \beta(g, m) \chi_{V \times U'}(g, m) d\rho(g) d\mu(m) \\
&= \int_M \int_G x_B(m) \chi_{V \times U'}(g, m) d\rho(g) d\mu(m) = (B'\alpha, I).
\end{aligned}$$

This completes the proof of Theorem 3.1.

COROLLARY 3.1. *If E is the ring of operators B on \mathfrak{H} of the form $(B\alpha)(g, m) = x_B(m)\alpha(g, m)$ for $x_B \in C(M)$, then E is both the commutant and center of R ; moreover, M is the maximal ideal space of E .*

Proof. Clearly $E \subset \mathfrak{C} \subset R$ and, on the basis of Theorem 3.1, the commutant of R is E . It follows that E is the center of R . The mapping $B \leftrightarrow x_B$ is an isometry of E onto $C(M)$, so that M is the maximal ideal space of E .

LEMMA 3.4. *Let S be the subset of \mathfrak{H} defined by $S = \{A_\alpha I : A_\alpha \in \mathfrak{C}\}$. Then*

- (a) S is dense in \mathfrak{H} ,
- (b) $AS \subset S$ for each $A \in R$, and
- (c) each $L_2(\rho \times \mu)$ equivalence class of S contains exactly one continuous function.

Proof. (a) follows directly from the definition of \mathfrak{C} . The set R_1 of all A in R such that $AS \subset S$ is a norm-closed subring of R ; moreover, $A_\alpha S \subset S$ for $A_\alpha \in \mathfrak{C}$ and $T_{U, g'} S \subset S$ for $T_{U, g'} \in T$, hence $\mathfrak{C}UT \subset R_1$ and, hence, $R_1 = R$. (c) follows from the fact that the carrier of $\rho \times \mu$ is $G \times M$.

LEMMA 3.5. For each $A \in R$ and $m \in M$, there is a unique operator $A_m \in B(H)$ such that $(A\alpha)(g, m) = (A_m\alpha_m)(g)$, where $\alpha_m \in H$ and is the function determined by $\alpha_m(g) = \alpha(m, g)$ for $\alpha \in S$; finally, the correspondence $A \rightarrow A_m$ is a symmetric and norm-decreasing representation of the ring R onto a ring $R_m \subset B(H)$.

Proof. For a fixed $n \in M$, let R_1 be the set of all operators A in R for which there exists an operator $A_n \in B(H)$ such that $(A\alpha)(n, g) = (A_n\alpha_n)(g)$ for all $\alpha \in S$. R_1 is a norm-closed subring of R which contains \mathbb{C} and T , consequently, $R_1 = R$. The set $\{\alpha_n: \alpha \in S\}$ is dense in H and for each $A \in R$ and $\alpha \in S$, $A\alpha$ defines a unique continuous function in S . Hence, A_n is uniquely defined by A . Finally, it is easily seen that the correspondence $A \rightarrow A_n$ is a symmetric representation of R and $|A| = \sup_{n \in M} |A_n|$.

THEOREM 3.2. If, for $A \in R$, $F(A) = (AI, I)$, I the identically one function on $G \times M$, and $f_m(A) = \int_G (AI)(g, m) d\rho(g)$, then (a) $F(A) = \int_M f_m(A) d\mu(m)$ is the Tomita decomposition of F described in Theorem 2.1, and (b) the correspondence $A \rightarrow A_m$ of Lemma 3.5, is the representation of A induced by f_m .

Proof. $F(A) = (AI, I) = \int_M \int_G (AI)(g, m) d\rho(g) d\mu(m) = \int_M f_m(A) d\mu(m)$; also, for $B \in E$, $f_m(BA) = \int_G x_B(m) (AI)(g, m) d\rho(g) = x_B(m) f_m(A)$, and $f_m(A)$ is a continuous function of m for fixed A . Hence, in accordance with (1), (2) and (3) of Theorem 2.1, $F(A) = \int_M f_m(A) d\mu(m)$ is the Tomita decomposition of F . Now, from Lemma 3.5, $f_m(A) = \int_G (AI)(g, m) d\rho(g) = \int_G (A_m I_m)(g) d\rho(g) = (A_m I_m, I_m)$ in H . This equality, along with the fact that $\{A_m I_m: A \in R\}$ is dense in H , uniquely defines the representation generated by f_m to within unitary equivalence. Hence, $A \rightarrow A_m$ is this representation.

THEOREM 3.3. No one of the functionals f_m , described in Theorem 3.2, is indecomposable and no one of the rings R_m is irreducible.

Proof. We shall prove the theorem by showing that the spaces H_m^0 and H_m^1 are invariant under the ring R_m . To this end, let R_m^0 denote the norm-closed ring consisting of all operators $A_m \in R_m$ such that $A_m H_m^0 \subset H_m^0$ and $A_m H_m^1 \subset H_m^1$ (cf. (d) of Definition 3.2). Clearly, $\mathfrak{C}_m = \{A_m: A \in \mathfrak{C}\} \subset R_m^0$. Let $T_m = \{A_m: A \in T\}$. If $T_{U, g'} \in T$, then $g'(m) = 0$ for $m \in U$, $(T_{U, g'}\alpha)(g, m) = \alpha(g + g', m)$ for $m \in U$, and $(T_{U, g'}\alpha)(g, m) = 0$ for $m \notin U$. Hence, if $\alpha \in S$ and $\alpha_m \in H_m^0$, then $(T_{U, g'}\alpha)(g, m) = 0$ for $m \notin U$ and $g \in V_m^1$. Hence, $A_m \alpha_m \in H_m^0$ for $\alpha_m \in H_m^0$ and $A \in T$. A similar argument shows that the same is true for the space H_m^1 . Since $\mathfrak{C}_m \cup T_m$ generates R_m and $\mathfrak{C}_m \cup T_m \subset R_m^0 \subset R_m$, we have $R_m = R_m^0$ and, consequently, H_m^0 and H_m^1 are invariant subspaces of R_m . Hence, no R_m is irreducible and no f_m is indecomposable.

This concludes the proof of Theorem 3.3; however, a few observations about the functional f_m may be worthwhile. For each $m \in M$, the functional f_m can be written as $f_m^0 + f_m^1$ where

$$f_m^0(A) = \int_{V_m^0} (A_m I_m)(g) d\rho \quad \text{and} \quad f_m^1(A) = \int_{V_m^1} (A_m I_m)(g) d\rho, \quad A \in R.$$

The functionals f_m^0 and f_m^1 are distinct positive functionals on R having the interesting property that $\int_M f_m^0(A) d\mu = \int_M f_m^1(A) d\mu = 2^{-1} \int_M f_m(A) d\mu$ for $A \in R$. This emphasizes one essential drawback of the Tomita decomposition, namely, unless R is weakly closed, the rings R_m may have important properties individually which are not reflected in the behavior of R as a ring of operators in \mathfrak{S} .

IV. Remark. Since the publication of Tomita's paper, several papers have appeared on the more general problem of expressing a point in a convex-compact set K as a direct integral over the set N of extremal points of K . Choquet [2], [3], Bishop and deLeeuw [1], and Loomis [4] have all obtained results on this problem. Referring to Loomis's paper, we note that these results can be applied to the Tomita problem when K is the set of normalized positive functionals on a ring R which contains its commutant E . The results of Loomis yield a unique measure μ on K having the properties (1) $F(A) = \int_K f(A) d\mu(f)$, and (2) μ is zero on any G_δ set which misses N . The carrier of μ is homeomorphic to M ; however, Theorem 3.3 shows that the carrier of μ may be disjoint from N in spite of (2).

It is possible to use the measure μ to define a measure ω on the set algebra consisting of all sets of the form $N \cap \Delta$ where Δ is a Baire set of K . With a slight modification of the definition of direct integrals, ω yields a direct integral decomposition of R into irreducible rings R_f . Unfortunately ω is not a regular Borel measure on a compact space, so that some of the important properties of a direct integral are lost.

BIBLIOGRAPHY

1. E. Bishop and K. deLeeuw, *The representation of linear functionals by measures on sets of extreme points*, Ann. Inst. Fourier (Grenoble) **9** (1959), 305-331.
2. G. Choquet, *Unicité des représentations intégrales au moyen de points extrémaux dans les cônes convexes réticulés*, C. R. Acad. Sci. Paris **243** (1956), 555-557.
3. ———, *Existence des représentations intégrales au moyen de points extrémaux dans les cônes convexes*, C. R. Acad. Sci. Paris. **243** (1956), 699-702.
4. L. H. Loomis, *Unique direct integral decomposition on convex sets*, Amer. J. Math. **84** (1962), 509-526.
5. M. A. Naimark, *Normed rings*, Noordoff, Groningen, 1959.
6. M. Tomita, *Representations of operator algebras*, Math. J. Okayama Univ. **3** (1954), 147-173.

LOUISIANA STATE UNIVERSITY,
BATON ROUGE, LOUISIANA