

DUAL SPACES OF A VECTOR LATTICE AND ITS CUT-COMPLETION

BY
J. J. MASTERSON

Introduction. Let L be an Archimedean vector lattice. The space L_n^\sim of all normal integrals (order continuous linear functionals) on L was first studied by H. Nakano [6], and has been the object of recent investigation [1]–[5].

The central theorem in this paper shows that if \hat{L} is the cut-completion of L , then L_n^\sim and $(\hat{L})_n^\sim$ are isomorphic vector lattices, a known result when $L = C(X)$, the continuous functions on a compact Hausdorff space (cf. [3, §10]). This serves as a convenient tool for examining the role that completeness of L plays in determining properties of L_n^\sim . It will be shown that a number of results previously obtained under the assumption that L is complete can be extended to Archimedean vector lattices.

1. **The isomorphism of L_n^\sim and $(\hat{L})_n^\sim$.** We first recall some definitions and other facts basic to our study of the cut-completion (or Dedekind completion) of a vector lattice. We will say that a subvector lattice D of L is *order dense* in \hat{L} (or *dense* in L) if for every positive element a in L , $a = \inf(B)$ for some set B of positive elements taken from D . \hat{L} then is said to be a *cut-completion* of the vector lattice L if \hat{L} is a complete vector lattice which contains a dense subvector lattice isomorphic to L ("isomorphism" always denotes vector lattice isomorphism). To simplify notation we will consider L to be a subvector lattice of \hat{L} .

The following three properties of \hat{L} are well known and will be used in the paper without proof:

- (a) \hat{L} is unique up to isomorphism.
- (b) If B is an arbitrary subset of elements of L and the supremum of B taken in L exists, then it is identical with the supremum of B taken in \hat{L} .
- (c) If \hat{a} is a positive element of \hat{L} , then there exist subsets B and C of L such that $\hat{a} = \sup(B) = \inf(C)$.

Let L be an Archimedean vector lattice and \hat{L} its cut-completion. Then,

THEOREM 1.1. L_n^\sim and $(\hat{L})_n^\sim$ are isomorphic vector lattices⁽¹⁾.

Proof. Let $f \geq 0$ be a normal integral on L and extend it to the positive cone \hat{L}^+ of \hat{L} by

$$\hat{f}(\hat{a}) = \sup \{ f(x) : 0 \leq x \leq \hat{a}, x \in L \}, \quad \hat{a} \in \hat{L}.$$

Received by the editors March 1, 1966 and, in revised form, October 26, 1966.

⁽¹⁾ The author wishes to thank the referee for simplifying the proof of Theorem 1.1.

The supremum of a set in L exists and agrees with that in \hat{L} from which it follows that $\hat{f}|_{L^+} = f$. Moreover, \hat{f} is monotone on \hat{L}^+ .

(a) $\hat{f}(\hat{a}) = \inf \{f(y) : \hat{a} \leq y \in L\}$. Clearly, $\hat{f}(\hat{a}) \leq \inf \{f(y) : \hat{a} \leq y \in L\}$. Suppose there is an $\epsilon > 0$ such that $\hat{f}(\hat{a}) + \epsilon \leq \inf \{f(y) : \hat{a} \leq y \in L\}$. Then,

$$\inf_{\hat{L}} \{y - x : 0 \leq x \leq \hat{a} \leq y; x, y \in L\} = 0 = \inf_{L} \{y - x : 0 \leq x \leq \hat{a} \leq y; x, y \in L\}.$$

But f would not then be order continuous, contradicting the fact that it is a normal integral.

(b) If $B \subset \hat{L}^+$, $\hat{f}(\inf B) = \inf(\hat{f}(B))$. Again, $\hat{f}(\inf B) \leq \inf(\hat{f}(B))$ is clear. Let $\epsilon > 0$ and $\inf B = \hat{b}_0$. There exists $x \in L$, $x \geq \hat{b}_0$ with $\hat{f}(\hat{b}_0) + \epsilon \geq f(x)$. Then by the monotonicity of f there is $\hat{b} \in B$ with $\hat{b} \leq x$, whence $\hat{f}(\hat{b}_0) + \epsilon \geq \hat{f}(\hat{b}) \geq \inf(\hat{f}(B))$, then, is order continuous on \hat{L} .

(c) \hat{f} is additive on \hat{L}^+ . Let \hat{a}_1 and \hat{a}_2 be in \hat{L} .

$$\{[0, \hat{a}_1] \cap L\} \cup \{[0, \hat{a}_2] \cap L\} \subset [0, \hat{a}_1 + \hat{a}_2] \cap L.$$

So,

$$\begin{aligned} \hat{f}(\hat{a}_1) + \hat{f}(\hat{a}_2) &= \sup(f([0, \hat{a}_1] \cap L)) + \sup(f([0, \hat{a}_2] \cap L)) \\ &= \sup(f(\{[0, \hat{a}_1] \cap L\} \cup \{[0, \hat{a}_2] \cap L\})) \\ &\leq \sup(f([0, \hat{a}_1 + \hat{a}_2] \cap L)) \\ &= \hat{f}(\hat{a}_1 + \hat{a}_2). \end{aligned}$$

Let $\epsilon > 0$. Then, there exists $b \in [0, \hat{a}_1 + \hat{a}_2] \cap L$ with $\hat{f}(\hat{a}_1 + \hat{a}_2) - \epsilon \leq f(b)$. Let $b = \hat{b}_1 + \hat{b}_2$. Then, $\hat{f}(\hat{b}_2) = \hat{f}(b - \hat{b}_1) = f(\inf \{b - x : 0 \leq x \leq \hat{b}_1, x \in E\}) = f(b) - \hat{f}(\hat{b}_1)$ applying the decomposition lemma for vector lattices. So, $\hat{f}(\hat{a}_1 + \hat{a}_2) - \epsilon \leq f(b) = \hat{f}(\hat{b}_1) + \hat{f}(\hat{b}_2) \leq \hat{f}(\hat{a}_1) + \hat{f}(\hat{a}_2)$. Equality then follows.

(d) Positive homogeneity is clear. Setting $\hat{f}(\hat{x}) = \hat{f}(\hat{x}^+) - \hat{f}(\hat{x}^-)$ for arbitrary $\hat{x} \in \hat{L}$, then, \hat{f} becomes a normal integral on \hat{L} . \hat{f} is uniquely determined by f and it follows rather easily that the mapping $f \rightarrow \hat{f}$ is the desired isomorphism.

If $L_{\tilde{n}}$ is separating on L , then L can be embedded in $L_{\tilde{n}} \sim (L_{\tilde{n}} \sim = (L_{\tilde{n}})_{\tilde{n}})$ as a subvector lattice. If L is complete, then under this embedding it is an ideal in $L_{\tilde{n}} \sim$ (cf. [1, 3.8]). So, \hat{L} is an ideal in $(\hat{L})_{\tilde{n}} \sim$. With the above theorem, this gives us the following:

COROLLARY 1.2. \hat{L} can be embedded in $L_{\tilde{n}} \sim$ as an ideal.

Moreover, it is easily seen that \hat{L} is the ideal in $L_{\tilde{n}} \sim$ generated by L , a somewhat surprising relation between the concepts of cut-completion and continuous linear functional.

We will now apply the theorem to obtain a result concerning absolute weak topologies on $L_{\tilde{n}}$. The absolute weak topology on $L_{\tilde{n}}$ generated by L is the locally-convex topology defined in $L_{\tilde{n}}$ by the polars of the bounded sets in L . Equivalently, if for each positive element a in L , we define a seminorm $\| \cdot \|_a$ on $L_{\tilde{n}}$ by

$\|f\|_a = |f|(a)$, the absolute weak topology is the topology defined by the family of all such seminorms. We will denote this topology by $|w|(L_{\tilde{n}}, L)$.

Using the seminorm definition of the topology and the fact that for every positive element \hat{a} in \hat{L} there is a positive element a in L such that $a \geq \hat{a}$, and for every positive element b in L there is a positive element \hat{b} in \hat{L} such that $\hat{b} \geq b$, we obtain the following:

COROLLARY 1.3. *The $|w|(L_{\tilde{n}}, L)$ and $|w|(L_{\tilde{n}}, \hat{L})$ topologies on $L_{\tilde{n}}$ are identical.*

The following two results remove the hypothesis that L be a complete vector lattice from theorems appearing in [2] and [3]; ([3, 2.8] and [2, 3.3] respectively).

COROLLARY 1.4. *If I is a dense ideal in L , then $L_{\tilde{n}}$ can be embedded in $I_{\tilde{n}}$ as a dense ideal.*

Proof. The cut-completion of I has a natural embedding in \hat{L} as a dense ideal. We call this ideal \hat{I} . By Theorem 1.1, $I_{\tilde{n}} \cong (\hat{I})_{\tilde{n}}$ and $L_{\tilde{n}} \cong (\hat{L})_{\tilde{n}}$. Applying the analogous theorem in [3] to \hat{I} and \hat{L} , the result follows.

A subvector lattice D of L is said to be *closed* in L if whenever $B \subset D$ is a set of positive elements such that $\sup(B)$ is in L , then $\sup(B)$ is in D .

COROLLARY 1.5. *If I is a closed ideal in $L_{\tilde{n}}$, then I is $w(L_{\tilde{n}}, L)$ closed.*

Proof. (The proof is a variation of that in [2], letting J be the null ideal of ϕ in \hat{L} instead of L .) $L_{\tilde{n}} = I \oplus I'$ since I is closed. It suffices to show that if $\phi \in I'$ and $\phi \neq 0$, then ϕ is not a $w(L_{\tilde{n}}, L)$ limit point of I . Now, the null ideal J of ϕ in \hat{L} is closed ([2], 3.3) hence $\hat{L} = J \oplus J'$. So, $L_{\tilde{n}} = J^{\perp} \oplus J'^{\perp}$. Let us first show that J'^{\perp} is $w(L_{\tilde{n}}, L)$ closed. Let $\{f_{\alpha}\} \subset J'^{\perp}$ and $f \in L_{\tilde{n}}$ be such that $f_{\alpha} \rightarrow f(w(L_{\tilde{n}}, L))$ i.e., $f_{\alpha}(a) \rightarrow f(a)$ for all $a \in L$. But then for $a \in J' \cap L$, $f_{\alpha}(a) = 0$ for all α , since $f_{\alpha} \in J'^{\perp}$. So, $f(a) = 0$ for each $a \in J' \cap L$. Hence for $\hat{a} \in J'$, $f(\hat{a}) = 0$ (f being continuous and L being order dense in \hat{L}). So, $f \in J'^{\perp}$ and hence J'^{\perp} is $w(L_{\tilde{n}}, L)$ closed. Now $\phi \in J^{\perp}$ hence not in J'^{\perp} . So, we can show that ϕ is not a weak limit point of I by showing that $I \subset J'^{\perp}$ (J'^{\perp} being weakly closed). Since $L_{\tilde{n}} = J^{\perp} \oplus J'^{\perp}$ we establish $I \subset J'^{\perp}$ by proving that $I \cap J^{\perp} = \{0\}$.

Consider any $\psi \in I$, $\psi > 0$. We have to show $\psi(a) \neq 0$ for some $a \in J$. Now, $\psi \wedge |\phi| = 0$. So, for $\epsilon > 0$ and $b \in \hat{L}^+$, there is c in \hat{L} such that $0 \leq c \leq b$ and $|\phi|(c) + \psi(b - c) \leq \epsilon$. It follows easily that there is a sequence $c_1 \geq c_2 \geq \dots \geq 0$ in \hat{L} satisfying $|\phi|(c_n) \leq 1/2^n$ and $\psi(c_n) \geq \lambda > 0$ for all n . Let $a = \bigwedge_n c_n$. Then, $|\phi|(a) = 0$ which implies that $a \in J$ and since ψ is continuous, $\psi(a) > 0$.

REMARK. The proof of 1.5 would be trivial if the topologies $w(L_{\tilde{n}}, L)$ and $w(L_{\tilde{n}}, \hat{L})$ were identical, as is the case with the $|w|$ -topologies. This is not true, however, as the following example shows: Let $L = c$ (convergent sequences). Then $L_{\tilde{n}} = l^1$ and $\hat{L} = l^{\infty}$. Define $x_n = (x_n^{(1)}, x_n^{(2)}, \dots)$ by: $x_n^{(k)} = 0, k \neq n, n + 1, x_n^{(n)} = 1, x_n^{(n+1)} = -1$. Then $\{x_n\}$ is a sequence in l^1 which converges to 0 in the $w(l^1, c)$ topology but not in the $w(l^1, l^{\infty})$ topology.

2. **The space of normal integrals.** In [5], Zaanen and Luxemburg extend a number of the results of [4] from complete to Archimedean vector lattices. New (and often simpler) proofs of these results will be given based on Theorem 1.1 and the elementary propositions listed below. Moreover, natural extensions to the Archimedean case are obtained for several other theorems in [4].

We will first recall some definitions from [4] and state some elementary facts that we need.

If $\phi \in L_n^\sim$, then $N_\phi = \{a \in L : |\phi| |a| = 0\}$ is called the *null ideal* of ϕ in L . We denote the null ideal of ϕ in \hat{L} by \hat{N}_ϕ . The *carrier* of ϕ in L is the complement of N_ϕ in L . I.e., $C_\phi = (N_\phi)^p = \{b \in L : |a| \wedge |b| = 0 \text{ for all } a \in N_\phi\}$. \hat{C}_ϕ is the carrier of ϕ in \hat{L} . If $A \subset L$, the *annihilator* of A in L_n^\sim is the set $\{\phi \in L_n^\sim : \phi(a) = 0 \text{ for all } a \in A\}$ and will be denoted by $A^\#$. If $B \subset L_n^\sim$, ${}^0B = \{a \in L : \phi(a) = 0 \text{ for all } \phi \in B\}$ is called the *annihilator of B in L*. ${}^{\hat{0}}B$ denotes the annihilator of B in \hat{L} .

Theorem 1.1 tells us, then, that a normal integral on L may be regarded as a normal integral on \hat{L} and *vice versa*. We will use this fact without mention in proving the following propositions.

From the definition of N_ϕ and the above-mentioned fact we obtain immediately

PROPOSITION 2.1. *If $\phi \in L_n^\sim$, then $N_\phi = \hat{N}_\phi \cap L$.*

PROPOSITION 2.2. *If \hat{I} is an ideal in \hat{L} and $I = \hat{I} \cap L$, then $(\hat{I})^p \cap L = I^p$.*

Proof. I is dense in \hat{I} . Then, if $a \in I^p$, $|a| \wedge |b| = 0$ for all $b \in I$. Given $\hat{b} \in \hat{I}$, there is a net $\{b_\alpha\} \subset I$ such that $|b_\alpha| \uparrow |\hat{b}|$. But $|a| \wedge |b_\alpha| = 0$ for all α . Hence $|a| \wedge |\hat{b}| = 0$ which implies that $a \in (\hat{I})^p \cap L$, hence $I^p \subset (\hat{I})^p \cap L$.

Let $a \in (\hat{I})^p \cap L$. Then $|a| \wedge |\hat{b}| = 0$ for all $\hat{b} \in \hat{I}$ implies $|a| \wedge |b| = 0$ for all $b \in I$ since $I \subset \hat{I}$. So, $a \in I^p$ and hence $(\hat{I})^p \cap L \subset I^p$. Thus, $(\hat{I})^p \cap L = I^p$.

Since $C_\phi = (N_\phi)^p$ by definition, the following is an immediate consequence of 2.1 and 2.2.

COROLLARY 2.3. *If $\phi \in L_n^\sim$, $C_\phi = \hat{C}_\phi \cap L$.*

PROPOSITION 2.4. *If \hat{I} is an ideal in L and $I = \hat{I} \cap L$, then for $\phi \in L_n^\sim$, $\phi(I) = 0$ if and only if $\phi(\hat{I}) = 0$, i.e., $(\hat{I})^\# = I^\#$.*

Proof. Since $I \subset \hat{I}$, $\phi(\hat{I}) = 0$ implies $\phi(I) = 0$. Suppose $\phi(I) = 0$. Then, if $\hat{a} \in \hat{I}^+$, there is a net $\{a_\alpha\}$ in I such that $a_\alpha \uparrow \hat{a}$. Since ϕ is continuous $\phi(a_\alpha) \rightarrow \phi(\hat{a})$. Hence $\phi(\hat{a}) = 0$. So, $\phi(\hat{I}) = 0$.

By definition, 0B is precisely those elements of $\hat{0}B$ which lie in L . We state more precisely

PROPOSITION 2.5. *If B is a subset of L_n^\sim , then $({}^{\hat{0}}B) \cap L = {}^0B$.*

PROPOSITION 2.6. *If $\hat{L} = \hat{I} \oplus \hat{J}$, $I = \hat{I} \cap L$ and $J = \hat{J} \cap L$, then $L = \{I \oplus J\}$.*

Proof. By [5, Theorem 29.10], $L = \{I \oplus I^p\}$. Since $\hat{L} = \hat{I} \oplus \hat{J}$, $\hat{J} = (\hat{I})^p$. Hence $J = \hat{J} \cap L = (\hat{I})^p \cap L = I^p$ by Proposition 2.2. So $L = \{I \oplus J\}$.

The following theorems are proved in [5]. Proofs will be given here based on Theorem 1.1 and the results for Dedekind complete vector lattices in [4]. We denote the set of normal integrals on both L and \hat{L} by L_n^\sim .

THEOREM 2.8. *If L is Archimedean and if $\phi, \Psi \in L_n^\sim$, then the following conditions are mutually equivalent: (a) $|\phi| \wedge |\Psi| = 0$, (b) $C_\phi \subset N_\Psi$, (c) $C_\Psi \subset N_\phi$, (d) $C_\phi \perp C_\Psi$ (cf. [5, 31.2]).*

Proof. By [4, Theorem 27.6], the following are equivalent: (a') $|\phi| \wedge |\psi| = 0$, (b') $\hat{C}_\phi \subset \hat{N}_\psi$, (c') $\hat{C}_\psi \subset \hat{N}_\phi$, (d') $\hat{C}_\phi \perp \hat{C}_\psi$. To complete the argument, it suffices to show that (b) is equivalent to (b') and (d) is equivalent to (d'). By 2.1 and 2.3, (b') implies (b). Assume (b) i.e., $C_\phi \subset N_\psi$. Let $a \in \hat{C}^+$. Then, there is a net $\{a_\alpha\} \subset C_\phi$ such that $a_\alpha \uparrow a$. But then $\{a_\alpha\} \subset N_\psi$. So $|\psi|(|a_\alpha|) = 0$ for all α and hence $|\psi|(|a|) = 0$. Thus, $a \in \hat{N}_\psi$ hence $\hat{C}_\phi \subset \hat{N}_\psi$. Using the density of C_ϕ in \hat{C}_ϕ and 2.3, the equivalence of (d) and (d') follows easily.

COROLLARY 2.9 (HAHN DECOMPOSITION). *Let L be Archimedean. Given $\phi \in L_n^\sim$, there exist normal subspaces P and N such that $P \cap N = \{0\}$, $P \oplus N$ is (order) dense in L , $\phi(u) \leq 0$ for all $0 \leq u \in N$ and $\phi(u) \geq 0$, for $0 \leq u \in P$. Moreover, $P = C_{\phi^+}$ and $N = N_{\phi^+}$ (cf. [5, 31.3]).*

Proof. By [4, Corollary 27.7], there exist \hat{P} and \hat{N} such that $\hat{L} = \hat{P} \oplus \hat{N}$ (\hat{P} and \hat{N} are normal subspaces) such that $\phi(u) \geq 0$ for all $0 \leq u \in \hat{P}$ and $\phi(u) \leq 0$ for all $0 \leq u \in \hat{N}$. Moreover, $\hat{P} = \hat{C}_{\phi^+}$ and $\hat{N} = \hat{N}_{\phi^+}$. Let $P = \hat{P} \cap L$ and $N = \hat{N} \cap L$. Then, by 2.1 and 2.3, $N = N_{\phi^+}$ and $P = C_{\phi^+}$. $\hat{P} \cap \hat{N} = \{0\}$ hence $P \cap N = \{0\}$. Also, since $P \subset \hat{P}$ and $N \subset \hat{N}$, $\phi(u) \leq 0$, $0 \leq u \in N$ and $\phi(u) \geq 0$, $0 \leq u \in P$. By 2.6, $\{P \oplus N\} = L$. Since \hat{P} is a normal subspace of \hat{L} , $P = \hat{P} \cap L$ is a normal subspace of L and likewise for N . This completes the argument.

Theorem 31.5 in [5] can be generalized without using the lemma which precedes it.

THEOREM 2.10. *Let L be Archimedean. If B is an ideal in L_n^\sim , then $({}^0B)^\# = \{B\}$.*

Proof. Since $\{B\}$ is a normal subspace, it suffices to consider B to be a normal subspace of L_n^\sim . Then, by [4, Theorem 27.10], $({}^0B)^\# = B = \{B\}$. But $({}^0B)^\# = ({}^0B \cap L)^\# = ({}^0B)^\#$ by 2.5 and 2.4. Applying [4, Theorem 27.10], we have $({}^0B)^\# = ({}^0B)^\# = \{B\}$.

If B is an ideal in L_n^\sim , then the normal subspace of L generated by all carriers C_ϕ , where $\phi \in B$, is called the *carrier* of B and is denoted by C_B .

Let \hat{C}_B denote the carrier of B in \hat{L} . Then, we note that $\hat{C}_B \cap L = C_B$. Clearly, $C_B \subset \hat{C}_B \cap L$ since for each $\phi \in B$, $C_\phi \subset \hat{C}_\phi \cap L$. Now suppose $a \in \hat{C}_B \cap L$ is a positive element. Then, there is a net $\{\hat{a}_\alpha\}$ such that for each α , $\hat{a}_\alpha \in \hat{C}_\phi$ for some $\phi \in B$, and such that $\hat{a}_\alpha \uparrow a$. For each α , there is a net $\{a_{\alpha,\beta}\} \subset L$ such that $a_{\alpha,\beta} \uparrow \hat{a}_\alpha$. Then, $a = \vee a_{\alpha,\beta} - in - \hat{L}$ and hence $a = \vee a_{\alpha,\beta} - in - L$. But $a_{\alpha,\beta} \in C_\phi$ for some ϕ , given α and β . Hence $a \in C_B$.

THEOREM 2.11. (i) *If L is Archimedean and B is an ideal in $L_{\tilde{n}}$, then $C_B = ({}^0B)^p$. Hence, $L = \{{}^0B \oplus C_B\}$. In addition, the ideals B and B_1 in $L_{\tilde{n}}$ are disjoint if and only if C_B and C_{B_1} are disjoint.*

(ii) *If L is Archimedean and B is an ideal in $L_{\tilde{n}}$, then $C_B = C_{(B)}$ (cf. [5, 31.8]).*

Proof. By [4, Theorem 27.12], $\hat{C}_B = ({}^0B)^p$. By the remark made above, $C_B = \hat{C}_B \cap L$. But then, $C_B = \hat{C}_B \cap L = ({}^0B)^p \cap L = ({}^0B)^p$ by 2.2 and 2.5. Using again the remark made above, it is easily shown that $C_B \perp C_{B_1}$ if and only if $\hat{C}_B \perp \hat{C}_{B_1}$. But it is proved in [4, Theorem 27.12] that ideals B and B_1 in $L_{\tilde{n}}$ are disjoint if and only if \hat{C}_{B_1} and \hat{C}_B are disjoint. The last statement in (i) then follows.

Since $C_B = \hat{C}_B \cap L$, and from [4, Theorem 27.12], $\hat{C}_B = \hat{C}_{(B)}$, (ii) is proved.

$L_{\tilde{n}}$ is a normal subspace in L^{\sim} . Then, let $L_0 = {}^0(L_{\tilde{n}})$ and let L_1 be the carrier of $L_{\tilde{n}}$ in L . Then, $L_1 = (L_0)^p$ and $L = \{L_0 \oplus L_1\}$.

THEOREM 2.12. (i) *If L is Archimedean and A is a normal subspace of L_1 , then ${}^0(A^{\#}) = \{A \oplus L_0\}$. (ii) *If L is Archimedean and A is a normal subspace of L_1 , then A is the carrier of some normal subspace of $L_{\tilde{n}}$ (cf. [5, Theorem 31.10]).**

Proof. (i) If A is a normal subspace of L_1 , then $\hat{A} = \{b \in \hat{L} : |b| \leq a \text{ for some } a \in L\}$ is a normal subspace of \hat{L}_1 . So, by [4, Theorem 27.14], ${}^0(\hat{A}^{\#}) = \hat{A} \oplus \hat{L}_0$. Now ${}^0(\hat{A}^{\#}) \cap L = {}^0(A^{\#})$ by 2.5 and 2.4. By 2.6, $(\hat{A} \oplus \hat{L}_0) \cap L = \{A \oplus L_0\}$; hence (i) follows.

(ii) If A is a normal subspace of L_1 , then \hat{A} is a normal subspace of L_1 . But then, \hat{A} is by [4, Theorem 27.14], the carrier of some normal subspace \hat{B} of $L_{\tilde{n}}$. So, by 2.2 and 2.5, and the fact that $A = \hat{A} \cap L$, A is the carrier of \hat{B} in L .

It is also possible, using 1.1, to prove Archimedean analogs of other results proved in [4] for Dedekind complete vector lattices.

Recall that a vector lattice L has the *Egoroff property* in case $0 \leq u_{n,k} \uparrow u$, $n = 1, 2, \dots$ implies the existence of a sequence $0 \leq v_m \uparrow u$ such that for every pair of indices (m, n) there exists $j = j(m, n)$ satisfying $v_m \leq u_{n,j}$. The concept “super-Dedekind complete” can be extended in a natural way to an Archimedean vector lattice L by omitting the condition that L be Dedekind complete: An Archimedean vector lattice is called *super-Archimedean* in case $0 \leq u_{\tau}$ and $u_{\tau} \uparrow u_0$ implies the existence of a subsequence $\{u_{\tau_n}\}$ of u_{τ} such that $u_{\tau_n} \uparrow u_0$. Then [4, Theorem 27.17] can be extended using the following lemma.

LEMMA 2.13. *If $\hat{I} \subset \hat{L}$ is an ideal and I is super-Dedekind complete and has the Egoroff property, then $I = \hat{I} \cap L$ is super-Archimedean and has the Egoroff property.*

Proof. I is clearly super-Archimedean since it inherits this property from \hat{I} . Suppose $0 \leq u_{n,k} \uparrow u$ for each $n = 1, 2, \dots$; $u \in I$, $u_{n,k} \in I$, $n, k = 1, 2, \dots$. Then, there exists a sequence $\{\hat{v}_m\} \subset \hat{I}$ such that $\hat{v}_m \uparrow u$ and such that for every pair of indices m, n , there exists $j = j(m, n)$ such that $\hat{v}_m \leq u_{n,j}$. Now, there is a net $\{v_{m,\tau}\} \subset I$ such that $0 \leq v_{m,\tau} \uparrow \hat{v}_m$ for each m and hence, since \hat{I} is super-Dedekind complete, a sequence $\{v_{m,k}\} \subset I$ such that $v_{m,k} \uparrow \hat{v}_m$ for each m . Extend the countable set $\{v_{m,k}, k = 1, 2, \dots$; $m = 1, 2, \dots\}$ to a countable directed set $\{w_v\}$ by adding all suprema of finite

subsets of $\{v_{m,k}\}$. $\{w_\nu\}$ is then a sequence of elements of I such that $w_\nu \uparrow u$ (since $v_{m,k} \uparrow \hat{v}_m$ for each m and $\hat{v}_m \uparrow u$). Moreover, given ν , $w_\nu = v_{m_1,k_1} \vee \cdots \vee v_{m_p,k_p}$ and $v_{m_1,k_1} \leq \hat{v}_{m_1}, \dots, v_{m_p,k_p} \leq \hat{v}_{m_p}$. But then, since $\{\hat{v}_m\}$ is directed there is an m' such that $\hat{v}_{m_1} \vee \cdots \vee \hat{v}_{m_p} \leq \hat{v}_{m'}$ so, $w_\nu \leq \hat{v}_{m'}$. Then, if ν and n are given, there is j , namely $j = j(m', n)$ such that $\hat{v}_{m'} \leq u_{n,j}$ and hence $w_\nu \leq u_{n,j}$. So, I has the Egoroff property.

We recall from [4] the definition of *Riesz direct sum*. $A \subset L$ will be called the Riesz direct sum of the collection of normal subspaces $\{A_\tau\}$ if all A_τ are mutually disjoint and if A is the smallest normal subspace containing all A_τ . We use the notation $A = \bigcup \oplus A_\tau$.

LEMMA 2.14. *If $\hat{A} \subset \hat{L}$ and $\hat{A} = \bigcup \oplus \hat{A}_\tau$, then $A = \bigcup \oplus A_\tau$ where $A = \hat{A} \cap L$ and $A_\tau = \hat{A}_\tau \cap L$.*

Proof. Let $B = \bigcup \oplus A_\tau$. Then B is a normal subspace of L and $B \subset A$. Suppose $B \neq A$. Then, since B and A are both normal subspaces of L , B is not dense in A . Let $B_A^p = B^p \cap A$. Then by [5, Theorem 29.10] $B^p \neq \{0\}$. Let $\hat{B}^p = \{b \in \hat{L} : |b| \leq c \text{ for some } c \in B^p\}$. Then, $\{\hat{B}_A^p\}$ is a normal subspace in \hat{L} , and $\{\hat{B}_A^p\} \neq \{0\}$. Moreover, $A_\tau \perp B_A^p$ for each τ and have $\hat{A}_\tau \perp \hat{B}_A^p$ for each τ . But then since \hat{A}_τ is a normal subspace of \hat{L} , $\hat{A}_\tau \perp \{\hat{B}_A^p\}$ for each τ . So, since $\{\hat{B}_A^p\} \neq \{0\}$, we have that $\hat{A} \cap \{\hat{B}_A^p\}^p$ is a normal subspace of \hat{L} which contains \hat{A}_τ for each τ and yet, \hat{A} contains it properly. This contradicts the hypothesis.

THEOREM 2.15. *Let L be Archimedean. The carrier L_1 of L_n^\sim is the Riesz direct sum of normal subspaces A_τ such that A_τ is super-Archimedean and has the Egoroff property, and in addition, each A_τ possesses a strictly positive normal integral.*

Proof. By [4, Theorem 27.17], $\hat{L}_1 = \bigcup \oplus \hat{A}_\tau$ where \hat{A}_τ is a normal subspace of \hat{L} for each τ . Then $L_1 = \hat{L}_1 \cap L = \bigcup \oplus A_\tau$ where $A_\tau = \hat{A}_\tau \cap L$ by 2.14. But A_τ is a normal subspace of L and possesses the required properties by 2.13.

The corollary to Theorem 27.17 in [4], can be generalized independently of the theorem.

THEOREM 2.16. *If L_n^\sim has a finite or countable order basis, then the carrier L_1 of L_n^\sim is super-Archimedean.*

Proof. By Corollary 27.18 in [4], \hat{L}_1 is super-Dedekind complete and hence by 5.1, L_1 is super-Archimedean, noting that $L_1 = \hat{L}_1 \cap L$.

Corollary 27.19 in [4] generalizes quite easily to Archimedean vector lattices, using 1.1:

THEOREM 2.17. (i) *L_n^\sim is the Riesz direct sum of normal subspaces B_τ such that each B_τ is super-Dedekind complete and has the Egoroff property.*

(ii) *If the carrier L_1 of L_n^\sim has a finite or countable order basis, then L_n^\sim is super-Dedekind complete.*

Proof. (i) is an immediate consequence of 1.1. If \hat{L}_1 has a countable order basis

then L_1 has a countable order basis since L_1 is order-dense in \hat{L}_1 . (ii) then follows from 1.1 and [4, Corollary 27.19].

Added in proof. In a subsequent paper, the authors of [4] give a proof of Theorem 1.1. They apply it to different problems.

REFERENCES

1. S. Kaplan, *The second dual of the space of continuous functions*. II, Trans. Amer. Math. Soc. **93** (1959), 329–350.
2. ———, *The second dual of the space of continuous functions*. III, Trans. Amer. Math. Soc. **101** (1961), 34–51.
3. ———, *The second dual of the space of continuous functions*. IV, Trans. Amer. Math. Soc. **113** (1964), 512–546.
4. W. A. J. Luxemburg and A. C. Zaanen, *Notes on Banach function spaces*. VIII, Nederl. Akad. Wetensch. Proc. Ser. A **67** (1964), 104–119.
5. ———, *Notes on Banach function spaces*. IX, Nederl. Akad. Wetensch. Proc. Ser. A **67** (1964), 360–376.
6. H. Nakano, *Modern spectral theory*, Maruzen, Tokyo, 1950.

PURDUE UNIVERSITY,
LAFAYETTE, INDIANA